

# On Einstein metrics, normalized Ricci flow and smooth structures on $3\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$

Rafael Torres

ABSTRACT. In this paper, first we consider the existence and nonexistence of Einstein metrics on the topological 4-manifolds  $3\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$ , the connected sum of  $\mathbb{C}\mathbb{P}^2$  with both choices of orientation, by using the idea of Răşdeaconu–Şuvaina, 2009, and the constructions in Park–Park–Shin, 2013. Then, we study the existence or nonexistence of nonsingular solutions of the normalized Ricci flow on the exotic smooth structures of these topological manifolds by employing the obstruction developed in Ishida, 2008.

## CONTENTS

1. Introduction	179
2. Homeomorphism type	181
2.1. Description of the minimal surfaces of general type built by H. Park, J. Park and D. Shin	182
3. Existence of Einstein metrics	183
3.1. Proof of Proposition 8	183
4. Nonexistence of Einstein metrics: Exotic smooth structures	185
4.1. Nonexistence of anti-self-dual Einstein metrics	186
5. Proof of Proposition 2	186
References	186

## 1. Introduction

Recent years have witnessed a drastic increase in our understanding of the topology and geometry of 4-manifolds and complex surfaces. The newest developments can be exemplified by the construction of simply connected surfaces of general type with small topology [22, 23], by the unveiling of a myriad of exotic smooth structures on small 4-manifolds [1], and by how these manifolds have provided an adequate environment for the study of

Received May 9, 2013.

2010 *Mathematics Subject Classification*. Primary 53C25; Secondary 57R55.

*Key words and phrases*.  $\mathbb{Q}$ -Gorenstein smoothings, surfaces of general type, exotic smooth structures, Einstein metrics.

fundamental questions in Riemannian geometry that were previously out of reach.

In particular, intriguing questions regarding Einstein metrics ([17, 24]), and the relation between smooth and geometric structures (like the Yamabe invariant and the normalized Ricci flow) on a given topological 4-manifold ([17, 13, 15]) have been immediate beneficiaries of the novel constructions. In this paper we employ the procedure of R. Răşdeaconu and I. Şuvaina ([24]) to the constructions of H. Park, J. Park and D. Shin ([23]), and to those of A. Akhmedov and B.D. Park ([1]) to study the (non)-existence of Einstein metrics, and the (non)-existence of nonsingular solutions to the normalized Ricci flow on manifolds with small topological invariants (although bigger than those considered in [24]).

Our main results are the following.

**Theorem 1.** *Let  $k \in \{11, 13, 14, 15, 16, 17, 18\}$ . Each of the topological 4-manifolds*

$$3\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$$

*admits a smooth structure that has an Einstein metric of scalar curvature  $s < 0$ , and infinitely many nondiffeomorphic smooth structures that do not admit Einstein metrics.*

Regarding the nonsingular solutions to the normalized Ricci flow on the exotic smooth structures of the manifolds from Theorem 1 and in the spirit of [15], the following result is proven.

**Proposition 2.** *Let  $k \in \{11, 13, 14, 15, 16, 17, 18\}$ . The topological 4-manifold  $M := 3\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$  satisfies the following properties:*

- (1)  *$M$  admits a smooth structure of negative Yamabe invariant on which there exist nonsingular solutions to the normalized Ricci flow.*
- (2)  *$M$  admits infinitely many smooth structures, all of which have negative Yamabe invariant, and on which there are no nonsingular solutions to the normalized Ricci flow for any initial metric.*

We are also able to prove that for  $k \geq 9$ , each of the spaces of Theorem 1 have infinitely many reducible smooth structures that do not carry an Einstein metric, all of which have negative Yamabe invariant, and on which the only solutions to the normalized Ricci flow for any initial metric are all singular (Proposition 11). Moreover, for  $k \geq 8$ , the manifolds of our theorem do not admit anti-self-dual Einstein metrics (Lemma 12). Theorem 1 and Proposition 2 extend the results in [24] and [15], and improve results of [14] and [5].

The (non)-existence of Einstein metrics on different smooth structures on  $3\mathbb{C}\mathbb{P}^2$  blown-up at a small number of points was previously considered by V. Braungart and D. Kotschick in [5]. In that paper, the authors proved instances  $k = 17$  and  $k = 18$  of Theorem 1. By a result of F. Catanese [8],

in the case  $k = 18$ , the manifolds with Kähler–Einstein metrics in this paper and in [5] are diffeomorphic.

The paper is organized as follows. In Section 2 we determine the homeomorphism types of the complex surfaces built by H. Park, J. Park and D. Shin; the second part of the section provides a description of their surfaces. The third section contains the construction of an Einstein metric on each of these surfaces of general type. The nonexistence of these metrics on the topological prototypes is addressed in Section 4. The proof of Theorem 1 is spread throughout the first four sections. In Section 5, we study the sign of the Yamabe invariant and the solutions to the normalized Ricci flow on the exotic smooth structures. That is, Proposition 2 is proven in the fifth and last section.

## 2. Homeomorphism type

The following theorem was proven in [23].

**Theorem 3** (H. Park–J.Park–D. Shin). *There exist simply connected minimal surfaces of general type with  $p_g = 1$ ,  $q = 0$  and  $K^2 = 1, 2, 3, 4, 5, 6, 8$ .*

Here  $K$  denotes the canonical divisor class of the complex surface. Our enterprise starts by pinning down a homeomorphism type for each of these complex surfaces. From now on, let  $S$  be one of such surfaces. Surfaces of general type are Kähler (see, for example, [21, Lemma 2]). Thus, one has

$$b_2^+(S) = 2p_g + 1 = 3.$$

On the other hand, we have:

**Lemma 4.**  $b_2^-(S) = 19 - c_1^2(S)$ .

**Proof.** The Thom–Hirzebruch Index Theorem ([4, Theorem I 3.1, p.22]) states

$$\sigma(S) = \frac{1}{3}(c_1^2(S) - 2c_2(S)) = \frac{1}{3}(c_1^2(S) - 2e(S)).$$

The claim follows by substituting  $\sigma(S) = b_2^+ - b_2^- = 3 - b_2^-$  and  $e(S) = b_2^+ + b_2^- + 2 = 5 + b_2^-$ . □

From these computations we also observe

**Corollary 5.** *These manifolds satisfy the Hitchin–Thorpe inequality [12, Theorem 1].*

**Proof.** The claim is  $2\chi + 3\sigma > 0$ . Indeed, we have

$$2\chi + 3\sigma = 19 - b_2^-,$$

which is always positive for the manifolds considered in this paper. □

It follows from Rokhlin's Theorem [25] that the manifolds built in [23] are non-spin. We are now ready to conclude on the topological prototypes of the minimal surfaces of general type in question by using Freedman's Theorem [10], and Donaldson's results [9]. The possible homeomorphism types are arranged in the following proposition.

**Proposition 6.** *Let  $S$  be a simply connected surface of general type with  $p_g = 1$ , and  $q = 0$ . The homeomorphism type of  $S$  is given as follows:*

- If  $K_S^2 = 1$  :  $S \cong_{C^0} 3\mathbb{CP}^2 \# 18\overline{\mathbb{CP}^2}$ .
- If  $K_S^2 = 2$  :  $S \cong_{C^0} 3\mathbb{CP}^2 \# 17\overline{\mathbb{CP}^2}$ .
- If  $K_S^2 = 3$  :  $S \cong_{C^0} 3\mathbb{CP}^2 \# 16\overline{\mathbb{CP}^2}$ .
- If  $K_S^2 = 4$  :  $S \cong_{C^0} 3\mathbb{CP}^2 \# 15\overline{\mathbb{CP}^2}$ .
- If  $K_S^2 = 5$  :  $S \cong_{C^0} 3\mathbb{CP}^2 \# 14\overline{\mathbb{CP}^2}$ .
- If  $K_S^2 = 6$  :  $S \cong_{C^0} 3\mathbb{CP}^2 \# 13\overline{\mathbb{CP}^2}$ .
- If  $K_S^2 = 8$  :  $S \cong_{C^0} 3\mathbb{CP}^2 \# 11\overline{\mathbb{CP}^2}$ .

**Remark 1.** In particular notice that the 4-manifolds of H. Park, J. Park and D. Shin (Theorem 3) are exotic symplectic copies of the reducible manifolds of Proposition 6.

**2.1. Description of the minimal surfaces of general type built by H. Park, J. Park and D. Shin.** The minimal complex surfaces of Theorem 3 have a very similar nature. We proceed to give a sketch of the construction for the example with  $K^2 = 6$ . The reader is referred to the quoted papers for details.

The starting manifold is a particular rational elliptic surface  $E(1)$ , which is obtained out of blowing up a well-chosen pencil of cubics in  $\mathbb{CP}^2$ . Take the double cover of this rational elliptic surface  $E(1)$ , and call it  $Y$ . The complex surface  $Y$  is an elliptic  $K3$  surface; this complex manifold  $Y$  is a common material in all of the minimal surfaces produced in [23].

In particular for the surface with  $K^2 = 6$  we are describing in this section, one considers (within  $Y$ ) two  $I_8$ -singular fibers, two  $I_2$ -singular fibers, one nodal singular fiber, and three sections.

By blowing up  $Y$  18 times at rightly selected points (see [23, Section 4.5, Fig. 13]), one obtains a surface  $Z := Y \# 18\overline{\mathbb{CP}^2}$ . The surface  $Z$  contains five disjoint linear chains of  $\mathbb{CP}^1$ 's including the proper transforms of the sections. The linear chains are denoted by the following dual graphs, which have been labeled for the purposes in Section 3:

$$\begin{array}{c}
 \circ_{G_1}^{-2} - \circ_{G_2}^{-2} - \circ_{G_3}^{-3} - \circ_{G_4}^{-9} - \circ_{G_5}^{-2} - \circ_{G_6}^{-2} - \circ_{G_7}^{-2} - \circ_{G_8}^{-2} - \circ_{G_9}^{-3} - \circ_{G_{10}}^{-4}, \\
 \circ_{H_1}^{-2} - \circ_{H_2}^{-3} - \circ_{H_3}^{-7} - \circ_{H_4}^{-2} - \circ_{H_5}^{-2} - \circ_{H_6}^{-3} - \circ_{H_7}^{-3}, \\
 \circ_{I_1}^{-7} - \circ_{I_2}^{-2} - \circ_{I_3}^{-2} - \circ_{I_4}^{-2}, \\
 \circ_{J_1}^{-4} - \circ_{J_2}^{-3} - \circ_{J_3}^{-2}, \\
 \circ_L^{-4}
 \end{array}$$

One proceeds to contract these five chains of  $\mathbb{C}\mathbb{P}^1$ 's from  $Z$ . Since Artin's criteria is satisfied ([2]), the contraction produces a projective surface with special quotient singularities. Denote it by  $X$ . At this step, H. Park, J. Park and D. Shin use  $\mathbb{Q}$ -Gorenstein smoothings to deal with the singularities. Each singularity admits a local  $\mathbb{Q}$ -Gorenstein smoothing. In Section 3 of [23], they prove that the local smoothings can actually be glued to a global  $\mathbb{Q}$ -smoothing of the entire singular surface by proving there is no obstruction to do so. The surface of general type  $S$  with  $p_g = 1$ ,  $q = 0$  and  $K^2 = 6$  is a general fiber of the smoothing of  $X$ ; in the papers of H. Park, J. Park and D. Shin,  $S$  is denoted by  $X_t$ .

The argument regarding the minimality of  $S$  goes as follows. Let  $f : Z \rightarrow X$  be the contraction map of the chains of  $\mathbb{C}\mathbb{P}^1$ 's from  $Z$  to the singular surface  $X$ . By using the technique in, for example, Section 5 in [22], one sees that the pullback  $f^*K_X$  of the canonical divisor  $K_X$  of  $X$  is effective and nef. Therefore,  $K_X$  is nef as well, which implies the minimality of  $S$ .

### 3. Existence of Einstein metrics

The existence of an Einstein metric on a certain manifold is hard to prove. In the case of interest of this paper, where the manifold is a minimal complex surface of general type that does not contain any  $(-2)$ -curves, the following criterion was found independently by T. Aubin and by S.T. Yau.

**Theorem 7** (Aubin [3], Yau [26]). *A compact complex manifold  $(M^4, J)$  admits a compatible Kähler–Einstein metric with  $s < 0$  if and only if its canonical line bundle  $K_M$  is ample. When such a metric exists, it is unique, up to an overall multiplicative constant.*

In order to apply Theorem 7, the following result needs to be proven.

**Proposition 8.** *There exist simply connected surfaces of general type with  $p_g = 1$ ,  $q = 0$ ,  $K^2 = 1, 2, 3, 4, 5, 6$  or 8, and ample canonical bundle.*

The rest of the section is devoted to such endeavor. We carry out the argument for the surface with  $K^2 = 6$ . The other examples can be dealt with in a similar fashion.

**3.1. Proof of Proposition 8.** The following proof follows closely the argument of R. Răşdeaconu and I. Şuvaina used to prove Theorem 1.1 in [24].

**Proof.** Theorem 3 settles the existence part of the proposition. According to [23], in  $Z$  there are five disjoint linear chains. Using the labels we put on their dual graphs in Section 2, let us denote them by  $G = \sum_{i=1}^{11} G_i$ ,  $H = \sum_{i=1}^7 H_i$ ,  $I = \sum_{i=1}^4 I_i$ ,  $J = \sum_{i=1}^3 J_i$ , and let  $L$  be the chain of length one. Name  $F_i$ ,  $i = 1, \dots, 11$  the eleven smooth curves of self-intersection  $-1$  represented by dotted lines labeled  $-1$  in Fig 13 of [23]. We point out that the Poincaré duals of the irreducible components of the five chains and those of the curves  $F_i$ 's form a basis of  $H^2(Z; \mathbb{Q})$ .

Let  $f : Z \rightarrow X$  be the contraction map. Then, one has

$$f^*K_X \equiv_{\mathbb{Q}} \sum_{i=1}^{11} a_i F_i + \sum_{i=1}^{10} b_i G_i + \sum_{i=1}^7 c_i H_i + \sum_{i=1}^4 d_i I_i + \sum_{i=1}^3 e_i J_i + l_1 L.$$

The coefficients that appear above can be computed explicitly (see [22]). However, for our agenda it suffices to know that they are positive rational numbers. In particular the pullback of the canonical divisor of the singular variety to its minimal resolution is effective. Set the exceptional divisor of  $f$  to be  $\text{Exc}(f) = \sum G_i + \sum H_i + \sum I_i + \sum J_i + L$ .

We wish to show that the canonical bundle  $K_X$  of the  $\mathbb{Q}$ -Gorenstein smoothing is ample. This implies our claim: indeed, remember  $S$  is a general fiber of the  $\mathbb{Q}$ -Gorenstein smoothing  $X$ , and ampleness is an open property ([16]). Moreover, we know  $K_X$  is nef. To show it is ample as well, we proceed by contradiction.

Suppose  $K_X$  is not ample. By its nefness and according to the Nakai–Moishezon criterion ([16]), there exists an irreducible curve  $C \subset X$  such that  $(K_X \cdot C) = 0$ .

The total transform of  $C$  in  $Z$  is

$$f^*C \equiv_{\mathbb{Q}} C' + \sum_{i=1}^{10} w_i G_i + \sum_{i=1}^7 x_i H_i + \sum_{i=1}^4 y_i I_i + \sum_{i=1}^3 z_i J_i + tL.$$

Here  $C'$  stands for the strict transform of  $C$ , and the coefficients  $w_i, x_i, y_i, z_i, t$  are nonnegative rational numbers. It is straight-forward to see that  $C'$  is not numerically equivalent to 0 ([24]).

We compute

$$\begin{aligned} (K_X \cdot C) &= (f^*K_X \cdot f^*C) \\ &= (f^*K_X \cdot C') \\ &= \sum_{i=1}^{11} a_i (F_i \cdot C') + \sum_{i=1}^{10} b_i (G_i \cdot C') + \sum_{i=1}^7 c_i (H_i \cdot C') \\ &\quad + \sum_{i=1}^4 d_i (I_i \cdot C') + \sum_{i=1}^3 e_i (J_i \cdot C') + l_1 (L \cdot C'). \end{aligned}$$

The intersection number of the curve  $C'$  with any component of the exceptional divisor  $\text{Exc}(f)$  is greater or equal to zero. The equality is achieved only in the case when  $C'$  is disjoint to all the irreducible components of  $\text{Exc}(f)$ ; this is equivalent to the curve  $C$  missing the singular points of  $X$ . This is

$$\sum_{i=1}^{10} b_i (G_i \cdot C') + \sum_{i=1}^7 c_i (H_i \cdot C') + \sum_{i=1}^4 d_i (I_i \cdot C') + \sum_{i=1}^3 e_i (J_i \cdot C') + l_1 (L \cdot C') \geq 0.$$

Thus, we have  $\sum_{i=1}^{11} a_i(F_i \cdot C') \leq 0$ . At this point there are two possible scenarios:

- There is an  $i_0 \in \{1, \dots, 11\}$  such that  $(C' \cdot F_{i_0}) < 0$ .
- The equality  $(C' \cdot F_i) = 0$  holds for all  $i = 1, \dots, 11$ .

The first scenario requires  $C'$  to coincide with  $F_{i_0}$ . This is not the case, since given that  $f^*K_X$  is nef,  $(f^*K_X \cdot F_i) > 0$  holds for all  $i = 1, \dots, 11$ , which is impossible by our assumption. Thus, the intersection number of the curve  $C'$  with all the  $F_i$ 's and with all of the irreducible components of  $\text{Exc}(f)$  must be zero. However, as it was remarked earlier, the Poincaré duals of the  $F_i$ 's and those of the irreducible components of  $\text{Exc}(f)$  generate  $H^2(Z; \mathbb{Q})$ . This implies that  $C'$  would have to be numerically trivial on  $Z$ . This is a contradiction.

Thus,  $K_X$  is ample. The proposition now follows from Aubin–Yau’s criterion (Theorem 7). □

**Corollary 9.** *There exist a minimal complex structure on  $3\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$ , for each  $k = 11, 13, 14, 15, 16, 17, 18$ , which admits a Kähler–Einstein metric of negative scalar curvature.*

#### 4. Nonexistence of Einstein metrics: Exotic smooth structures

Topologically there is no obstruction for the existence of an Einstein metric on the surfaces of general type we are working with (cf. Corollary 5). We now proceed to study the nonexistence of Einstein metrics with respect to their exotic differential structures.

When one considers different smooth structures on 4-manifolds, the main obstruction to the existence of an Einstein metric is the following result, which generalizes work done by C. LeBrun in [18].

**Theorem 10** (LeBrun, [20]). *Let  $X$  be a compact oriented 4-manifold with a nontrivial Seiberg–Witten invariant and with  $(2\chi + 3\sigma)(X) > 0$ . Then*

$$M = X \#_r \overline{\mathbb{C}\mathbb{P}^2}$$

*does not admit an Einstein metric if  $r \geq \frac{1}{3}(2\chi + 3\sigma)(X)$ .*

As a corollary we have:

**Proposition 11.** *Let  $9 \leq k \leq 18$ . The topological manifolds*

$$3\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$$

*support infinitely many smooth structures that do not admit an Einstein metric. Moreover, each of these manifolds admits infinitely many smooth structures, all of which have negative Yamabe invariant, and on which there are no nonsingular solutions to the normalized Ricci flow for any initial metric.*

**Proof.** We make use of the infinite family  $\{X_n\}$  of pairwise nondiffeomorphic 4-manifolds (with nontrivial SW) sharing the topological prototype  $3\mathbb{C}\mathbb{P}^2\#4\overline{\mathbb{C}\mathbb{P}^2}$  built in [1]. The first part of the lemma now follows by setting  $r \geq 5$  in LeBrun’s result (Theorem 10); notice that the blow-up formula [11, Theorem 1.4] allows us to conclude that the manifolds in the infinite family  $\{X_n\#(4+r)\overline{\mathbb{C}\mathbb{P}^2}\}$  are pairwise nondiffeomorphic. For the claims regarding the Yamabe invariant and the solutions to the normalized Ricci flow see Section 5 below.  $\square$

**4.1. Nonexistence of anti-self-dual Einstein metrics.** Using another obstruction theorem of LeBrun in [20, Theorem 3.8] we obtain the following lemma.

**Lemma 12.** *Let  $8 \leq k \leq 18$ . The topological manifolds*

$$3\mathbb{C}\mathbb{P}^2\#k\overline{\mathbb{C}\mathbb{P}^2}$$

*support infinitely many smooth structures that do not admit an anti-self-dual Einstein metric.*

The reducible smooth structures on the spaces of Lemma 12 are obtained by blowing up a number of points on each member of the infinite family  $\{X_n\}$  of pairwise nondiffeomorphic manifolds that are homeomorphic to  $3\mathbb{C}\mathbb{P}^2\#4\overline{\mathbb{C}\mathbb{P}^2}$  [1] (cf. Proof of Proposition 11).

## 5. Proof of Proposition 2

The following argument is based on the proof of Theorem B in [15].

**Proof.** We start with the part of (1) concerning the sign of the Yamabe invariant. Consider the smooth structure related to the minimal surfaces of general type taken from [23]. By [19] their Yamabe invariant is negative. The existence of nonsingular solutions to the normalized Ricci flow follows from Cao’s theorem ([6], [7]) by taking as an initial metric the Kähler metric with Kähler form the cohomology class of the canonical line bundle.

For Property (2), consider the smooth structures used in Theorem 1 that were built by A. Akhmedov and B.D. Park in [1]: the infinite family  $\{X_n\}$  of pairwise nondiffeomorphic minimal manifolds homeomorphic to  $3\mathbb{C}\mathbb{P}^2\#4\overline{\mathbb{C}\mathbb{P}^2}$ . These manifolds have nontrivial Seiberg–Witten invariants, and for all of them  $c_1^2 > 0$  holds. Thus, by [18], their Yamabe invariant is strictly negative. By a result of M. Ishida (Theorem B in [13]), there are no solutions to the normalized Ricci flow on  $X_i$  for any  $i$  and any initial metric.  $\square$

## References

- [1] AKHMEDOV, ANAR; PARK, B. DOUG. Exotic smooth structures on small 4-manifolds with odd signatures. *Invent. Math.* **181** (2010), no. 3, 577–603. [MR2660453](#)



- (2012b:57048), [Zbl 1206.57029](#), [arXiv:math/0701829](#), doi: [10.1007/s00222-010-0254-y](#).
- [2] ARTIN, MICHAEL. Some numerical criteria for contractability of curves on algebraic surfaces. *Amer. J. Math.* **84** (1962), 485–496. [MR0146182](#) (26 #3704), [Zbl 0105.14404](#), doi: [10.2307/2372985](#).
- [3] AUBIN, THIERRY. Équations du type Monge–Ampère sur les variétés kähleriennes compactes. *C.R. Acad. Sci. Paris Sér. A-B* **283** (1976), no. 3, Aiii, A119–A121. [MR0433520](#) (55 #6496), [Zbl 0333.53040](#).
- [4] BARTH, WOLF P.; HULEK, KLAUS; PETERS, CHRIS A. M.; VAN DE VEN, ANTONIUS. Compact complex surfaces. Second edition. *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Springer-Verlag, Berlin*, 2004. xii+436 pp. ISBN: 3-540-00832-2. [MR2030225](#) (2004m:14070), [Zbl 1036.14016](#).
- [5] BRAUNGARDT, V.; KOTSCHICK, D. Einstein metrics and the number of smooth structures on a four-manifold. *Topology* **44** (2005), no. 3, 641–659. [MR2122219](#) (2005i:57040), [Zbl 1068.57017](#), [arXiv:math/0306021](#), doi: [10.1016/j.top.2004.11.001](#).
- [6] CAO, HUAI DONG. Deformation of Kähler metrics to Kähler–Einstein metrics on compact Kähler manifolds. *Invent. Math.* **81** (1985), no. 2, 359–372. [MR0799272](#) (87d:58051), [Zbl 0574.53042](#), doi: [10.1007/BF01389058](#).
- [7] CAO, HUAI–DONG; CHOW, BENNETT. Recent developments on the Ricci flow. *Bull. Amer. Math. Soc. (N.S.)* **36** (1999), no. 1, 59–74. [MR1655479](#) (99j:53044), [Zbl 0926.53016](#), [arXiv:math/9811123](#), doi: [10.1090/S0273-0979-99-00773-9](#).
- [8] CATANESE, FABRIZIO. Surfaces with  $K^2 = p_g = 1$  and their period mapping. Algebraic geometry, 1–29. *Lecture Notes in Math.*, 732, *Springer, Berlin*, 1979. [MR0555688](#) (81c:14018), [Zbl 0423.14019](#).
- [9] DONALDSON, S. K. An application of gauge theory to four dimensional topology. *J. Differential Geom.* **18** (1983), no. 2, 279–315. [MR0710056](#) (85c:57015), [Zbl 0507.57010](#).
- [10] FREEDMAN, MICHAEL HARTLEY. The topology of four-dimensional manifolds. *J. Differential Geom.* **17** (1982), no. 3, 357–453. [MR0679066](#) (84b:57006), [Zbl 0528.57011](#).
- [11] FINTUSHEL, RONALD; STERN, RONALD J. Immersed spheres in 4-manifolds and the immersed Thom conjecture. *Turkish J. Math.* **19** (1995), no. 2, 145–157. [MR1349567](#) (96j:57036), [Zbl 0869.57016](#).
- [12] HITCHIN, NIGEL. Compact four-dimensional Einstein manifolds. *J. Differential Geom.* **9** (1974), 435–441. [MR0350657](#) (50 #3149), [Zbl 0281.53039](#).
- [13] ISHIDA, MASASHI. The normalized Ricci flow on four-manifolds and exotic smooth structures. 2008. [arXiv:0807.2169](#).
- [14] ISHIDA, MASASHI; LEBRUN, CLAUDE. Spin manifolds, Einstein metrics, and differential topology. *Math. Res. Lett.* **9** (2002), no. 2–3, 229–240. [MR1909640](#) (2003f:53079), [Zbl 1017.53045](#), [arXiv:math/0107111](#).
- [15] ISHIDA, MASASHI; RĂSDEACONU, RAREȘ; ȘUVAINA, IOANA. On normalized Ricci flow and smooth structures on four-manifolds with  $b^+ = 1$ . *Arch. Math.* **92** (2009), no. 4, 355–365. [MR2501291](#) (2010j:53130), [Zbl 1171.53042](#), [arXiv:0810.2465](#), doi: [10.1007/s00013-009-3098-1](#).
- [16] KOLLÁR, JÁNOS; MORI, SHIGEFUMI. Birational geometry of algebraic varieties. With the collaboration of C. H. Clemens and A. Corti. Translated from the 1998 Japanese original. *Cambridge Tracts in Mathematics*, 134. *Cambridge University Press, Cambridge*, 1998. viii+254 pp. ISBN: 0-521-63277-3. [MR1658959](#) (2000b:14018), [Zbl 0926.14003](#), doi: [10.1017/CBO9780511662560](#).
- [17] KOTSCHICK, D. Einstein metrics and smooth structures. *Geom. Topol.* **2** (1998), 1–10. [MR1489351](#) (99h:57060), [Zbl 0885.57014](#), [arXiv:math/9801156](#), doi: [10.2140/gt.1998.2.1](#).

- [18] LEBRUN, CLAUDE. Four-manifolds without Einstein metrics. *Math. Res. Lett.* **3** (1996), no. 2, 133–147. [MR1386835](#) (97a:53072), [Zbl 0856.53035](#), [arXiv:dg-ga/9511015](#).
- [19] LEBRUN, CLAUDE. Kodaira dimension and the Yamabe problem. *Comm. Anal. Geom.* **7** (1999), no. 1, 133–156. [MR1674105](#) (99m:58056), [Zbl 0996.32009](#), [arXiv:dg-ga/9702012](#).
- [20] LEBRUN, CLAUDE. Ricci curvature, minimal volumes, and Seiberg–Witten theory. *Invent. Math.* **145** (2001), no. 2, 279–316. [MR1872548](#) (2002h:53061), [Zbl 0999.53027](#), [arXiv:math/0003068](#), doi: [10.1007/s002220100148](#).
- [21] LEBRUN, CLAUDE. Einstein metrics, complex surfaces, and symplectic 4-manifolds. *Math. Proc. Cambridge Philos. Soc.* **147** (2009), no. 1, 1–8. [MR2507306](#) (2010h:53056), [Zbl 1175.57014](#), [arXiv:0803.3743](#).
- [22] PARK, HEESANG; PARK, JONGIL; SHIN, DONGSOO. A simply connected surface of general type with  $p_g = 0$  and  $K^2 = 3$ . *Geom. Topol.* **13** (2009), no. 2, 743–767. [MR2469529](#) (2009m:14057), [Zbl 1181.14042](#), [arXiv:0708.0273](#), doi: [10.2140/gt.2009.13.743](#).
- [23] PARK, HEESANG; PARK, JONGIL; SHIN, DONGSOO. Surfaces of general type with  $p_g = 1$  and  $q = 0$ . *J. Korean Math. Soc.* **50** (2013), no. 3, 493–507. [arXiv:0906.5195](#).
- [24] RĂSDEACONU, RAREȘ; ȘUVAINA, IOANA. Smooth structures and Einstein metrics on  $\mathbb{C}P^2 \# 5, 6, 7\overline{\mathbb{C}P^2}$ . *Math. Proc. Cambr. Phil. Soc.* **147** (2009), no. 2, 409–417. [MR2525934](#) (2010i:53080), [Zbl 1185.53049](#). doi: [10.1017/S0305004109002527](#).
- [25] ROHLIN, V. A. New results in the theory of four dimensional manifolds. *Doklady Akad. Nauk. SSSR* **84** (1952), 221–224. [MR0052101](#) (14,573b).
- [26] YAU, SHING TUNG. Calabi’s conjecture and some new results in algebraic geometry. *Proc. Nat. Acad. Sci. U.S.A.* **74** (1997), no. 5, 1789–1799. [MR0451180](#) (56 #9467), [Zbl 0355.32028](#), doi: [10.1073/pnas.74.5.1798](#).

CALIFORNIA INSTITUTE OF TECHNOLOGY — MATHEMATICS, 1200 E CALIFORNIA BLVD,  
 PASADENA, CA 91125  
[rtorresr@caltech.edu](mailto:rtorresr@caltech.edu)

This paper is available via <http://nyjm.albany.edu/j/2013/19-12.html>.