

A gauge theoretic approach to Einstein 4-manifolds

Joel Fine, Kirill Krasnov and Dmitri Panov

ABSTRACT. This article investigates a new gauge theoretic approach to Einstein’s equations in dimension 4. Whilst aspects of the formalism are already explained in various places in the mathematics and physics literature, our first goal is to give a single coherent account of the theory in purely mathematical language. We then explain why the new approach may have important mathematical applications: the possibility of using the calculus of variations to find Einstein 4-manifolds, as well as links to symplectic topology. We also carry out some of the technical groundwork to attack these problems.

CONTENTS

1. Introduction	294
Acknowledgements	294
1.1. Main idea	294
1.2. Drawbacks	295
1.3. Applications	295
2. Einstein’s equations as a gauge theory	296
2.1. Definite connections as potentials for conformal classes	296
2.2. The sign of a definite connection	296
2.3. Definite connections as potentials for metrics	297
2.4. Reformulation of the Einstein equations	298
2.5. The action	300
2.6. Topological bounds	301
2.7. Potential extensions of the formalism	302
3. Related action principles	303
3.1. The Einstein–Hilbert action	303
3.2. Eddington’s action for torsion-free affine connections	303

Received January 13, 2014; revised February 27, 2014.

2010 *Mathematics Subject Classification*. 53C25, 53C07, 53D35, 58E30.

Key words and phrases. Einstein manifolds, 4-manifolds, gauge theory.

JF was supported by an Action de Recherche Concertée and by an Interuniversity Action Poles grant.

KK was supported by ERC Starting Grant 277570-DIGT.

DP is a Royal Society Research Fellow.

3.3. Hitchin’s functional and stable forms	304
4. The symplectic geometry of definite connections	305
4.1. The symplectic manifold associated to a definite connection	305
4.2. Symplectic Fano manifolds	306
4.3. A gauge theoretic “sphere” conjecture for positive definite connections	306
4.4. Negative definite connections	309
5. The Hessian and gradient flow of the volume functional	310
5.1. The Hessian is elliptic modulo gauge	310
5.2. The Hessian as a spin Laplacian	318
5.3. Short time existence of the gradient flow of S	320
References	321

1. Introduction

The focus of this article is a new approach to Einstein 4-manifolds, introduced in the physics literature by the second named author [23] and independently, albeit in a weaker form focusing on *anti-self-dual* Einstein metrics, by the first named author [7]. As things stand, the full description of this method is somewhat inaccessible to mathematicians, the necessary background material being spread over several articles written primarily for physicists. Our first goal is to rectify this by giving a single coherent account of the theory in purely mathematical language (§2 below). Our discussion is also more complete in several places than that currently available.

As we will explain, this formalism potentially has important mathematical applications: it opens up a new way to use the calculus of variations to find Einstein metrics and also has possible applications to symplectic topology. Our second goal is to lay some of the ground work in these directions, ask questions and state some conjectures which we hope will inspire future work (see §4 and §5).

Acknowledgements. We would like to thank Claude LeBrun and Misha Verbitsky for helpful discussions, particularly concerning §4.3. We would also like to thank the anonymous referee of the first draft of this article for suggesting we consider the situation treated in Theorem 4.12.

1.1. Main idea. The key idea is to rewrite Einstein’s equations in the language of *gauge theory*, placing them in a similar framework to Yang–Mills theory over 4-manifolds. This reformulation is special to dimension four. We give here the Riemannian version, but one can work equally with Lorentzian signature, replacing $\mathrm{SO}(3)$ throughout by $\mathrm{SL}(2, \mathbb{C})$, as is explained in [23]. We give the details in §2, but put briefly it goes as follows:

- Given an $\mathrm{SO}(3)$ -connection A over a 4-manifold X which satisfies a certain curvature inequality, we associate a Riemannian metric g_A on X , defined algebraically from the curvature of A . One can think of A as a “potential” for the metric, analogous to the relationship between the electromagnetic potential and field.
- To such connections we also associate an action $S(A) \in \mathbb{R}$, which is simply the volume of the metric g_A . Critical points of the action solve a second order PDE which implies that g_A is Einstein.
- There is an a priori topological bound for $S(A)$ which is attained precisely when A solves a first order PDE (which then implies the second-order equation alluded to above). When this happens the corresponding metric g_A is both anti-self-dual and Einstein, with nonzero scalar curvature.

The parallel with Yang–Mills theory is clear: S plays the rôle of Yang–Mills energy, with Einstein metrics corresponding to Yang–Mills connections and anti-self-dual Einstein metrics being the instantons of the theory. In §3, we briefly discuss other related action principles: the Einstein–Hilbert action, Eddington’s action for affine connections and Hitchin’s volume functional for stable forms.

1.2. Drawbacks. Before explaining the theory in more detail, it is important to note its current principal failing: the Einstein metrics which arise are precisely those for which $\frac{s}{12} + W^+$ is a definite endomorphism of Λ^+ . (Here s is the scalar curvature and W^+ is the self-dual Weyl curvature.) We discuss briefly what modifications may be needed to accommodate more general Einstein metrics in §2.7, but there remains much to be done in this direction.

The only possible compact Einstein manifolds for which $\frac{s}{12} + W^+$ is positive definite are the standard metrics on S^4 and $\mathbb{C}\mathbb{P}^2$ (with the noncomplex orientation). This is proved in Theorem 4.14, following an argument which was explained to us by Claude LeBrun.

When $\frac{s}{12} + W^+$ is negative definite the only known compact examples are hyperbolic and complex-hyperbolic metrics (the latter again having the noncomplex orientation). Note that, just as for the positive case, here one even has $W^+ = 0$. It is an interesting open question as to whether these are the only such examples (Question 4.17 below). Some candidate manifolds which have a chance to support such Einstein metrics are described in §4.4.

1.3. Applications. We believe this reformulation of Einstein’s equations will have important applications. Firstly, it reveals a new link between Einstein 4-manifolds and symplectic Fano and Calabi–Yau 6-manifolds with a certain geometric structure, expanding on that discussed by the first and third named authors [8]. We explain this in §4, where we state Conjecture 4.8, which claims that certain 6-dimensional symplectic Fanos are actually algebraic. Proving such a result seems out of reach by current methods

in symplectic topology so it is intriguing that the formalism described here suggests a line of attack. Conjecture 4.8 can also be viewed as a “gauge theoretic sphere theorem”, saying that only the four-sphere and complex projective plane admit connections whose curvature satisfies a certain inequality. (This might be compared to “sphere theorems” in Riemannian geometry, which say that only certain special manifolds admit Riemannian metrics satisfying certain Riemannian curvature inequalities.)

Secondly, the formalism described here gives a new variational approach to Einstein metrics. We lay the groundwork for this in §5. The traditional action principle for Einstein metrics, via the Einstein–Hilbert action, is not well suited to the calculus of variations, ultimately because its Hessian has an infinite number of eigenvalues of both signs. In our setting however, this problem does not arise: *the Hessian of the volume functional is elliptic with finitely many positive eigenvalues*. This is Theorem 5.1. We also briefly discuss the gradient flow of S , the analogue of the Yang–Mills flow in this context, proving short time existence in Theorem 5.18.

2. Einstein’s equations as a gauge theory

2.1. Definite connections as potentials for conformal classes. We begin with the curvature inequality for an $\mathrm{SO}(3)$ -connection mentioned above, and which first appeared in [8].

Definition 2.1. A metric connection A in an $\mathrm{SO}(3)$ -bundle $E \rightarrow X$ over a 4-manifold is called *definite* if whenever u, v are independent tangent vectors, $F_A(u, v) \neq 0$.

Plenty of examples of definite connections are given in [8]. Particularly important to our discussion are those carried by S^4 , $\mathbb{C}\mathbb{P}^2$, hyperbolic and complex-hyperbolic 4-manifolds. For each of these Riemannian manifolds the Levi-Civita connection on Λ^+ is definite (where for the complex surfaces we take self-dual forms with respect to the noncomplex orientation).

Given a definite connection A there is a unique conformal class for which A is a self-dual instanton. To see this note that on a 4-manifold a conformal class is determined by the corresponding sub-bundle $\Lambda^+ \subset \Lambda^2$ of self-dual 2-forms. Now, given a local frame e_1, e_2, e_3 for $\mathfrak{so}(E)$, write $F_A = \sum F_i \otimes e_i$ for a triple of 2-forms F_i . We then take the span of the F_i to be Λ^+ . We must check that this sub-bundle satisfies the necessary algebraic condition to be the self-dual 2-forms of some conformal class, namely that the matrix $F_i \wedge F_j$ of volume forms is definite. This, it turns out, is equivalent to A being a definite connection. In what follows we write Λ_A^+ for the bundle of self-dual 2-forms to emphasise its dependence on A . (Notice that this also implicitly orients X : given any nonzero $\alpha \in \Lambda_A^+$, the square $\alpha \wedge \alpha$ is positively oriented.)

2.2. The sign of a definite connection. Unlike for an arbitrary connection, it is possible to give a sign to the curvature of a definite connection.

The starting point is the fact that the Lie algebra $\mathfrak{so}(3)$ carries a natural orientation. One way to see this is to begin with two linearly independent vectors $e_1, e_2 \in \mathfrak{so}(3)$ and then declare $e_1, e_2, [e_1, e_2]$ to be an oriented basis. One must then check that this orientation does not depend on the initial choice of e_1, e_2 . Alternatively, and more invariantly, pick an orientation on \mathbb{R}^3 ; the cross product then gives an isomorphism $\mathbb{R}^3 \rightarrow \mathfrak{so}(3)$ which one can use to push the orientation from \mathbb{R}^3 to $\mathfrak{so}(3)$. If one begins with the opposite orientation on \mathbb{R}^3 , the cross-product changes sign and so the resulting orientation on $\mathfrak{so}(3)$ is unchanged.

One consequence of this is that when X^4 is given a conformal structure, the resulting bundle Λ^+ carries a natural orientation. Again, there are various ways to see this. For example, picking a metric in the conformal class, the splitting $\Lambda^2 = \Lambda^+ \oplus \Lambda^-$ corresponds to the Lie algebra isomorphism $\mathfrak{so}(4) = \mathfrak{so}(3) \oplus \mathfrak{so}(3)$, making Λ^+ into a bundle of $\mathfrak{so}(3)$ Lie algebras. Equivalently, given an orthonormal basis $\omega_1, \omega_2, \omega_3$ of Λ^+ at some point p , using the metric to “raise an index” on $\sqrt{2}\omega_i$ defines a triple J_1, J_2, J_3 of almost complex structures on T_pX . There are two possibilities: either the J_i satisfy the quaternion relations, or the $-J_i$ do; the first corresponds to the original choice of basis ω_i having positive orientation.

Definition 2.2. Let A be a definite connection in an $SO(3)$ -bundle $E \rightarrow X$. Let e_1, e_2, e_3 be an oriented local frame of $\mathfrak{so}(E)$ and write $F_A = \sum F_i \otimes e_i$. A is called *positive definite* if F_1, F_2, F_3 is an oriented basis for Λ_A^+ and *negative definite* otherwise.

For the standard metrics on S^4 and $\overline{\mathbb{C}\mathbb{P}^2}$ the Levi–Civita connections on Λ^+ are positive definite, whilst hyperbolic and complex-hyperbolic metrics give negative definite connections.

2.3. Definite connections as potentials for metrics. In what follows we fix an orientation on E . Together with the fibrewise metric this gives orientation preserving isomorphisms $E^* \cong E \cong \mathfrak{so}(E)$ and we will freely identify all these bundles.

We have explained how a definite connection A gives rise to a conformal structure on X , the unique one making A into a self-dual instanton. To specify a metric in this conformal class we need to chose a volume form. To see how to do this, pick an arbitrary positively oriented volume form ν . This gives a fibrewise inner-product on Λ_A^+ . Now we can interpret

$$F_A \in \Lambda_A^+ \otimes \mathfrak{so}(E) \cong \text{Hom}(E, \Lambda_A^+)$$

as an isomorphism $E \rightarrow \Lambda_A^+$. Pulling back the inner-product from Λ_A^+ via this isomorphism gives a new metric in E which differs from the old one by a self-adjoint endomorphism $M_\nu \in \text{End}(E)$. In terms of an orthonormal local trivialisation e_1, e_2, e_3 of E in which $F_A = \sum F_i \otimes e_i$, the matrix representative of M_ν is determined by

$$M_{ij} \nu = F_i \wedge F_j.$$

Note that ν and M_ν are in inverse proportion, so specifying a volume form is the same as fixing the scale of M_ν .

A particularly judicious choice of volume form is the following: let ν be any background choice of (positively oriented) volume form and let Λ be any nonzero constant whose sign agrees with that of A . Set

$$(1) \quad \nu_A = \frac{1}{\Lambda^2} \left(\text{Tr} \sqrt{M_\nu} \right)^2 \nu,$$

where \sqrt{M} denotes the positive definite square root of M . The homogeneity of (1) means that ν_A does not depend on the choice of ν . To ease the notation, in what follows we write M_A for M_{ν_A} . An equivalent definition of ν_A is to demand that if A is positive definite, $\text{Tr} \sqrt{M_A} = \Lambda$ and if A is negative definite, we have $\text{Tr} \sqrt{M_A} = -\Lambda$.

Definition 2.3. Given a definite connection A , we write g_A for the resulting Riemannian metric with volume form ν_A defined via (1) and which makes the definite connection A a self-dual instanton.

The justification for this definition of ν_A is the following result.

Lemma 2.4. *Let g be an Einstein metric, $\text{Ric} = \Lambda g$ and suppose that $\frac{\Lambda}{3} + W^+$ is a definite endomorphism of Λ^+ . Then the Levi-Civita ∇ connection on Λ^+ is definite and $g_\nabla = g$.*

Proof. Write ν for the volume form of the Einstein metric. Calculation gives

$$(2) \quad M_\nu = \left(\frac{\Lambda}{3} + W^+ \right)^2.$$

When $\frac{\Lambda}{3} + W^+$ is positive definite, ∇ is a positive definite connection. We can recover the curvature of Λ^+ via

$$\frac{\Lambda}{3} + W^+ = \sqrt{M_\nu}.$$

Now $\text{Tr} \sqrt{M_\nu} = \Lambda$ is constant. Meanwhile, when $\frac{\Lambda}{3} + W^+$ is negative definite, ∇ is a *negative* definite connection. In this case we have

$$\frac{\Lambda}{3} + W^+ = -\sqrt{M_\nu}$$

and $-\text{Tr} \sqrt{M_\nu} = \Lambda$ is again equal to the Einstein constant. In either case, one sees that $\nu = \nu_\nabla$ and so $g = g_\nabla$. \square

2.4. Reformulation of the Einstein equations. We will now give a second order PDE for A which implies that g_A is Einstein. The key is the following well-known observation: an oriented Riemannian 4-manifold (X, g) is Einstein if and only if the Levi-Civita connection on Λ^+ is a self-dual instanton (see for example [1]).

When A is a definite connection in a bundle E , it is automatically a self-dual instanton for g_A and, moreover, E is *isomorphic* to Λ_A^+ . To compare

A and the Levi-Civita connection on Λ_A^+ we need an isometry $E \rightarrow \Lambda_A^+$. As explained above, the curvature provides an isomorphism $F_A: E \rightarrow \Lambda_A^+$ but it is not necessarily isometric, the defect being measured by $M_A \in \text{End}(E)$. To correct for this, define $\Phi_A: E \rightarrow \Lambda_A^+$ by

$$(3) \quad \Phi_A = \pm F_A \circ M_A^{-1/2}$$

where the sign here agrees with that of the connection A . This is an isometry, by definition of M_A , which is moreover orientation preserving thanks to the sign.

Definition 2.5. Given a definite connection A and $\Phi_A: E \rightarrow \Lambda_A^+$ defined as in (3), we write $\text{LC}(A) = \Phi_A^* \nabla$ for the pull-back to E of the g_A -Levi-Civita connection in Λ_A^+ .

In the following result, Φ_A is treated as a 2-form with values in E (via the identification $\text{Hom}(E, \Lambda_A^+) \cong \Lambda_A^+ \otimes E$). The connection A defines a coupled exterior derivative d_A on E -valued forms and so $d_A \Phi_A$ is a 3-form with values in E .

Theorem 2.6 ([23]). *Let A be a definite connection, Λ a nonzero constant whose sign agrees with that of A and Φ_A defined as in (3). If*

$$d_A \Phi_A = 0,$$

then $A = \text{LC}(A)$, g_A is an Einstein metric with $\text{Ric}(g_A) = \Lambda g_A$ and $\frac{\Lambda}{3} + W^+$ definite. Conversely all such metrics are attained this way.

Sketch of proof. If $A = \text{LC}(A)$ then the fact that A is a self-dual instanton implies the Levi-Civita connection is also and hence that g_A is Einstein.

We now need a way to recognise $\text{LC}(A)$ amongst all metric connections in E , something we can do via *torsion*. Given any metric connection ∇ on Λ^+ , its torsion $\tau(\nabla) \in \text{Hom}(\Lambda^+, \Lambda^3)$ is defined, just as for affine connections, as the difference $\tau(\nabla) = d - \sigma \circ \nabla: \Omega^+ \rightarrow \Omega^3$ where d is the exterior derivative and $\sigma: \Lambda^1 \otimes \Lambda^+ \rightarrow \Lambda^3$ is skew-symmetrisation. Paralleling the standard definition of the Levi-Civita connection on the tangent bundle, *the Levi-Civita connection on Λ^+ is the unique metric connection which is torsion free.* (A proof of this well-known fact can be found in [7].)

We can interpret this from the point of view of E . Given a metric connection B in E , we push it forward via Φ_A to a metric connection in Λ_A^+ . The torsion $\tau(\Phi_{A*}(B))$ is identified via Φ_A with the 3-form $d_B \Phi_A$: here the isometry $\Phi_A: E \rightarrow \Lambda_A^+$ is viewed as an E^* -valued 2-form, its coupled exterior derivative is then a section of $\Lambda^3 \otimes E^* \cong \text{Hom}(\Lambda_A^+, \Lambda^3)$ which matches up with $\tau(\Phi_{A*}(B))$. (This calculation is also given in [7].) We now see that $\text{LC}(A)$ is the unique metric connection B for which $d_B \Phi_A = 0$.

The upshot of this discussion is that if $d_A \Phi_A = 0$ then $A = \text{LC}(A)$ and hence g_A is Einstein. It remains to show that $\text{Ric}(g_A) = \Lambda g_A$ and $\frac{\Lambda}{3} + W^+$ is definite.

Since the Levi-Civita connection in Λ^+ is definite, it follows from (2) that $\frac{\Lambda}{3} + W^+$ is necessarily invertible. Assume for a contradiction that $\frac{\Lambda}{3} + W^+$ is indefinite, so that Λ^+ splits into positive and negative eigenbundles. This induces a splitting of E via Φ_A . Now one can check that the equation $d_A \Phi_A = 0$ forces the sub-bundle of rank one to be A -parallel. This implies the curvature of F_A has kernel and so contradicts the fact that A is definite. We also now see from (2) that $\text{Ric}(g_A) = \Lambda g_A$, since $\pm \text{Tr} \sqrt{M_A} = \Lambda$, (with sign corresponding to that of A).

To prove that all such metrics arise, one simply checks that for an Einstein metric g of the given sort the Levi-Civita connection on Λ^+ is a definite connection A for which $\Lambda_A^+ = \Lambda^+$ is unchanged. Moreover, $\nu_A = \text{dvol}_g$, whilst $\Phi_A: \Lambda^+ \rightarrow \Lambda^+$ is the identity. Hence $d_A \Phi_A = 0$ and $g_A = g$. \square

An important special case arises when M_A is a multiple of the identity. Since $\text{Tr} \sqrt{M_A}$ is constant it follows that M_A is a *constant* multiple of the identity and so Φ_A is a constant multiple of F_A . Now $d_A \Phi_A = 0$ follows automatically from the Bianchi identity. Equation (2) shows that g_A is in fact anti-self-dual and Einstein, with nonzero scalar curvature. We state this as a separate result:

Theorem 2.7 ([7]). *Let A be a definite connection. If M_A is a multiple of the identity, then g_A is an anti-self-dual Einstein metric with nonzero scalar curvature. Conversely all such metrics are attained this way.*

2.5. The action. A remarkable feature of definite connections and the corresponding metrics is that *the total volume is an action functional for the theory.*

Definition 2.8 ([23]). Let A be a definite connection in an $\text{SO}(3)$ -bundle E over a compact 4-manifold X . The *action* of A is defined to be

$$S(A) = \frac{\Lambda^2}{12\pi^2} \int_X \nu_A$$

where ν_A is defined by (1).

Theorem 2.9 ([23]). *A definite connection A is a critical point of S if and only if $d_A \Phi_A = 0$. So critical points of S give Einstein metrics.*

Proof. Pick a local orthonormal frame e_1, e_2, e_3 for E and write $F_A = \sum F_i \otimes e_i$ (where we have as always identified $\mathfrak{so}(E) \cong E$ via the cross product). Then $M_{ij} \nu_A = F_i \wedge F_j$, where $M_{ij} = (F_i, F_j)$ is the matrix of pointwise innerproducts of the F_i with respect to g_A . Making an infinitesimal change $\dot{A} = a$ of the connection gives

$$\dot{M}_{ij} \nu_A + M_{ij} \dot{\nu}_A = (d_A a)_i \wedge F_j + F_i \wedge (d_A a)_j$$

where $d_A a = \sum (d_A a)_i \otimes e_i$. Now the fact that $\text{Tr}(\sqrt{M_A})$ is constant implies that $\text{Tr}(M_A^{-1/2} \dot{M}_A) = 0$. Multiply the above equation by $M_A^{-1/2}$ and taking

the trace; then use the fact that for any 2-form α , $F_i \wedge \alpha = (F_i, \alpha)\nu_A$ (since the F_i are self dual) to obtain

$$(4) \quad \dot{\nu}_A = \frac{2}{\Lambda}(\Phi_A, d_A a)\nu_A.$$

Note that the sign in (3) is correctly captured by the factor of Λ here. It follows that

$$\dot{S} = \frac{\Lambda}{6\pi^2} \int_X (d_A^* \Phi_A, a)\nu_A$$

and so the critical points are those A with $d_A^* \Phi_A = 0$. On E -valued 2-forms, $d_A^* = - * d_A *$. Since Φ_A is a self-dual 2-form, $*\Phi_A = \Phi_A$ and so $d_A^* \Phi_A = - * d_A \Phi_A$. Hence the critical points of S are precisely those A for which $d_A \Phi_A = 0$. \square

2.6. Topological bounds. We next explain topological bounds on S .

Proposition 2.10 (cf. [7]). *There is an a priori bound*

$$\frac{1}{3} (2\chi(X) + 3\tau(X)) < S(A) \leq 2\chi(X) + 3\tau(X)$$

for all definite connections. Moreover, $S(A) = 2\chi(X) + 3\tau(X)$ if and only if M_A is a multiple of the identity and hence g_A is anti-self-dual and Einstein with nonzero scalar curvature.

Proof. Given a self-adjoint positive-definite 3-by-3 matrix M ,

$$\text{Tr}(M) < \left(\text{Tr}(\sqrt{M}) \right)^2 \leq 3 \text{Tr}(M)$$

with equality on the right hand side if and only if M is a multiple of the identity. It follows from (1) that for any choice of volume form ν , $\frac{1}{\Lambda^2} \text{Tr}(M_\nu)\nu < \nu_A \leq \frac{3}{\Lambda^2} \text{Tr}(M_\nu)\nu$ with equality on the right if and only if M_ν (and hence M_A), is a multiple of the identity.

Now $\text{Tr}(M_\nu)\nu$ is a multiple of the first Pontryagin form of A . To see this, in a local trivialisation in which $F_A = \sum F_i \otimes e_i$, we have $\text{Tr}(M_\nu)\nu = \sum F_i \wedge F_i$. Meanwhile, $p_1(A) = \frac{1}{4\pi^2} \sum F_i \wedge F_i$. Hence

$$\frac{4\pi^2}{\Lambda^2} p_1(A) < \nu_A \leq \frac{12\pi^2}{\Lambda^2} p_1(A).$$

This, together with the fact that $p_1(E) = p_1(\Lambda^+) = 2\chi(X) + 3\tau(X)$, gives the result. \square

As an immediate corollary we see that the existence of a definite connection implies “one-half” of the Hitchin–Thorpe inequality satisfied by Einstein 4-manifolds [15, 29]:

Corollary 2.11 ([8]). *If X carries a definite connection then*

$$2\chi(X) + 3\tau(X) > 0.$$

We remark that to date *this is the only known obstruction to the existence of definite connections.*

2.7. Potential extensions of the formalism. The Einstein metrics that can be reached using the approach given above are precisely those for which $\frac{\Lambda}{3} + W^+$ is a definite endomorphism of Λ^+ . We now discuss briefly how one might extend the formalism to include more Einstein metrics.

One case that is currently missing is when (M, g) is Einstein and $\frac{\Lambda}{3} + W^+$ is invertible, but not definite. In this case, the Levi-Civita connection A on Λ^+ is still a definite connection, but $g_A \neq g$. This is because in the above recipe for g_A , we fix $\text{Tr}(\sqrt{M}) = |\Lambda|$, where \sqrt{M} is the *positive* square root of $(\frac{\Lambda}{3} + W^+)^2$. But this is only equal to $\pm(\frac{\Lambda}{3} + W^+)$ when all eigenvalues of the later have the same sign. The solution is, of course, to choose a different branch of the square root. Using the right choice for \sqrt{M} , the normalisation $\text{Tr}(\sqrt{M}) = |\Lambda|$ gives $g_A = g$ and the theory proceeds as before. This raises two important questions. Firstly, given a definite connection A , how can one tell in advance which branch of the square root to take? Secondly, one could imagine a situation in which the Levi-Civita connection A of an Einstein metric was, say, positive definite, i.e., that $\det(\frac{\Lambda}{3} + W^+) > 0$, and yet $\Lambda < 0$. This means that one might not know in advance what sign to take for Λ .

The second type of Einstein metric which does not fit directly in the above set-up are those for which eigenvalues of $\frac{\Lambda}{3} + W^+$ vanish somewhere on the manifold. Given such an Einstein metric, with Levi-Civita connection A in Λ^+ , one can proceed as above up to the definition of $\Phi_A = \pm F_A \circ M_A^{-1/2}$, which is not immediately valid where eigenvalues of M_A vanish. On the other hand, since we started with an Einstein metric, one can check that Φ_A extends smoothly over the seemingly singular locus to a globally defined isometry $\Lambda^+ \rightarrow \Lambda^+$. To include such Einstein metrics, we should consider more general connections, which we call *semi-definite*. These are connections for which $M_A \geq 0$ and that $M_A > 0$ in at least one point. We demand moreover that $\Phi_A = \pm F_A \circ M_A^{-1/2}$ (defined initially where $M_A > 0$) extends smoothly to an isometry $E \rightarrow \Lambda^+$ over the whole manifold.

An example which exhibits some of the problems mentioned above is the Page metric [27]. This is an Einstein metric with positive scalar curvature on the one-point blow-up X of $\mathbb{C}\mathbb{P}^2$. It is cohomogeneity-one, being invariant under the $\text{SU}(2)$ -action on X lifting that on $\mathbb{C}\mathbb{P}^2$ which fixes the centre of the blow-up. It can be described via a path $g(t)$ of left-invariant metrics on $\text{SU}(2)$ parametrised by $t \in (0, 1)$, which collapse the fibres of the Hopf map $\text{SU}(2) \rightarrow \mathbb{C}\mathbb{P}^1$ as $t \rightarrow 0, 1$. The metric $dt^2 + g(t)$ on $\text{SU}(2) \times (0, 1)$ extends to give a smooth metric on the blow-up X . The $\mathbb{C}\mathbb{P}^1$ s which compactify the ends of $\text{SU}(2) \times (0, 1)$ are the exceptional curve and the line in $\mathbb{C}\mathbb{P}^2$ which is orthogonal to the centre of the blow-up.

One can check, by direct computation, that the Levi-Civita connection on Λ^+ of the Page metric is positive definite everywhere except for $\text{SU}(2) \times \{t_0\}$ for a single value of t_0 . For $t < t_0$, $\frac{\Lambda}{3} + W^+ > 0$ and we are in exactly the situation treated in this article. At $t = t_0$, two eigenvalues of $\frac{\Lambda}{3} + W^+$ vanish and for $t > t_0$ these same two eigenvalues become negative. Provided one

uses the correct branch of the square root as t crosses t_0 , the whole Page metric can be seen using the gauge theoretic approach. Indeed one can “find” the Page metric by solving the equation $d_A \Phi_A = 0$ for an appropriate cohomogeneity-one definite connection. This directly gives the Page metric over $t \in [0, t_0)$ and one sees clearly that two eigenvalues vanish at $t = t_0$. Then taking a different square root gives the metric over $(t_0, 1]$ and it is a simple matter to see that the two parts combine smoothly to give the whole Page metric.

3. Related action principles

This section gives some additional context to the above formalism by describing some related action principles.

3.1. The Einstein–Hilbert action. The traditional action principle for Einstein metrics is that of Einstein–Hilbert. Given a Riemannian metric g , define the Einstein–Hilbert action of g to be

$$S_{\text{EH}}(g) = \int_X (s - 2\Lambda) \text{dvol}$$

where s is the scalar curvature of g and Λ is the cosmological constant. Critical points of S_{EH} are those metrics with $\text{Ric}(g) = \Lambda g$.

Note that the Einstein–Hilbert functional evaluated on an Einstein metric is a multiple of the volume. By contrast in the gauge-theoretic formulation the action is always the volume of the space, even away from the critical points.

As is well known, the Einstein–Hilbert action is beset with difficulties from both the mathematical and physical points of view. Physically, the resulting quantum theory is not renormalizable. Mathematically, one hopes to use the calculus of variations to find critical points of an action. For S_{EH} this is extremely hard: at a critical point the Hessian has infinitely many positive and negative eigenvalues which leads to seemingly unresolvable problems in min-max arguments. As we discuss in §5, it is precisely these problems that the gauge-theoretic approach outlined above aims to avoid.

3.2. Eddington’s action for torsion-free affine connections. The idea of using a connection as the independent variable, rather than the metric, goes back to Eddington [5]. Let ∇ be a torsion-free affine connection in $TX \rightarrow X$. Such connections have a Ricci tensor, defined exactly as for the Levi–Civita connection: the curvature tensor of ∇ is a section of $\Lambda^2 \otimes TX \otimes T^*X$ and so one can contract the TX factor with the Λ^2 factor to produce a tensor $\text{Ric}_\nabla \in T^*X \otimes T^*X$.

The fact that ∇ is torsion-free implies that Ric_∇ is symmetric. We now focus on those ∇ for which Ric_∇ is a definite form. Pick a nonzero constant Λ which is positive if Ric_∇ is positive definite and negative if Ric_∇ is negative

definite. We can then define a Riemannian metric on X by $g = \Lambda^{-1} \text{Ric}_\nabla$. Eddington's action is again the volume:

$$S_{\text{Edd}}(\nabla) = \int_X \text{dvol}_g.$$

The Euler–Lagrange equations for S_{Edd} are $\nabla \text{Ric}_\nabla = 0$. In other words, ∇ is both torsion-free and makes g parallel, making it the Levi–Civita connection of g . So $\text{Ric}_\nabla = \text{Ric}_g$ is the Ricci curvature and g is Einstein by virtue of its definition as $g = \Lambda^{-1} \text{Ric}_g$.

The parallels with our gauge-theoretic approach are obvious. An important difference, however, is the size of the space of fields. For us the connections are given locally by 12 real-valued functions ($\Lambda^1 \otimes \mathfrak{so}(E)$ has rank 12). The relevant gauge group is that of fibrewise linear isometries $E \rightarrow E$ (not necessarily covering the identity) which has functional dimension 7 (4 diffeomorphisms plus 3 gauge rotations), leaving the space of definite connections modulo gauge with functional dimension 5.

On the other hand, torsion-free affine connections on a 4-manifold are given locally by 40 real-valued functions ($S^2 T^* X \otimes TX$ has rank 40). Moreover, the gauge group is rather small and consists of just diffeomorphisms which have functional dimension 4. This means that Eddington's theory has a configuration space of functional dimension 36. Finally the traditional theory of metrics modulo diffeomorphisms gives a configuration space of functional dimension 6. This vast increase in the “number of fields” makes Eddington's theory even more complicated than the traditional Einstein–Hilbert theory. This is why, in spite of the fact that the action in this formulation is a multiple of the volume, it is unlikely to give a useful variational principle.

3.3. Hitchin's functional and stable forms. The use of total volume as an action has also appeared in the work of Hitchin [17, 18]. In these articles, Hitchin studies “stable forms”. A form $\rho \in \Lambda^p \mathbb{R}^n$ is called stable if its $\text{GL}(n, \mathbb{R})$ -orbit is open in $\Lambda^p \mathbb{R}^n$. A symplectic form on \mathbb{R}^{2m} is one example, but there are also “exceptional” examples with $p = 3$ and $n = 6, 7, 8$. A form ρ on a manifold M is called stable if it is stable at each point. The existence of such a form gives a reduction of the structure group of TM to $G \subset \text{GL}(n, \mathbb{R})$, the stabiliser of ρ at a point. In the three exceptional cases mentioned above, G is $\text{SL}(3, \mathbb{C})$, G_2 and $\text{PSU}(3)$ in dimensions 6, 7 and 8 respectively. In each case G preserves a volume form and so the stable form defines in turn a volume form $\phi(\rho)$ on M .

Hitchin proceeds to study the corresponding volume functional

$$\rho \mapsto \int_M \phi(\rho)$$

restricted those ρ which are closed and lie in a fixed cohomology class. The critical points define interesting geometries. For example, when $p = 3, n = 6$, a critical stable form defines a complex structure on M with trivial canonical

bundle, whilst when $p = 3, n = 7$, a critical stable form defines a metric on M with holonomy G_2 (in particular an Einstein metric).

There is a formal parallel with the set-up described in §2. The curvature F_A of a definite connection is modelled on a “stable” element of $\Lambda^2(\mathbb{R}^4) \otimes \mathbb{R}^3$, in the sense that its orbit under the action of $GL(4, \mathbb{R}) \times SO(3)$ is open. Moreover, the stabiliser preserves the volume form defined by (1). The condition that the stable form be closed has been replaced by the Bianchi identity $d_A F_A = 0$ and the restriction to a fixed cohomology class can be thought of as analogous to considering connections on a fixed bundle: $F_{A+a} = F_A + d_A a + a \wedge a$. The main difference between the situation studied by Hitchin and our set-up is that we work with differential forms taking values in a vector bundle over the manifold (curvature 2-forms), while Hitchin studies ordinary differential forms. Because of this, there are more options for constructing a volume form, leading to a family of actions “deforming” the action S considered here. This is discussed in [22].

4. The symplectic geometry of definite connections

4.1. The symplectic manifold associated to a definite connection.

A definite connection in $E \rightarrow X$ is not just a “potential” for a Riemannian metric on X , it also defines a symplectic form on the unit sphere bundle $\pi: Z \rightarrow X$ of E . The symplectic point of view on definite connections was the original motivation for their introduction in [8]. We briefly recall the construction here.

Given any (not necessarily definite) connection A we define a closed 2-form ω_A on Z as follows. Let $V \rightarrow Z$ be the vertical tangent bundle (i.e., the sub-bundle $\ker \pi_* \leq TZ$). If we orient the fibres of E , then V becomes an oriented $SO(2)$ -bundle, or equivalently a Hermitian complex line bundle. The form ω_A is the curvature form of a certain unitary connection ∇ in V defined as follows. Given a section of V , we differentiate it vertically in Z by using the Levi-Civita connection on the S^2 -fibres. Meanwhile to differentiate horizontally, we use A to identify nearby fibres of $Z \rightarrow X$. Together this defines ∇ and hence the closed form $\omega_A = \frac{i}{2\pi} F_\nabla$.

Lemma 4.1 ([8]). *The form ω_A is symplectic if and only if A is a definite connection.*

Remark 4.2. This is inspired in part by twistor theory. The isometric identification $E \rightarrow \Lambda_A^+$ gives a diffeomorphism of Z with the twistor space of (X, g_A) . However, the twistor almost complex structures are rarely compatible with ω_A . One can check that the Atiyah–Hitchin–Singer almost complex structure J_+ is compatible with ω_A if and only if g_A is anti-self-dual Einstein with positive scalar curvature whilst the Eells–Salamon almost complex structure J_- is compatible with ω_A if and only if g_A is anti-self-dual Einstein with negative scalar curvature. (See [1] and [6] for the definitions and properties of J_+ and J_- respectively.)

4.2. Symplectic Fano manifolds. The symplectic geometry of (Z, ω_A) depends dramatically on the sign of the definite connection. We begin with the following definition (which is not universally standard).

Definition 4.3. A symplectic manifold (M, ω) is called a *symplectic Fano* if $[\omega]$ is a positive multiple of $c_1(M, \omega)$. It is called a *symplectic Calabi–Yau* if $c_1(M, \omega) = 0$.

Proposition 4.4 ([8]). *If A is a positive definite connection then (Z, ω_A) is a symplectic Fano. If A is negative definite then (Z, ω_A) is a symplectic Calabi–Yau.*

Definition 4.3 is of course directly inspired by the similar terms in use in algebraic geometry. The motivation is that in algebraic geometry the study of Fanos and Calabi–Yaus has special features and one is curious to see to what extent these features extend to symplectic geometry. For example, it is known that in each dimension there is a finite number of deformation families of smooth algebraic Fano varieties (see, e.g., the discussion in [20]).

The next two results show the current knowledge on the extent to which symplectic and algebraic Fanos diverge:

Theorem 4.5 (McDuff [25]). *Let (M, ω) be a symplectic Fano 4-manifold. Then there is a compatible complex structure on M making it a smooth algebraic Fano variety.*

Theorem 4.6 (Fine–Panov [9]). *There are symplectic Fanos which are not algebraic, starting at least from dimension 12.*

This leads to the following question (to the best of our knowledge, this first appeared in the literature in [9], but had surely been discussed before):

Question 4.7. *What is the lowest dimension $2n$ in which all symplectic Fanos are necessarily algebraic? (We see from the above that $2 \leq n < 6$.)*

4.3. A gauge theoretic “sphere” conjecture for positive definite connections. Since positive definite connections give rise to symplectic Fano 6-manifolds they are useful for probing the $n = 3$ case of Question 4.7. In this regard we have the following conjecture, which first appeared in [8].

Conjecture 4.8. *Let A be a positive definite connection over a compact 4-manifold. Then the underlying 4-manifold is either S^4 or $\overline{\mathbb{C}\mathbb{P}^2}$ whilst the resulting symplectic Fano is algebraic, being symplectomorphic to the standard structure on either $\mathbb{C}\mathbb{P}^3$ or the complete flag $F(\mathbb{C}^3)$ respectively.*

An initial motivation for this conjecture is the stark contrast between the known examples of positive and negative definite connections. The only known compact positive definite connections are isotopic to the Levi-Civita connections on $\Lambda^+ \rightarrow S^4$ or $\Lambda^+ \rightarrow \overline{\mathbb{C}\mathbb{P}^2}$. The resulting symplectic 6-manifolds are then $\mathbb{C}\mathbb{P}^3$ or $F(\mathbb{C}^3)$. On the other hand, in the many known

examples of negative definite connections over compact 4-manifolds, the resulting symplectic 6-manifold never admits a compatible complex structure. See, for example, the discussions in [8, 9, 10].

Leaving aside Question 4.7, one can view Conjecture 4.8 as a gauge theoretic “sphere” conjecture. Sphere theorems in Riemannian geometry start with a compact Riemannian manifold whose curvature satisfies a certain inequality and deduce that the underlying manifold is diffeomorphic to a sphere, or spherical space form. There is a long history of such theorems. For a survey of classical results and recent developments, see the article of Wilking [31]. A proof of Conjecture 4.8 would be, to the best of our knowledge, the first example of such a theorem with purely gauge theoretic hypotheses, involving the curvature of an auxiliary connection rather than a Riemannian metric. It would also imply a more traditional sphere-type theorem. To see this we first need the following result from [8]:

Theorem 4.9 (Fine–Panov [8]). *Let g be a Riemannian metric on an oriented 4-manifold X . Write Ric_0 for the trace free Ricci curvature of g , interpreted as a map $\Lambda^+ \rightarrow \Lambda^-$. The Levi-Civita connection on Λ^+ is definite if*

$$(5) \quad \left(\frac{s}{12} + W^+\right)^2 > \text{Ric}_0^* \text{Ric}_0.$$

In this case the sign of the connection agrees with that of $\det(s/12 + W^+)$.

With this result in hand, we see that the following purely Riemannian conjecture is implied by the more general Conjecture 4.8:

Conjecture 4.10. *S^4 and $\overline{\mathbb{C}\mathbb{P}^2}$ are the only compact 4-manifolds which admit Riemannian metrics satisfying the curvature inequality (5) and with $s/12 + W^+$ a positive definite endomorphism of Λ^+ .*

A small step in the direction of Conjecture 4.10 was made in [8], where the following result appears:

Theorem 4.11 (Fine–Panov [8]). *Let (M, g) be a compact oriented Riemannian 4-manifold which satisfies the curvature inequality (5) and with $s/12 + W^+$ a positive definite endomorphism of Λ^+ . Then M is homeomorphic to the connected sum of n copies of $\overline{\mathbb{C}\mathbb{P}^2}$ for some $n = 0, 1, 2, 3$.*

More evidence for Conjecture 4.10 is provided by the following result.

Theorem 4.12. *Let (M, g) be a compact oriented Riemannian 4-manifold which satisfies the curvature inequality (5). If g has positive scalar curvature and is anti-self-dual then it is conformal to the standard metric on either S^4 or $\overline{\mathbb{C}\mathbb{P}^2}$.*

The proof is based on the following result of Verbitsky. We also give here a shorter variation of Verbitsky’s proof. (The result we prove here is Theorem 1.1 of [30]; see Definition 1.2 of [30] to pass from the statement there to the language of taming forms.)

Theorem 4.13 (Verbitsky [30]). *Let (M, g) be a compact oriented 4-manifold with anti-self-dual Riemannian metric. If the twistor space Z admits a taming symplectic form then the metric is conformal to the standard metric on either S^4 or $\overline{\mathbb{C}\mathbb{P}^2}$.*

Proof. We begin by showing that Z is Moishezon, using a result of Campana (Theorem 4.5 of [2]). Let C denote the space of analytic cycles of dimension 1 in Z and write C_t for the irreducible 4-dimensional component which contains the vertical twistor lines. Campana's Theorem says that if C_t is compact then Z is Moishezon. In our case, the complex structure is tamed by a symplectic form and so compactness of C_t follows by Gromov's compactness result.

Next, we note that if ω is a taming symplectic form then so is $\omega - \gamma^*\omega$, where $\gamma: Z \rightarrow Z$ is the real involution on the twistor space. This second form is in a multiple of the anti-canonical class. (This follows from the form of the cohomology of Z which is given by Leray–Hirsch, and is described in, for example, [16].) So we can assume the taming form represents $c_1(K^{-1})$. As a consequence, for any k -dimensional subvariety V of Z , $\int_V c_1(K^{-1})^k > 0$. We can now apply the Nakai–Moishezon criterion, which holds for Moishezon manifolds (Theorem 6 of [26], see also Theorem 3.11 of [21]). This tells us that K^{-1} is ample and so Z is a projective manifold. Finally, a theorem of Hitchin [16] guarantees that the standard conformal structures on S^4 and $\overline{\mathbb{C}\mathbb{P}^2}$ are the only ones which yield compact Kähler twistor spaces. \square

Proof of Theorem 4.12. The inequality (5) implies that there is a symplectic form on the twistor space. Moreover, it follows from Theorem 4.4 of [8] that this symplectic form tames the twistor complex structure. The result now follows from Verbitsky's Theorem. \square

As further motivation for both believing Conjecture 4.8 and perhaps seeing how to attack it, we now show it holds for those positive definite connections which are also critical points of the volume function. The following argument was explained to us by Claude LeBrun.

Theorem 4.14. *Let A be a positive definite connection on a compact 4-manifold, solving the equation $d_A\Phi_A = 0$ ensuring that g_A is Einstein. Then g_A is also anti-self-dual. It follows that g_A is the standard metric on either S^4 or $\overline{\mathbb{C}\mathbb{P}^2}$ and the corresponding Fano is symplectomorphic to $\mathbb{C}\mathbb{P}^3$ or $F(\mathbb{C}^3)$ respectively.*

Proof. Since A is positive definite, the scalar curvature $s = 4\Lambda$ of g_A is positive. If g_A is also anti-self-dual then a theorem of Hitchin [16] ensures that the only possibilities are the standard metrics on S^4 and $\overline{\mathbb{C}\mathbb{P}^2}$. The twistor spaces of these 4-manifolds are $\mathbb{C}\mathbb{P}^3$ and $F(\mathbb{C}^3)$ respectively and it is a simple matter to check that the symplectic form ω_A is the standard Kähler form on these spaces (e.g., for symmetry reasons).

Assume then for a contradiction that $W^+ \neq 0$. The key is the following inequality due to Gursky (this first appeared in [12], see also the streamlined proof in [13]). Let g be an Einstein metric with $\text{Ric} = \Lambda g$, where $\Lambda > 0$, on a compact 4-manifold X . If W^+ is *not* identically zero then

$$(6) \quad \int |W^+|^2 \text{dvol} \geq \frac{2\Lambda^2}{3} \text{vol}(g).$$

To exploit this, write $w_1 \leq w_2 \leq w_3$ for the eigenvalues of W^+ . Now $w_1 + w_2 + w_3 = 0$ implies $0 \leq w_3 \leq -2w_1$ and $w_1 \leq w_2 \leq -\frac{1}{2}w_1$. Because $w_1 < 0$ this last two-sided bound implies $|w_2| \leq |w_1|$. Hence $|W^+|^2 = w_1^2 + w_2^2 + w_3^2 \leq 6w_1^2$. Meanwhile, since $\Lambda/3 + W^+$ is positive definite, we see that $w_1^2 < \Lambda^2/9$. Integrating the bounds $|W^+|^2 \leq 6w_1^2 < \frac{2\Lambda^2}{3}$ and applying Gursky’s inequality (6) we obtain a contradiction. \square

Remark 4.15. We remark that this result also follows from recent work of Richard and Seshadri [28]. They study Riemannian 4-manifolds with positive isotropic curvature on self-dual 2-forms (a weakening of the usual positive isotropic curvature inequality, which is required to hold for all 2-forms and which makes sense in any dimension). They prove that the only such compact Einstein metrics are the standard metrics on S^4 and $\overline{\mathbb{C}\mathbb{P}^2}$. They give a different proof of this fact, but as they say it also follows from Gursky’s inequality. The argument is almost identical to that given here: having positive isotropic curvature on self-dual 2-forms is equivalent to $W^+ < s/6$ and so, for an Einstein metric, $w_3 \leq \frac{2\Lambda}{3}$. Reasoning as above then shows that $|W^+|^2 < \frac{2\Lambda^2}{3}$ and so by Gursky’s inequality the metric must actually be anti-self-dual.

We can rephrase Theorem 4.14 as saying that when A is positive definite and a critical point of the volume functional S then in fact $S(A)$ attains the topological maximum $S(A) = 2\chi + 3\tau$. In particular, for positive definite connections over a compact manifold, intermediate critical points are ruled out. One possible approach to proving Conjecture 4.8 would be to use the calculus of variations, or even the gradient flow, to find a maximum of S . The goal would be to prove the following result, which, in view of Theorem 4.14 and Moser’s proof of local rigidity of symplectic structures, implies Conjecture 4.8.

Conjecture 4.16. *Let A be a positive definite connection over a compact 4-manifold. Then A can be smoothly deformed through such connections to a critical point of S .*

In §5 we lay the basic analytic groundwork for this line of attack: we prove that the Hessian of S is an elliptic second-order operator (modulo gauge) and that its gradient flow exists for short time.

4.4. Negative definite connections. Recall that using definite connections, we are able to describe all Einstein metrics for which $\Lambda/3 + W^+$

is definite. Theorem 4.14 says that when $\Lambda/3 + W^+$ is positive definite, the only compact examples are the standard metrics on S^4 and $\overline{\mathbb{C}P}^2$. In the negative definite case, the only *known* examples are hyperbolic and complex-hyperbolic manifolds. (Again, for the complex surfaces we use the noncomplex orientation.)

Question 4.17. *Do there exist compact Einstein 4-manifolds with Einstein constant $\Lambda < 0$, for which $\frac{\Lambda}{3} + W^+$ is negative definite, besides hyperbolic and complex-hyperbolic manifolds?*

We remark that there is a related well-known problem: do there exist compact anti-self-dual Einstein manifolds with negative scalar curvature, besides hyperbolic and complex-hyperbolic manifolds?

One way to try and produce an Einstein manifold answering Question 4.17 is to start with a negative definite connection and deform it to a critical point of the volume functional. Some possible starting connections are described in [8], exploiting a construction of Gromov–Thurston [11]. We recall the idea here.

Let M be an oriented compact hyperbolic 4-manifold with a totally-geodesic nulhomologous surface $\Sigma \subset M$. Since Σ is nulhomologous, for each m there is an m -fold branched cover $X_m \rightarrow M$, with branch locus Σ . Pulling back the hyperbolic metric from M gives a singular metric on X_m . It can be smoothed in such a way as to give a metric for which (5) of Theorem 4.9 is satisfied, with $s/12 + W^+$ a negative definite endomorphism of Λ^+ . It follows that the Levi-Civita connection on $\Lambda^+ \rightarrow X_m$ is negative definite.

Finiteness considerations show that not all X_m achieved this way can admit hyperbolic metrics and in fact Gromov and Thurston conjecture that none of them do. Now one can check that $\tau(X_m) = 0$. This follows from the fact that Σ is nulhomologous and an equivariant version of the signature theorem (see e.g., Equation (15) in the article [14] of Hirzebruch.) Since the signature is a multiple of $\int (|W^+|^2 - |W^-|^2)$ any anti-self-dual metric on X_m is automatically conformally flat. It follows that any anti-self-dual Einstein metric on X_m has constant curvature and hence would be hyperbolic. In particular, those nonhyperbolic X_m (conjecturally all of them) do not admit anti-self-dual Einstein metrics. To answer Question 4.17 then, one might start with the negative definite connection on X_m and try to deform it to a critical point of S , giving an Einstein metric which is not anti-self-dual.

5. The Hessian and gradient flow of the volume functional

In the previous section we discussed finding Einstein metrics by searching for critical points of the volume functional S . In this section we lay the basic groundwork for such an approach.

5.1. The Hessian is elliptic modulo gauge. We begin with the Hessian of the volume function, S , defined on the space of definite connections.

Definite connections satisfy a weak inequality and hence form an open set in the space of *all* connections (in, for example, the C^1 -topology). We use the natural affine structure on the space of connections to take the Hessian of S at A , to get a symmetric bilinear form H on $\Omega^1(X, E)$. (Recall we implicitly identify $\mathfrak{so}(E) \cong E$ throughout.) Using the L^2 -innerproduct defined by g_A we can then realise this as

$$H(a, b) = \int_X (Da, b) \nu_A$$

for a uniquely determined self-adjoint second-order operator D on $\Omega^1(X, E)$, which we also call the Hessian. Our main result concerning D is:

Theorem 5.1. *The Hessian of S is elliptic modulo gauge, with finitely many positive eigenvalues.*

As was mentioned above, this is in sharp contrast to the Hessian of the Einstein–Hilbert action.

The Hessian in general and this result in particular are far simpler to understand at a definite connection for which $M_A = (\Lambda/3)^2 1_E$ is a multiple of the identity (with the multiple chosen so that $\text{Tr}(\sqrt{M_A}) = |\Lambda|$). Recall that for such a connection g_A is anti-self-dual Einstein; meanwhile the functional S attains its a priori topological maximum (equal to $2\chi(X) + 3\tau(X)$) precisely at connections corresponding to ASD Einstein metrics. At such points the functional is clearly concave so must have nonpositive Hessian. On a first reading of the proof of Theorem 5.1, it is useful to focus on this case $M_A = (\Lambda/3)^2 1_E$ where a great many of the formulae and interpretations become simpler. We highlight these simplifications at various points.

5.1.1. Gauge fixing. Before proving Theorem 5.1, we first make precise what we mean by “elliptic modulo gauge”. The gauge group \mathcal{G} is the group of all fibrewise linear isometries $E \rightarrow E$. This acts on connections by pulling back, preserving both the space of definite connections and the action S . Differentiating this action at a connection A gives a map $R_A: \text{Lie}(\mathcal{G}) \rightarrow \Omega^1(X, E)$. Since S is gauge-invariant, it follows that its Hessian D vanishes on $\text{im } R_A$. The claim in Theorem 5.1 is that on the orthogonal complement of $\text{im } R_A$, the Hessian is the restriction of an elliptic operator whose spectrum is bounded above.

To describe this orthogonal complement, we need a concrete description of R_A . First consider the subgroup \mathcal{G}_0 of gauge transformations covering the identity on X (the usual gauge group in Yang–Mills theory). It has Lie algebra $\text{Lie}(\mathcal{G}_0) = C^\infty(X, E)$ and here R_A is given by the familiar formula: $R_A(\xi) = -d_A \xi$.

Next, we use the connection A to determine a vector-space complement to $\text{Lie}(\mathcal{G}_0) \subset \text{Lie}(\mathcal{G})$ by horizontally lifting vector fields on X to E . This gives

$$\text{Lie}(\mathcal{G}) = \text{Lie}(\mathcal{G}_0) \oplus \text{Hor}_A \cong \text{Lie}(\mathcal{G}_0) \oplus C^\infty(X, TX)$$

where $\text{Hor}_A \cong C^\infty(X, TX)$ are the horizontal lifts to E of vector fields on X . Of course, Hor_A is not a Lie subalgebra precisely because A has curvature.

Lemma 5.2. *Given $u \in \text{Hor}_A$ its infinitesimal action at A is*

$$R_A(u) = -\iota_u F_A.$$

Proof. We switch to the principal bundle formalism. Let $P \rightarrow X$ be the principal frame bundle of E . We interpret A as an $\text{SO}(3)$ -equivariant 1-form on P with values in $\mathfrak{so}(3)$ whilst $\text{Lie}(\mathcal{G})$ is the Lie algebra of $\text{SO}(3)$ -invariant vector fields on P . Given *any* element $u \in \text{Lie}(\mathcal{G})$, the corresponding infinitesimal action on A is $R_A(u) = -L_u(A) = -d(A(u)) - \iota_u dA$. Now Hor_A is precisely those u with $A(u) = 0$. For such vectors, $\iota_u[A \wedge A] = 2[A(u), A] = 0$. It follows that $\iota_u dA = \iota_u F_A$, since $F_A = dA + \frac{1}{2}[A \wedge A]$. \square

So, given A we have an isomorphism $\text{Lie}(\mathcal{G}) \cong C^\infty(X, E) \oplus C^\infty(X, TX)$ with respect to which the infinitesimal action at A is given by

$$(7) \quad R_A(\xi, u) = -d_A \xi - \iota_u F_A.$$

Note that the action is first order in derivatives of ξ , but zeroth order in u . This means that the gauge fixing will involve a mixture of algebraic and differential conditions. The orthogonal complement to $\text{im } R_A$ is $\ker R_A^*$. Write $f_A: TX \rightarrow \Lambda^1 \otimes E$ for the bundle homomorphism $f_A(u) = \iota_u F_A$ and set $W_A = \ker f_A^* = (\text{im } f_A)^\perp$. Then $R_A^* = -(d_A^* \oplus f_A^*)$ and so $\ker R_A^* = \ker d_A^* \cap W_A$.

Lemma 5.3. *There is an L^2 -orthogonal decomposition,*

$$\Omega^1(X, E) = \text{im } R_A \oplus \ker R_A^*.$$

Proof. This follows from combining the orthogonal decomposition

$$\Omega^1(X, E) = \text{im } d_A \oplus \ker d_A^*$$

provided by elliptic theory and the pointwise orthogonal decomposition

$$\Lambda^1 \otimes E = \text{im } f_A \oplus \ker f_A^*. \quad \square$$

By gauge invariance, the Hessian D of S preserves sections of W_A . We will prove Theorem 5.1 by showing that on those sections a of W_A for which $d_A^* a = 0$, D is equal to a genuinely elliptic second order operator on $C^\infty(W_A)$.

Remark 5.4. When M_A is a multiple of the identity, the splitting $\Lambda^1 \otimes E = \text{im } f_A \oplus W_A$ coincides with a standard decomposition coming from Riemannian geometry. Write S_\pm for the spin bundles of an oriented Riemannian 4-manifold and S_\pm^m for the m^{th} symmetric power of S_\pm . Whilst S_\pm are complex vector bundles, even powers— $S_-^p \otimes S_+^q$ with $p+q$ even—carry real structures whose fixed loci are real vector bundles of half the dimension. In what follows we will only encounter these even powers, which exist globally even when X is not spin. In these cases we use $S_-^p \otimes S_+^q$ to denote the real

vector bundle, of real rank $p + q + 2$. For example, by $S_- \otimes S_+$ we mean the spin bundle isomorphic to the real cotangent bundle.

There is an orthogonal decomposition

$$\Lambda^1 \otimes \Lambda^+ \cong \Lambda^1 \oplus W,$$

where $W = S_- \otimes S_+^3$. This follows from the isomorphisms $\Lambda^1 \cong S_+ \otimes S_-$ and $\Lambda^+ \cong S_+^2$ and the irreducible decomposition

$$S_- \otimes S_+ \otimes S_+^2 \cong (S_- \otimes S_+) \oplus (S_- \otimes S_+^3).$$

Using the identification $\Phi_A: E \rightarrow \Lambda_A^+$, this splitting pulls back to $\Lambda^1 \otimes E$. One can check that this agrees with $\Lambda^1 \otimes E = \text{im } f_A \oplus W_A$ precisely when M_A is a multiple of the identity (in which case F_A is a multiple of Φ_A).

5.1.2. The principal part of the Hessian. We next describe the leading order part of the Hessian of S . We break the calculation up into a series of small steps. We begin by introducing a first order operator $\delta_A: \Omega^1(X, E) \rightarrow C^\infty(S^2E)$ defined as follows. Let e_i be a local oriented orthonormal frame for E in which $F_A = \sum F_i \otimes e_i, d_A a = \sum d_A a_i \otimes e_i$. Then

$$(\delta_A a)_{ij} \nu_A = d_A a_i \wedge F_j + F_i \wedge d_A a_j,$$

defines the (i, j) -components of $\delta_A a$.

As we will see in the next lemma, δ_A appears in the first variation of ν_A and M_A . This ultimately leads to the principal part of the Hessian having the form $\delta_A^* \circ L_A \circ \delta_A$ for some endomorphism L_A of the bundle S^2E .

Lemma 5.5. *Given an infinitesimal change $\dot{A} = a$ in a definite connection, the corresponding changes in ν_A and M_A are given by*

$$(8) \quad \dot{\nu}_A = \frac{1}{|\Lambda|} \text{Tr} \left(M_A^{-1/2} \delta_A a \right) \nu_A,$$

$$(9) \quad \dot{M}_A = \delta_A a - \frac{1}{|\Lambda|} \text{Tr} \left(M_A^{-1/2} \delta_A a \right) M_A.$$

Proof. Equation (8) is just a restatement of (4) using δ_A . To prove (9), use the definition of M in a local orthonormal frame: $M_{ij} \nu_A = F_i \wedge F_j$. Differentiating this gives $\dot{M}_A \nu_A + M_A \dot{\nu}_A = (\delta_A a) \nu_A$ which together with (8) completes the proof. \square

Remark 5.6. When M_A is a multiple of the identity, all these formulae simplify. Identifying E and Λ^+ via Φ_A , we view $\delta_A: \Omega^1(X, \Lambda^+) \rightarrow C^\infty(S^2\Lambda^+)$. When $M_A = (\Lambda/3)^2 1_E$, this is simply the composition

$$\Omega^1(X, \Lambda^+) \xrightarrow{d_\nabla} \Omega^2(X, \Lambda^+) \longrightarrow C^\infty(S^2\Lambda^+)$$

where d_∇ is the covariant exterior derivative associated to the Levi-Civita connection and the second arrow is projection $\Lambda^2 \otimes \Lambda^+ \rightarrow S^2\Lambda^+$ onto the symmetric tensor product.

Notice also that the formulae for $\dot{\nu}_A$ and \dot{M}_A simplify. For example, \dot{M}_A is simply the trace-free part of $\delta_A a$.

Another ingredient we will need to describe the Hessian is a formula for the change in $M_A^{-1/2}$. This is determined from Equation (9) for \dot{M}_A by some linear algebra which we now describe.

Lemma 5.7. *Let M be a positive definite matrix and N any symmetric matrix regarded as an infinitesimal change in M . The corresponding infinitesimal change in $M^{-1/2}$ is the unique solution G to the equation*

$$(10) \quad GM^{-1/2} + M^{-1/2}G = -M^{-1}NM^{-1}$$

Proof. The equation follows from differentiating $M^{-1/2}M^{-1/2} = M^{-1}$. To see that it has a unique solution, define an endomorphism f of the space of symmetric matrices by

$$(11) \quad H: G \mapsto -MGM^{1/2} - M^{1/2}GM.$$

Now (10) says $H(G) = N$. But one can check H is an isomorphism and so G can be recovered from N . \square

With this lemma in hand, we define a map $G_A: S^2E \rightarrow S^2E$ by setting $G_A(N)$ to be the unique solution to (10) with $M = M_A$. To give a clean formula for the infinitesimal change in $M_A^{-1/2}$, we write $L_A: S^2E \rightarrow S^2E$ for the linear map given by

$$(12) \quad L_A(N) = G_A \left(N - \frac{1}{|\Lambda|} \text{Tr} \left(M_A^{-1/2} N \right) M_A \right).$$

Lemma 5.8. *Given an infinitesimal change $\dot{A} = a$ in a definite connection A , the corresponding change in $M_A^{-1/2}$ is $L_A(\delta_A a)$.*

Proof. This comes from putting together Equation (9) for \dot{M}_A with Lemma 5.7. \square

The somewhat complicated looking formula (12) for L_A (equivalently the expression (9) for \dot{M}_A) has a simple geometric explanation. One can check that G_A is self-adjoint and negative definite, i.e., that $\text{Tr}(G_A(N)N') = \text{Tr}(NG_A(N'))$ and $\text{Tr}(G_A(N)N) < 0$ if $N \neq 0$. (One way is to prove these facts first for H defined in (11).) This means we can use $-G_A$ to define a new fibrewise innerproduct on the bundle S^2E by $\langle P, Q \rangle := -\text{Tr}(G_A(P)Q)$. Now $G_A(M_A) = -\frac{1}{2}M_A^{-1/2}$ and so $\langle M_A, M_A \rangle = \frac{|\Lambda|}{2}$. We now see that

$$N - \frac{1}{|\Lambda|} \text{Tr} \left(M_A^{-1/2} N \right) M_A = N - \frac{\langle M_A, N \rangle}{\langle M_A, M_A \rangle} M_A$$

is simply the orthogonal projection of N away from the span of M_A in the new innerproduct $\langle \cdot, \cdot \rangle$. It follows that with respect to the *standard* innerproduct $\text{Tr}(PQ)$ on S^2E , L_A is self-adjoint, negative semi-definite with kernel spanned by M_A . We record this in the following Lemma.

Lemma 5.9. *The map $L_A: S^2E \rightarrow S^2E$ defined in (12) is a self-adjoint negative semi-definite operator with kernel spanned by M_A .*

Remark 5.10. Again, this all becomes much simpler when M_A is a multiple of the identity. In this case, $G_A(N) = -27(2|\Lambda|^3)^{-1}N$ is just a multiple of N and $L_A(N)$ is $-27(2|\Lambda|^3)^{-1}$ times the projection onto the trace-free part of N .

The final ingredient we will need is an explicit formula for δ_A^* .

Lemma 5.11. *The map $C^\infty(S^2E) \rightarrow \Omega^1(X, E)$ given by*

$$N \mapsto -2 * (d_A N \wedge F_A)$$

is the L^2 -adjoint of δ_A , where here the symbol \wedge denotes the operation $(\Lambda^1 \otimes \text{End}(E)) \otimes (\Lambda^2 \otimes E) \rightarrow \Lambda^3 \otimes E$ given by tensoring the wedge product on forms with the natural action of $\text{End}(E)$ on E .

Remark 5.12. Our convention for the Hodge star is that

$$\beta \wedge * \gamma = (\beta, \gamma) \text{dvol},$$

which means that the codifferential (on any degree form on a four-manifold) is given by $d^* = - * d *$.

Proof. First note that, since $d_A F_A = 0$, $d_A N \wedge F_A = d_A(N \cdot F_A)$ where the symbol \cdot here denotes the action of N on the E -factor of $F_A \in \Lambda^2 \otimes E$. Next note that since F_A is self-dual, $-2 * d_A(N \cdot F_A) = 2d_A^*(N \cdot F_A)$. We now see that

$$\int_X (-2 * (d_A N \wedge F_A), a) \nu_A = \int_X (N \cdot F_A, 2d_A a) \nu_A.$$

Write $d_A a = \sum \alpha_i \otimes e_i$, then

$$(N \cdot F_A, 2d_A a) = 2 \sum_{i,j,k} (N_{ij} F_i \otimes e_j, \alpha_k \otimes e_k) = 2 \sum_{i,j} N_{ij} (F_i, \alpha_j) = \text{Tr}(N \delta_A a)$$

which completes the proof. □

We can now give a precise description of the leading order part of the Hessian of S . As a warm up, and to motivate the general case, we first compute the Hessian of S at a point A for which M_A is a multiple of the identity.

Proposition 5.13. *When M_A is a multiple of the identity (so that g_A is anti-self-dual Einstein and $S(A) = 2\chi + 3\tau$ attains a global maximum), the Hessian of S is given by*

$$(13) \quad D = \frac{|\Lambda|}{12\pi^2} \delta_A^* \circ L_A \circ \delta_A.$$

Proof. Recall from the proof of Theorem 2.9 that the derivative of S in the direction a is given by

$$dS(a) = \frac{\Lambda}{6\pi^2} \int_X (\Phi_A, d_A a) \nu_A$$

where $\Phi_A = \pm F_A \circ M_A^{-1/2}$ (with sign corresponding to that of the definite connection). Writing the integrand in terms of a local frame for E , this reads

$$dS(a) = \frac{|\Lambda|}{6\pi^2} \int \sum_{i,j} (M_A^{-1/2})_{ij} F_i \wedge d_A a_j.$$

Differentiating with respect to A in the direction b gives

$$\begin{aligned} d^2 S(a, b) &= \frac{|\Lambda|}{6\pi^2} \int \left[\sum_{i,j} L_{ij}(\delta_A b) F_i \wedge d_A a_j + \frac{3}{|\Lambda|} \sum_i (d_A b_i \wedge d_A a_i + F_i[b, a]_i) \right], \end{aligned}$$

where in the last term we have used the fact that M_A is a multiple of the identity. It is now a matter of simple integration by parts in the last term to see that it vanishes. \square

In the general case, a similar calculation shows that the Hessian has the same principal part as in the anti-self-dual Einstein case.

Proposition 5.14. *At an arbitrary definite connection A , the principal part of the Hessian of S is*

$$(14) \quad \hat{D} = \frac{|\Lambda|}{12\pi^2} \delta_A^* \circ L_A \circ \delta_A.$$

Proof. Recall from the proof of Theorem 2.9 that the derivative of S in the direction a is given by

$$(15) \quad dS(a) = \frac{\Lambda}{6\pi^2} \int_X (d_A^* \Phi_A, a) \nu_A$$

where $\Phi_A = \pm F_A \circ M_A^{-1/2}$ (with sign corresponding to that of the definite connection). We now differentiate this with respect to A in the direction b .

The L^2 -innerproduct here depends on A , but the infinitesimal change in metric is *first*-order in b . Since we are only interested in the principal part, we can ignore this. The only piece which is second-order in b comes from $d_A^* \Phi_A$. In the notation of Lemma 5.11, one can check that

$$d_A^* \Phi_A = \pm \frac{1}{2} \delta_A^* (M_A^{-1/2}).$$

Again, the dependence of δ_A^* on A is first order. This means that we need only ultimately consider the linearisation of $A \mapsto M_A^{-1/2}$ in the direction of b (which will then be differentiated again by δ_A^* , becoming second order). The proposition now follows from Lemma 5.8. \square

We next turn to the symbol of D . Let η be a nonzero covector and write

$$\sigma(D, \eta): \Lambda^1 \otimes E \rightarrow \Lambda^1 \otimes E$$

for the symbol of D in the direction η .

Proposition 5.15. *The symbol $\sigma(D, \eta)$ is negative semi-definite with kernel equal to*

$$\ker \sigma(D, \eta) = \{\eta \otimes e + \iota_u F_A : e \in E, u \in TX\}$$

which is the span of the images of the symbols of d_A and f_A . (Recall $f_A: TX \rightarrow \Lambda^1 \otimes E$ is the map $f_A(u) = \iota_u F_A$.)

Proof. Write δ_η for the symbol of δ_A in the direction η . Then by Proposition 5.14, $\sigma(D, \eta) = (|\Lambda|/12\pi^2)\delta_\eta^* \circ L_A \circ \delta_\eta$ and this is negative semi-definite by Lemma 5.9. Moreover, the kernel of $\sigma(D, \eta)$ is precisely those a for which $\delta_\eta(a)$ is a multiple of M_A (possibly zero multiple).

Since the action is gauge invariant, $D \circ R_A = 0$ or, in other words, $D \circ d_A = 0 = D \circ f_A$. This means that the images of the symbols of d_A and f_A must lie in the kernel. Alternatively, we can check directly that

$$\sigma(D, \eta)(\eta \otimes e + \iota_u F_A) = 0.$$

The term $\eta \otimes e$ is killed immediately since Σ_η begins by wedging with η . For the second term, we work in a local frame for E in which $F_A = \sum F_i \otimes e_i$. We have $\Sigma_\eta(\iota_u F_A) = (\eta \wedge \iota_u F_i, F_j) + (\eta \wedge \iota_u F_j, F_i)$. Now the 5-form $\eta \wedge F_i \wedge F_j$ is necessarily zero. It follows that

$$0 = \iota_u(\eta \wedge F_i \wedge F_j) = \iota_u \eta \wedge F_i \wedge F_j - \eta \wedge \iota_u F_i \wedge F_j - \eta \wedge F_i \wedge \iota_u F_j.$$

Dividing by the volume form gives that $\delta_\eta(\iota_u F_A) = \eta(u)M_A$ is a multiple of M_A .

It remains to check that we have accounted for the entire kernel, i.e., that it has dimension 7. The map $\delta_\eta: \Lambda^1 \otimes E \rightarrow S^2 E$ is surjective. Indeed it is, up to a factor of 2, the composition of the surjective operations: wedge with η followed by self-dual projection $\Lambda^1 \otimes E \rightarrow \Lambda^+ \otimes E$; identification of $\Lambda^+ \otimes E$ and $E \otimes E$ via F_A and finally projection $E \otimes E \rightarrow S^2 E$. This means that those a sent to a multiple of M_A by δ_η —the kernel of $\sigma(D, \eta)$ —form a 7-dimensional subspace of $\Lambda^1 \otimes E$. \square

Proof of Theorem 5.1. We now prove that D is elliptic modulo gauge, with finitely many positive eigenvalues. Recall that if $\xi \in C^\infty(X, E)$ and $u \in C^\infty(X, TX)$, the infinitesimal gauge action $R_A(\xi, u) = -d_A \xi - \iota_u F_A$ is first order in ξ and zeroth order in u . To fix for this we will treat the two parts separately. Write $W_A \leq \Lambda^1 \otimes E$ for the orthogonal complement of the image map f_A (where $f_A(u) = \iota_u F_A$), i.e., $W_A = \ker f_A^*$. Write $\Pi: \Lambda^1 \otimes E \rightarrow W_A$ for the orthogonal projection onto W_A . Since $D \circ f_A = 0$ and D is self-adjoint we see that D restricts to a map $C^\infty(W_A) \rightarrow C^\infty(W_A)$. We will consider the gauge fixed operator $D': C^\infty(W_A) \rightarrow C^\infty(W_A)$ given by

$$(16) \quad D'(\Pi a) = D(\Pi a) - \Pi(d_A d_A^* \Pi a).$$

On the orthogonal complement $\ker R_A^* = \ker d_A^* \cap W_A$ of the gauge group action, $D' = D$. With Proposition 5.15 in hand, it is straight forward to

check that D' is genuinely elliptic. Its symbol in the direction of a unit length covector η is

$$\sigma(D', \eta) = \frac{|\Lambda|}{12\pi^2} \delta_\eta^* L_A \delta_\eta - \Pi p_\eta \Pi,$$

where $p_\eta: \Lambda^1 \otimes E \rightarrow \Lambda^1 \otimes E$ is given by tensoring 1_E with the orthogonal projection from Λ^1 onto the span of η .

Suppose that $\sigma(D', \eta)(a) = 0$ for some $a \in W_A$. The symbol of D' is a sum of negative semi-definite operators and so a is in the kernel of both terms. By Proposition 5.15, it follows that $a = \eta \otimes e + f_A(u)$ for some $e \in E$ and $u \in TX$, but now $p_\eta(a) = 0$ forces $e = 0$ and hence $a \in \text{im } f_A$. But since $a \in W_A$ is also orthogonal to $\text{im } f_A$, this forces $a = 0$ and so $\sigma(D', \eta)$ is negative definite and in particular an isomorphism.

We have shown that D' is elliptic and so it has a complete basis of eigenvectors. It follows that the restriction of D to $\ker R_A^* = \ker d_A^* \cap W_A$ also has a complete basis of eigenvectors (a sub-collection of the eigenvectors of D'). It remains to show that D' has finitely many positive eigenvalues, but this follows from the fact that its symbol is negative definite. In our case we can see this directly: replacing D by $\Pi \circ \hat{D} = \frac{|\Lambda|}{12\pi^2} \Pi \delta_A^* L_A \delta_A$ in (16) gives an elliptic operator \hat{D}' (with the same symbol) which is negative semi-definite and so has nonpositive spectrum. Now D and \hat{D} can be joined by a smooth path, leaving the principal part alone and interpolating between the lower order pieces. Using this path in place of D in (16) gives a path of elliptic operators (all with the same symbol). Along this path the spectrum—a discrete subset of \mathbb{R} —varies continuously. Since it is nonpositive at one end of the path, continuity implies only finitely many eigenvalues can become positive at the other end. \square

5.2. The Hessian as a spin Laplacian. When M_A is a multiple of the identity, so that g_A is anti-self-dual Einstein, the action $S(A)$ attains a global maximum and so the Hessian is nonpositive. In this section we will show that in fact the Hessian is invertible modulo gauge. This implies that anti-self-dual Einstein metrics with nonzero scalar curvature are locally rigid modulo diffeomorphisms.

This local rigidity was originally proved for positive scalar curvature by LeBrun [24] and negative scalar curvature by Horan [19]. In the traditional twistor-theoretic point of view they employ, one uses the Penrose transform to identify infinitesimal anti-self-dual Einstein deformations with the certain twisted harmonic spinors. One must then show that the only such harmonic spinors are zero. Our new formulation of the problem gives an alternative way to interpret the appearance of this Dirac operator: modulo gauge, the Hessian is the corresponding spin Laplacian.

When discussing gauge-fixing, we defined a sub-bundle W_A of $\Lambda^1 \otimes E$ which was orthogonal to the action $\text{im } f_A$ of tangent vectors. Recall from Remark 5.4 that when M_A is a multiple of the identity, Φ_A identifies W_A

with the twisted spin bundle $S_- \otimes S_+^3 \subset \Lambda^1 \otimes \Lambda^+$. This bundle carries a Dirac operator defined by coupling to the Levi-Civita connection on S_+^3 , which we denote

$$\not\partial: C^\infty(S_- \otimes S_+^3) \rightarrow C^\infty(S_+ \otimes S_+^3)$$

We will show that D is essentially $\not\partial^* \not\partial$. To compare these operators, we use Φ_A to identify $\Lambda^1 \otimes E \rightarrow \Lambda^1 \otimes \Lambda^+$. Given $a \in \Lambda^1 \otimes E$, we write \hat{a} for its image in $\Lambda^1 \otimes \Lambda^+$.

Proposition 5.16. *Let A be a definite connection with M_A a multiple of the identity. Let $a \in \Omega^1(X, E)$ be an infinitesimal deformation of A orthogonal to the gauge group action, i.e., $a \in C^\infty(W_A)$ with $d_A^* a = 0$. Then*

$$\widehat{D}(a) = -c \not\partial^* \not\partial(\hat{a})$$

for some positive constant c , which depends only on $|\Lambda|$.

Proof. Remark (5.4) explains that if $a \in W_A$ then $\hat{a} \in C^\infty(S_- \otimes S_+^3)$ lies in the domain of the Dirac operator. Next, since that Φ_A identifies A with the Levi-Civita connection ∇ , we see that $d_\nabla^* \hat{a} = 0$.

Now, Equation (13) for the Hessian of $S(A)$ when M_A is a multiple of the identity shows that the Hessian is, up to multiplication by a negative constant factor, the following composition

$$(17) \quad \Omega^1(X, \Lambda^+) \xrightarrow{d_\nabla} \Omega^2(X, \Lambda^+) \xrightarrow{p} \Omega^2(X, \Lambda^+) \xrightarrow{d_\nabla^*} \Omega^1(X, \Lambda^+)$$

where p is orthogonal projection from $\Lambda^2 \otimes \Lambda^+$ onto the subspace $S_0^2 \Lambda^+ \subset \Lambda^2 \otimes \Lambda^+$ of symmetric trace-free endomorphisms of Λ^+ .

Meanwhile, the Dirac operator $\not\partial$ can also be expressed in terms of d_∇ and d_∇^* . The range of $\not\partial$ is $S_+ \otimes S_+^3 \cong S_+^4 \oplus S_+^2$. Meanwhile the range of d_∇ is

$$\Lambda^2 \otimes \Lambda^+ \cong (S_-^2 \otimes S_+^2) \oplus (S_+^2 \otimes S_+^2) \cong (S_-^2 \otimes S_+^2) \oplus S_+^4 \oplus \mathbb{R} \oplus S_+^2$$

and that of d_∇^* is $\Lambda^+ \cong S_+^2$. The Dirac operator on $S_- \otimes S_+^3$ is built from the S_+^4 component of d_∇ and the whole of d_∇^* .

Now projection of a section of $\Lambda^2 \otimes \Lambda^+$ onto the S_+^4 component is precisely the above projection p . So when $d_\nabla^* \hat{a} = 0$, $\not\partial \hat{a} = p(d_\nabla \hat{a})$. It follows that $\not\partial^* \not\partial \hat{a} = (d_\nabla^* \circ p \circ d_\nabla) \hat{a}$ which is, up to a negative factor, $\widehat{D}(a)$. \square

Corollary 5.17 (cf. LeBrun [24] when $\Lambda > 0$ and Horan [19] when $\Lambda < 0$). *When M_A is a multiple of the identity, the Hessian of S at A is negative definite modulo gauge. It follows that anti-self-dual Einstein metrics with nonzero scalar curvature are locally rigid modulo diffeomorphism.*

Proof. By Proposition 5.16 it suffices to show that when the metric is anti-self-dual, Einstein with nonzero scalar curvature, the spin Laplacian $\not\partial^* \not\partial$ is invertible on sections of $S_- \otimes S_+^3$. This was proved for positive scalar curvature by LeBrun [24] and negative scalar curvature by Horan [19]. \square

At an arbitrary point A (for which M_A is no longer a multiple of the identity) there is a similar story for the principal part (14) of the Hessian. First, it is convenient to twist the tangent vector $a \in \Lambda^1 \otimes E$ by $\sqrt{M_A}$. Define the “twisted tangent” $\alpha \in \Lambda^1 \otimes E$ by $a = \sqrt{M_A}\alpha$. Explicitly, if $a = \sum a_i \otimes e_i$, then

$$\alpha = \sum (M_A^{-1/2})_{ij} a_j \otimes e_i.$$

One then finds that the story for the principal part of the Hessian on the space of twisted tangents α repeats the anti-self-dual Einstein case almost exactly. The only additional difference is that in this case instead of the projection p on the trace-free part one has to insert a more complicated operator on $S_0^2\Lambda^+ \cong S_+^4$ into (17) that depends nontrivially on M_A . One then finds that the principal part of the gauge-fixed operator can be written in a form analogous to Proposition 5.16, but with an additional endomorphism $Q: S_+ \otimes S_+^3 \rightarrow S_+ \otimes S_+^3$ (defined in terms of M_A) inserted:

$$\begin{aligned} C^\infty(S_- \otimes S_+^3) &\xrightarrow{\phi} C^\infty(S_+ \otimes S_+^3) \xrightarrow{Q} C^\infty(S_+ \otimes S_+^3) \xrightarrow{Q^*} C^\infty(S_+ \otimes S_+^3) \\ &\xrightarrow{\phi^*} C^\infty(S_- \otimes S_+^3). \end{aligned}$$

This shows that in general the principal part of the gauge-fixed Hessian is still given by the square of the Dirac operator, but with a nontrivial “twisting matrix” Q^*Q inserted. The main complication in the general case is that there are also other terms, apart from the principal part, which are absent in the anti-self-dual Einstein case.

5.3. Short time existence of the gradient flow of S . We now turn to the gradient flow of S , which is given by

$$(18) \quad \frac{\partial A}{\partial t} = \frac{\Lambda}{6\pi^2} d_A^* \Phi_A.$$

(The actual constant factor appearing here is unimportant as long as the sign agrees with that of the definite connection.) This is the analogue of the Yang–Mills flow in our theory.

The fact that the Hessian of S is elliptic modulo gauge implies that the gradient flow (18) is parabolic modulo gauge and hence exists for short time. This is standard in the study of geometric flows, going by the name of “de Turck’s trick” in the study of Ricci flow [3]; see also [4] for a similar proof of the short-time existence of Yang–Mills flow. Accordingly we are content to merely sketch the details.

Theorem 5.18. *Let A_0 be a definite connection. Then there is a short-time solution $A(t)$ to the gradient flow (18) with $A(t) = A_0$.*

Sketch of proof. We will first consider a gauge-equivalent flow:

$$(19) \quad \frac{\partial A}{\partial t} = \frac{\Lambda}{6\pi^2} d_A^* \Phi_A - d_A d_A^*(A - A_0) - f_A(u_A).$$

Here $u_A \in C^\infty(X, TX)$ is the unique vector field which solves the equation

$$(20) \quad f_A(u_A) = (1 - \Pi) \left(\frac{\Lambda}{6\pi^2} d_A^* \Phi_{\hat{A}} - d_{\hat{A}} d_A^*(A - A_0) \right).$$

(Recall that Π is the pointwise orthogonal projection of $\Lambda^1 \otimes E$ onto W_A and so $1 - \Pi$ is projection onto $\text{im } f_A$.) By definition of u_A this flow is tangent to W_A . We also ask that $A(0) = A_0$.

At $t = 0$, the principal part of the linearisation of this flow is the operator D' on $C^\infty(W_A)$ of (16). To see this note that the first two terms in (19) contribute $D - d_A d_A^*$ (since at $t = 0$ any effect on linearising $d_A d_A^*$ is then evaluated on $A(0) - A_0 = 0$). Moreover, f_A is first order in A and so the contribution to the principal part coming from $f_A(u_A)$ is determined purely by the change in u_A . From the definition (20) of u_A this cancels exactly $(1 - \Pi)(D - d_A d_A^*)$, leaving $D' = D - \Pi d_A d_A^*$ as the principal part as claimed.

As we have seen, the symbol of D' is negative definite. It follows from the standard theory of parabolic PDE that (19) has a solution starting, which exists at least for short time. Write A' for the solution to (19) with $A'(0) = A_0$. We now consider the path of gauge transforms $g(t) \in \mathcal{G}$ which are generated by $-d_{A'}^*(A' - A_0) - u_{A'}$, with $g(0)$ the identity. (Here, we have implicitly used A' to horizontally lift $u_{A'}$ and so consider it as an infinitesimal gauge transformation). By gauge invariance of $d_A^* \Phi_A$ and the formula (7) for the infinitesimal action of $-d_{A'}^*(A' - A_0) - u_{A'}$, it follows that $A(t) = g(t)^* A'(t)$ solves the original gradient flow (18) with $A(0) = A_0$. \square

References

- [1] ATIYAH, M. F.; HITCHIN, N. J.; SINGER, I. M. Self-duality in four-dimensional Riemannian geometry. *Proc. Roy. Soc. London. Ser. A.* **362** (1978), no. 1711, 425–461. [MR0506229](#) (80d:53023), [Zbl 0389.53011](#), doi: [10.1098/rspa.1978.0143](#).
- [2] CAMPANA, F. On twistor spaces of the class \mathcal{C} . *J. Differential Geom.* **33** (1991), no. 2, 541–549. [MR1094468](#) (92g:32059), [Zbl 0694.32017](#).
- [3] DETURCK, DENNIS M. Deforming metrics in the direction of their Ricci tensors. *J. Differential Geom.* **18** (1983), no. 1, 157–162. [MR0697987](#) (85j:53050), [Zbl 0517.53044](#).
- [4] DONALDSON, S. K.; KRONHEIMER, P. B. The geometry of four-manifolds. Oxford Mathematical Monographs. Oxford Science Publications. *The Clarendon Press Oxford University Press, New York*, 1990. x+440 pp. ISBN: 0-19-853553-8. [MR1079726](#) (92a:57036), [Zbl 0904.57001](#).
- [5] EDDINGTON, A. S. A generalisation of Weyl’s theory of the electromagnetic and gravitational fields. *London R. S. Proc.*, **99** (1921), 104–122. [Zbl 48.1040.04](#) doi: [10.1098/rspa.1921.0027](#).
- [6] EELLS, JAMES; SALAMON, SIMON. Constructions twistorielles des applications harmoniques. *C. R. Acad. Sci. Paris Sér. I Math.* **296** (1983), no. 15, 685–687. [MR0705691](#) (85a:58023), [Zbl 0531.58020](#).
- [7] FINE, JOEL. A gauge theoretic approach to the anti-self-dual Einstein equations. Preprint, 2011. [arXiv:1111.5005](#).

- [8] FINE, JOEL; PANOV, DMITRI. Symplectic Calabi–Yau manifolds, minimal surfaces and the hyperbolic geometry of the conifold. *J. Differential Geom.* **82** (2009), no. 1, 155–205. [MR2504773](#) (2011b:53206), [Zbl 1177.32014](#), [arXiv:0802.3648](#).
- [9] FINE, JOEL; PANOV, DMITRI. Hyperbolic geometry and non-Kähler manifolds with trivial canonical bundle. *Geom. Topol.* **14** (2010), no. 3, 1723–1763. [MR2679581](#) (2011k:32036), [Zbl 1214.53058](#), [arXiv:0905.3237](#), doi: [10.2140/gt.2010.14.1723](#).
- [10] FINE, JOEL; PANOV, DMITRI. The diversity of symplectic Calabi–Yau 6-manifolds. *J. Topology* **6** (2013), no. 3, 644–658. [MR3100885](#), [arXiv:1108.5944](#), doi: [10.1112/jtopol/jtt011](#).
- [11] GROMOV, M.; THURSTON, W. Pinching constants for hyperbolic manifolds. *Invent. Math.* **89** (1987), no. 1, 1–12. [MR0892185](#) (88e:53058), [Zbl 0646.53037](#), doi: [10.1007/BF01404671](#).
- [12] GURSKY, MATTHEW J. Four-manifolds with $\delta W^+ = 0$ and Einstein constants of the sphere. *Math. Ann.* **318** (2000), no. 3, 417–431. [MR1800764](#) (2002c:53074), [Zbl 1034.53032](#), doi: [10.1007/s002080000130](#).
- [13] GURSKY, MATTHEW J.; LEBRUN, CLAUDE. On Einstein manifolds of positive sectional curvature. *Ann. Global Anal. Geom.* **17** (1999), no. 4, 315–328. [MR1705915](#) (2000i:53068), [Zbl 0967.53029](#), [arXiv:math/9807055](#), doi: [10.1023/A:1006597912184](#).
- [14] HIRZEBRUCH, F. The signature of ramified coverings. *Global Analysis (Papers in Honor of K. Kodaira)*, 253–265. *Univ. Tokyo Press, Tokyo*, 1969. [MR0258060](#) (41 #2707), [Zbl 0208.51802](#).
- [15] HITCHIN, NIGEL. Compact four-dimensional Einstein manifolds. *J. Differential Geometry* **9** (1974), 435–441. [MR0350657](#) (50 #3149), [Zbl 0281.53039](#). <http://projecteuclid.org/euclid.jdg/1214432419>.
- [16] HITCHIN, N. J. Kählerian twistor spaces. *Proc. London Math. Soc.* (3) **43** (1981), no. 1, 133–150. [MR0623721](#) (84b:32014), [Zbl 0474.14024](#), doi: [10.1112/plms/s3-43.1.133](#).
- [17] HITCHIN, NIGEL. The geometry of three-forms in six dimensions. *J. Differential Geom.* **55** (2000), no. 3, 547–576. [MR1863733](#) (2002m:53070), [Zbl 1036.53042](#).
- [18] HITCHIN, NIGEL. Stable forms and special metrics. *Global differential geometry: the mathematical legacy of Alfred Gray (Bilbao, 2000)*, 70–89. *Contemp. Math.*, 288, *Amer. Math. Soc., Providence, RI*, 2001. [MR1871001](#) (2003f:53065), [Zbl 1004.53034](#), [arXiv:math/0107101](#), doi: [10.1090/conm/288](#).
- [19] HORAN, ROBIN. A rigidity theorem for quaternionic Kähler manifolds. *Differential Geom. Appl.* **6** (1996), no. 2, 189–196. [MR1395029](#) (97c:53068), [Zbl 0846.32022](#), doi: [10.1016/0926-2245\(96\)89149-2](#).
- [20] ISKOVSKIKH, V. A. Fano threefolds. I. *Izv. Akad. Nauk SSSR Ser. Mat.* **41** (1977), no. 3, 516–562, 717. [MR0463151](#) (80c:14023a), [Zbl 0363.14010](#), doi: [10.1070/IM1977v011n03ABEH001733](#).
- [21] KOLLÁR, JÁNOS. Projectivity of complete moduli. *J. Differential Geom.* **32** (1990), no. 1, 235–268. [MR1064874](#) (92e:14008), [Zbl 0684.14002](#), <http://projecteuclid.org/euclid.jdg/1214445046>.
- [22] KRASNOV, KIRILL. Gravity as a diffeomorphism-invariant gauge theory. *Phys. Rev. D; Physical Review D* **84** (2011). [arXiv:1101.4788](#). doi: [10.1103/PhysRevD.84.024034](#).
- [23] KRASNOV, KIRILL. Pure connection action principle for general relativity. *Phys. Rev. Lett.* **106** (2011). doi: [10.1103/PhysRevLett.106.251103](#).
- [24] LEBRUN, CLAUDE. A rigidity theorem for quaternionic-Kähler manifolds. *Proc. Amer. Math. Soc.* **103** (1988), no. 4, 1205–1208. [MR0955010](#) (89h:53105), [Zbl 0655.53056](#), doi: [10.1016/0926-2245\(96\)89149-2](#).
- [25] MCDUFF, DUSA. The structure of rational and ruled symplectic 4-manifolds. *J. Amer. Math. Soc.* **3** (1990), no. 3, 679–712. [MR1049697](#) (91k:58042), [Zbl 0723.53019](#), doi: [10.1090/S0894-0347-1990-1049697-8](#).

- [26] MOISHEZON, B. G. On n -dimensional compact complex manifolds having n algebraically independent meromorphic functions. I. *Izv. Akad. Nauk SSSR Ser. Mat.* **30** (1966), 133–174. [MR0216522](#) (35 #7355a), [Zbl 0161.17802](#).
- [27] PAGE, D. A compact rotating gravitational instanton. *Phys. Lett.* **79** (1979), 235 – 238. doi: [10.1016/0370-2693\(78\)90231-9](#).
- [28] RICHARD, THOMAS; SESHADRI, HARISH. Positive isotropic curvature and self-duality in dimension 4. Preprint, 2013. [arXiv:1311.5256](#).
- [29] THORPE, JOHN A. Some remarks on the Gauss–Bonnet integral. *J. Math. Mech.* **18** (1969), 779–786. [MR0256307](#) (41 #963), [Zbl 0183.50503](#).
- [30] M. Verbitsky. Rational curves and special metrics on twistor spaces. To appear in *Geometry and Topology*, 2012. [arXiv:1210.6725](#).
- [31] WILKING, BURKHARD. Nonnegatively and positively curved manifolds. *Surveys in differential geometry. XI*, 25–62. *Surv. Differ. Geom.*, 11, *Int. Press, Somerville, MA*, 2007. [MR2408263](#) (2009e:53048), [Zbl 1162.53026](#), [arXiv:0707.3091](#), doi: [10.4310/SDG.2006.v11.n1.a3](#).

(Joel Fine) DÉPARTEMENT DE MATHÉMATIQUE, UNIVERSITÉ LIBRE DE BRUXELLES, BRUXELLES 1050, BELGIUM
Joel.Fine@ulb.ac.be

(Kirill Krasnov) SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY PARK, NOTTINGHAM, NG7 2RD, UNITED KINGDOM
Kirill.Krasnov@nottingham.ac.uk

(Dmitri Panov) DEPARTMENT OF MATHEMATICS, KING'S COLLEGE, STRAND, LONDON WC2R 2LS, UNITED KINGDOM
Dmitri.Panov@kcl.ac.uk

This paper is available via <http://nyjm.albany.edu/j/2014/20-18.html>.