

Differential Borel equivariant cohomology via connections

Corbett Redden

ABSTRACT. For a compact Lie group acting on a smooth manifold, we define the differential cohomology of a certain quotient stack involving principal bundles with connection. This produces differential equivariant cohomology groups that map to the Cartan–Weil equivariant forms and to Borel’s equivariant integral cohomology. We show the Chern–Weil homomorphism for equivariant vector bundles with connection naturally factors through differential equivariant cohomology.

CONTENTS

1. Introduction	441
2. A cochain model	445
3. Differential cohomology of the differential quotient stack	449
4. Constructions of maps between stacks	455
5. Equivariant Chern–Weil theory	466
6. Details of the simplicial sheaf construction	472
Appendix A. Equivariant de Rham theory	478
References	485

1. Introduction

Differential cohomology, also known as Cheeger–Simons differential characters [CheeS] or Deligne cohomology [Bry], is a contravariant functor \check{H}^* from the category of manifolds to graded abelian groups. It sits in the


Received February 23, 2016.

2010 *Mathematics Subject Classification.* 53C08, 55N91, 53C05.

Key words and phrases. Differential cohomology, equivariant cohomology, Chern–Weil.

The author thanks the Hausdorff Institute for Mathematics for its support and hospitality during his visit.

character diagram



where the two diagonals are short exact sequences, B is the Bockstein homomorphism, and the subscript \mathbb{Z} denotes closed forms with \mathbb{Z} -periods. Because it captures both the torsion of integral cohomology and the local structure of differential forms, differential cohomology has proven to be useful in a number of contexts. In particular, the Chern–Weil homomorphism factors through \check{H}^* , making it a natural home for secondary invariants of principal bundles with connection. Differential cohomology has now been generalized in several directions, leading to the notion of a “differential extension” of a generalized cohomology theory.

The purpose of this paper is to construct a differential extension of Borel’s equivariant integral cohomology. This was also recently done by Kübel–Thom [KT], and our functors fit into the same short exact sequences. Assume that G is a compact Lie group acting smoothly on a manifold M . The equivariant cohomology of M , with coefficients in an abelian group A , is the cohomology of the homotopy quotient

$$H_G^*(M; A) := H^*(EG \times_G M; A).$$


The complex of equivariant forms $(\Omega_G(M), d_G)$ is given by the Weil model

$$(S\mathfrak{g}^* \otimes \Lambda\mathfrak{g}^* \otimes \Omega(M))_{\text{hor}}^G,$$

whose cohomology is naturally isomorphic to $H_G^*(M; \mathbb{R})$. One can equivalently use the Cartan model $(S\mathfrak{g}^* \otimes \Omega(M))^G$ in place of the Weil model, as the two complexes are isomorphic (not merely quasi-isomorphic). We summarize the paper’s main results in two theorems below. Theorem 1.1 contains the important user-friendly properties of \check{H}_G^* . It is simply Propositions 2.3, 4.4, and 4.16 combined into a single statement. In fact, we show in Proposition 4.18 that the functor \check{H}_G^* is characterized uniquely by properties A and C. Theorem 5.3 states that the equivariant Chern–Weil homomorphism, with equivariant forms defined by the equivariant curvature Ω_G [BeV], factors through \check{H}_G^* . For convenience, we restate it here as Theorem 1.2 in the specific case of equivariant complex vector bundles.

Theorem 1.1. *For any compact Lie group G , there exists a contravariant functor $\check{H}_G^*(-)$, from the category of smooth G -manifolds to the category of graded abelian groups, satisfying the following.*

(A) *There exist natural homomorphisms forming the following character diagram.*



(B) *For $G_1 \xrightarrow{\phi} G_2$ a Lie group homomorphism, a smooth map $M \xrightarrow{F} N$ that is ϕ -equivariant induces homomorphisms*

$$\check{H}_{G_2}^*(N) \xrightarrow{F^*_{\phi}} \check{H}_{G_1}^*(M)$$

compatible with the character diagrams.

(C) *A principal G -bundle with connection $(P, \Theta) \rightarrow M$ induces homomorphisms*

$$\check{H}_G^*(P) \xrightarrow{\Theta^*} \check{H}^*(M)$$

compatible with the character diagrams.

Theorem 1.2. *Let $(V, \nabla) \rightarrow M$ be a G -equivariant Hermitian vector bundle with G -invariant connection. There exist natural classes $\check{c}_k(\Theta_G) \in \check{H}_G^{2k}(M)$ satisfying*

- (1) $I(\check{c}_k(\Theta_G)) = c_k(EG \times_G V \rightarrow EG \times_G M) \in H_G^{2k}(M; \mathbb{Z})$.
- (2) $R(\check{c}_k(\Theta_G)) = c_k(\Omega_G) \in \Omega_G^{2k}(M)$.

The paper is organized as follows. Appendix A, which appears at the end of the paper, contains a review of principal bundles with connection and the equivariant de Rham complex $\Omega_G(M)$. Readers unfamiliar with the Weil model may find it helpful to begin here. While this appendix is expository and should probably be replaced by a citation to [GS], we use it to introduce notation and to emphasize the relationship between $W(\mathfrak{g})$ and differential forms on principal bundles with connection.

In Section 2 we define $\check{H}_G^*(M)$ using the Hopkins–Singer [HS] cochain model for $\check{H}^*(EG \times_G M)$, but with $\Omega(EG \times_G M)$ replaced by $\Omega_G(M)$. The short exact sequences and character diagram follow immediately. The advantage of this construction is that one need not leave the world of cochain complexes and topological spaces. When giving constructions such as the

Chern–Weil refinement, however, one is forced to play the tedious game of choosing classifying maps and checking that everything is well-defined.

To emphasize the geometric nature of our constructions, we begin working with the differential Borel quotient stack $\mathcal{E}_\nabla G \times_G M$ in Section 3. This is defined as a contravariant functor that associates to any test manifold X the groupoid of principal G -bundles with connection $(P, \Theta) \rightarrow X$, together with an equivariant map $P \rightarrow M$. In [FreH], Freed–Hopkins show that the natural map

$$\Omega_G^n(M) \xrightarrow{\cong} \Omega^n(\mathcal{E}_\nabla G \times_G M)$$

is an isomorphism, and we proceed to give the second definition

$$\check{H}_G^n(M) := \check{H}^n(\mathcal{E}_\nabla G \times_G M).$$

The details of this definition involve simplicial sheaves and are contained in Section 6. The virtue is that maps between various stacks induce homomorphisms in differential cohomology, as we demonstrate in Section 4.

For example, the homomorphism (B) from Theorem 1.1 is induced by a natural map

$$\mathcal{E}_\nabla G_1 \times_{G_1} M \xrightarrow{F_\phi} \mathcal{E}_\nabla G_2 \times_{G_2} N,$$

defined using the associated bundle construction. Similarly, if $(P, \Theta) \rightarrow M$ is a principal G -bundle with connection, the pullback gives a natural map

$$M \xrightarrow{\Theta} \mathcal{E}_\nabla G \times_G P,$$

and this induces the homomorphism in C. For $(Q, \Theta) \rightarrow M$ a G -equivariant principal K -bundle with G -invariant connection, the refined Chern–Weil homomorphism is induced by a natural map

$$\mathcal{E}_\nabla G \times_G M \longrightarrow \mathcal{B}_\nabla K,$$

as described in Section 5. Given a G -bundle with connection mapping to M , we pull back (Q, Θ) to form a $G \times K$ -bundle with connection, and then we quotient by G . The equivariant extensions of the Chern forms appear when forming the connection on the pullback of (Q, Θ) . We conclude Section 5 with a discussion of $\check{H}_G^n(M)$ as differential characters on $\mathcal{E}_\nabla G \times_G M$, and we show $\check{H}_G^2(M)$ is naturally equivalent to isomorphism classes of G -equivariant principal \mathbb{R}/\mathbb{Z} -bundles on M with invariant connection.

A definition of differential equivariant cohomology was previously given by Gomi [Gom], but it did not fully incorporate $\Omega_G(M)$, leading to groups that are not isomorphic to the ones in this paper. See Kübel–Thom [KT] for a more detailed discussion. In fact, many of our results can also be found in their paper, including the short exact sequences and the refined equivariant Chern–Weil homomorphism. However, the constructions and methods are different. The two definitions in this paper involve the topological space $EG \times_G M$ and the stack $\mathcal{E}_\nabla G \times_G M$, and most homomorphisms are induced by bundle constructions. On the other hand, Kübel–Thom use the simplicial manifold $\{G^\bullet \times M\}$ as the model for the homotopy quotient. We believe

that having multiple viewpoints will prove to be useful, just as it has for ordinary differential cohomology.

2. A cochain model

Our first definition of $\check{H}_G^*(-)$ is given by using the Hopkins–Singer construction for $\check{H}^*(EG \times_G M)$, with $\Omega(EG \times_G M)$ replaced by the Weil model of equivariant forms $\Omega_G(M)$. The definition, short exact sequences, and ring structure all follow immediately from the same arguments given in Sections 2 and 3 of the Hopkins–Singer paper [HS], so we will keep the proofs brief. (In fact, [HS] also contains our construction for the case where $M = \text{pt.}$) Readers unfamiliar with the Weil model of equivariant forms $\Omega_G(M)$ can find a full exposition in Appendix A. As noted in Section A.5, the Weil model may be replaced, via the Mathai–Quillen isomorphism, by the isomorphic Cartan model $(S\mathfrak{g}^* \otimes \Omega(M))^G$ if desired.

As discussed in Section A.4, let $(EG, \Theta_{EG}) \rightarrow BG$ be a universal principal G -bundle with connection, given as a direct limit of smooth finite-dimensional bundles. If M is a G -manifold, the Weil homomorphism induces an inclusion

$$\Omega_G^*(M) \xrightarrow{w(\Theta_{EG}) \otimes 1} \Omega^*(EG \times_G M) \hookrightarrow C^*(EG \times_G M; \mathbb{R})$$

that is a quasi-isomorphism of cochain complexes.

For the moment, we will use general coefficients. Let $\Lambda \subset V$ be a completely disconnected subgroup of a (possibly graded) vector space, and denote $\Omega^*(-; V) = \Omega^*(-) \otimes V$.

Definition 2.1. The differential equivariant cochain complex, denoted by $(\check{C}_G(q)^*(M; \Lambda), d)$, is defined as the homotopy pullback in the diagram

$$(2.2) \quad \begin{array}{ccc} \check{C}_G(q)^*(M; \Lambda) & \xrightarrow{I} & C^*(EG \times_G M; \Lambda) \\ \downarrow R & & \downarrow \\ \Omega_G^{*\geq q}(M; V) & \hookrightarrow & C^*(EG \times_G M; V). \end{array}$$

Explicitly, this is defined so that for $k \geq q$,

$$\begin{aligned} \check{C}_G(q)^k(M; \Lambda) &:= C^k(EG \times_G M; \Lambda) \times C^{k-1}(EG \times_G M; V) \times \Omega_G^k(M; V) \\ d(c, h, \omega) &:= (\delta c, \omega - c - \delta h, d_G \omega), \end{aligned}$$

and for $k < q$ we restrict to the subcomplex with $\omega = 0$.

For $\check{x} = (c, h, \omega)$, we call c the *characteristic cocycle* and ω the *curvature*, and we say \check{x} is *flat* if the curvature is zero. The degree n differential equivariant cohomology is defined as the abelian group

$$\check{H}_G^n(M; \Lambda) := H^n(\check{C}_G(n)^*(M; \Lambda)) = \frac{\check{Z}_G^n(M; \Lambda)}{d\check{C}_G^{n-1}(M; \Lambda)_{\text{flat}}}.$$

As one would expect, the groups $\check{H}_G^*(M; \Lambda)$ lie in short exact sequences that are completely analogous to those for ordinary differential cohomology. We denote the image of $H_G^n(M; \Lambda) \rightarrow H_G^n(M; V)$ by $H_G^n(M; V)_\Lambda$, and we denote the subgroup of closed n -forms with Λ -periods by $\Omega_G^n(M; V)_\Lambda$. This is defined so that under the de Rham isomorphism,

$$\frac{\Omega_G^n(M; V)_\Lambda}{d_G \Omega_G^{n-1}(M; V)} \xrightarrow{\cong} H_G^n(M; V)_\Lambda.$$

We will primarily be concerned with $\Lambda = \mathbb{Z}$, $V = \mathbb{R}$, and we use the notations

$$\check{H}_G^*(M) = \check{H}_G^*(M; \mathbb{Z}), \quad \check{H}_G^* = \check{H}_G^*(\text{pt}; \mathbb{Z}).$$

In later sections, we will simply use $\mathbb{Z} \subset \mathbb{R}$ to simplify notation, but all results will immediately generalize to general coefficients $\Lambda \subset V$.

Proposition 2.3. *The groups $\check{H}_G^*(M; \Lambda)$ lie in the short exact sequences*

$$(SES\ 1) \quad 0 \rightarrow H_G^{n-1}(M; V/\Lambda) \rightarrow \check{H}_G^n(M; \Lambda) \xrightarrow{R} \Omega_G^n(M; V)_\Lambda \rightarrow 0,$$

$$(SES\ 2) \quad 0 \rightarrow \frac{\Omega_G^{n-1}(M; V)}{\Omega_G^{n-1}(M; V)_\Lambda} \rightarrow \check{H}_G^n(M; \Lambda) \xrightarrow{I} H_G^n(M; \Lambda) \rightarrow 0,$$

$$(SES\ 3) \quad 0 \rightarrow \frac{H_G^{n-1}(M; V)}{H_G^{n-1}(M; V)_\Lambda} \rightarrow \check{H}_G^n(M; \Lambda) \rightarrow A_G^n(M; V) \rightarrow 0,$$

where $A_G^n(M)$ is the pullback (in sets) of the commutative square

$$(2.4) \quad \begin{array}{ccc} \check{H}_G^n(M; \Lambda) & \xrightarrow{I} & H_G^n(M; \Lambda) \\ \downarrow R & & \downarrow \\ \Omega_G^n(M)_\Lambda & \xrightarrow{dR} & H_G^n(M; V)_\Lambda. \end{array}$$

These sequences fit into the character diagram

$$\begin{array}{ccccc} & & & & 0 \\ & & & & \nearrow \\ & 0 & \searrow & & \\ & & H_G^{n-1}(M; V/\Lambda) & \xrightarrow{-B} & H_G^n(M; \Lambda) & \searrow & 0 \\ & H_G^{n-1}(M; V) & \nearrow & & \nearrow I & & H_G^n(M; V) \\ & & \Omega_G^{n-1}(M; V) & \xrightarrow{d_G} & \Omega_G^n(M; V)_\Lambda & \nearrow & \\ & 0 & \nearrow & & \searrow & & 0 \\ & & \Omega_G^{n-1}(M; V)_\Lambda & & & & \end{array}$$

where B is the Bockstein homomorphism.

Proof. The proof is straightforward and follows from the exact same arguments as those given in [HS]. Alternatively, one can use the homological

framework of Harvey–Lawson spark complexes, specifically Proposition 1.3 from [HaL]. The key point is that $\Omega_G^*(M) \rightarrow C^*(EG \times_G M; \mathbb{R})$ is an inclusion, as noted in Lemma A.17, and it induces an isomorphism in cohomology. \square

Proposition 2.5.

- (1) $\check{C}_G^*(M; \Lambda)$ and $\check{H}_G^*(M; \Lambda)$ define contravariant functors from $G\text{-Man}$ to cochain complexes and graded abelian groups, respectively.
- (2) An inclusion $\Lambda_1 \hookrightarrow \Lambda_2 \hookrightarrow V$ of totally disconnected subgroups of V induces natural transformations

$$\check{H}_G^*(-; \Lambda_1) \longrightarrow \check{H}_G^*(-; \Lambda_2).$$

- (3) The cup product and wedge product induce algebra homomorphisms

$$\check{H}_G^{k_1}(M; \Lambda_1) \otimes \check{H}_G^{k_2}(M; \Lambda_2) \longrightarrow \check{H}_G^{k_1+k_2}(M; \Lambda_1 \otimes \Lambda_2),$$

where $\Lambda_1 \otimes \Lambda_2 \subset V_1 \otimes V_2$.

- (4) The map $M \xrightarrow{\pi} \text{pt}$ makes $\check{H}_G^*(M; \Lambda)$ into a \check{H}_G^* -module.
- (5) If V is a ring with sub-ring Λ , then $\check{H}_G^*(M; \Lambda)$ is a graded commutative ring, and I and R are ring homomorphisms.

Proof. The constructions $\check{C}_G^*(M)$ and $\check{H}_G^*(M)$ are clearly functorial with respect to equivariant maps $f: M \rightarrow N$, as demonstrated in Appendix A, thus proving (1).

An inclusion $\Lambda_1 \hookrightarrow \Lambda_2$ defines natural maps between cochain complexes $C^*(-; \Lambda_1) \hookrightarrow C^*(-; \Lambda_2)$. This induces $\check{C}_G^*(-; \Lambda_1) \hookrightarrow \check{C}_G^*(-; \Lambda_2)$, thus proving (2).

For (3), first note that the homomorphism $\Omega_G^*(M) \rightarrow \Omega^*(EG \times_G M)$ is a homomorphism of DGAs, as the basic subcomplex of a G^* -algebra is a sub-DGA. Hence, the product structure from [HS] immediately defines the product

$$\begin{aligned} \check{C}_G(q_1)^{k_1}(M; \Lambda_1) \otimes \check{C}_G(q_2)^{k_2}(M; \Lambda_2) &\longrightarrow \check{C}_G(q_1 + q_2)^{k_1+k_2}(M; \Lambda_1 \otimes \Lambda_2) \\ \check{H}_G^{k_1}(M; \Lambda_1) \otimes \check{H}_G^{k_2}(M; \Lambda_2) &\longrightarrow \check{H}_G^{k_1+k_2}(M; \Lambda_1 \otimes \Lambda_2) \end{aligned}$$

by

$$(c_1, h_1, \omega_1) \cdot (c_2, h_2, \omega_2) := (c_1 \cup c_2, (-1)^{|c_1|} c_1 \cup h_2 + h_1 \cup \omega_2 + B(\omega_1, \omega_2), \omega_1 \wedge \omega_2).$$

Here $B(\omega_1, \omega_2) \in C^{k_1+k_2-1}(EG \times_G M; \mathbb{R})$ is any natural cochain homotopy between \wedge and \cup , and we denote the product in $\Omega_G(M)$ by \wedge in a slight abuse of notation.

For (4), the map $M \xrightarrow{\pi} \text{pt}$ is G -equivariant, so it induces natural ring homomorphisms $\check{H}_G^*(\text{pt}; \mathbb{Z}) \xrightarrow{\pi_G^*} \check{H}_G^*(M; \mathbb{Z})$. The composition

$$\check{H}_G^* \otimes \check{H}_G^*(M; \Lambda) \xrightarrow{\pi_G^* \otimes 1} \check{H}_G^*(M; \mathbb{Z}) \otimes \check{H}_G^*(M; \Lambda) \longrightarrow \check{H}_G^*(M; \Lambda)$$

makes $\check{H}_G^*(M; \Lambda)$ into an \check{H}_G^* -module.

Item (5) follows similarly. If Λ is a ring, the product in \check{H}_G^* can be composed with the product in Λ , giving

$$\check{H}_G^*(M; \Lambda) \otimes \check{H}_G^*(M; \Lambda) \xrightarrow{\cdot} \check{H}_G^*(M; \Lambda \otimes \Lambda) \longrightarrow \check{H}_G^*(M; \Lambda).$$

The homomorphisms I and R preserve products at the cochain level, so the induced maps $\check{H}_G^*(M; \Lambda) \xrightarrow{I} H_G^*(M; \Lambda)$ and $\check{H}_G^*(M; \Lambda) \xrightarrow{R} \Omega_G^*(M; V)_\Lambda$ are also ring homomorphisms. \square

Let us now use Propositions 2.3 and 2.5 to analyze $\check{H}_G^*(M)$ in a few simple examples.

Example 2.6. If $G = e$ is the trivial group, then $\Omega_G(M) \cong \Omega(M)$, and we can take $EG = \text{pt}$. This recovers the usual differential cohomology

$$\check{H}_e^*(M; \Lambda) \cong \check{H}^*(M; \Lambda).$$

Example 2.7. The point pt is trivially a G -manifold, and the Borel construction is $EG \times_G \text{pt} \cong BG$. For G compact we have $H^{2k}(BG; \mathbb{R}) \cong \Omega_G^{2k}(\text{pt}) = S^k(\mathfrak{g}^*)^G$ and $H^{2k+1}(BG; \mathbb{R}) = \Omega_G^{2k+1}(\text{pt}) = 0$. The short exact sequences (SES 2) and (SES 1) give isomorphisms

$$(2.8) \quad \check{H}_G^{2k} \cong H^{2k}(BG; \mathbb{Z}), \quad \check{H}_G^{2k+1} \cong H^{2k}(BG; \mathbb{R}/\mathbb{Z}).$$

Furthermore, (SES 2) becomes

$$(2.9) \quad 0 \longrightarrow \frac{S^k(\mathfrak{g}^*)^G}{S^k(\mathfrak{g}^*)_{\mathbb{Z}}^G} \longrightarrow \check{H}_G^{2k+1} \longrightarrow H^{2k+1}(BG; \mathbb{Z}) \longrightarrow 0,$$

where $\frac{S^k(\mathfrak{g}^*)^G}{S^k(\mathfrak{g}^*)_{\mathbb{Z}}^G}$ can also be written as $H^{2k}(BG; \mathbb{R}) \otimes \mathbb{R}/\mathbb{Z}$.

Example 2.10. If $G = \Gamma$ is a finite group, the Lie algebra is trivial. Hence, the equivariant forms $\Omega_\Gamma(M)$ are simply the invariant forms $\Omega(M)^\Gamma$. When M is a point, (SES 2) implies

$$\check{H}_\Gamma^* \cong H^*(B\Gamma; \mathbb{Z}).$$

Example 2.11. For $G = S^1$, we know that $H^*(BS^1; \mathbb{Z}) \cong H^*(BS^1; \mathbb{R})_{\mathbb{Z}} \cong \mathbb{Z}[t]$, where $|t| = 2$. The above identities (2.8) and (2.9) imply

$$\begin{aligned} \check{H}_{S^1}^{2k} &\cong H^{2k}(BS^1; \mathbb{Z}) \cong \mathbb{Z}t^k, \\ \check{H}_{S^1}^{2k+1} &\cong H^{2k}(BS^1; \mathbb{R}) \otimes \mathbb{R}/\mathbb{Z} \cong (\mathbb{R}/\mathbb{Z})t^k. \end{aligned}$$

We now show that as a graded commutative ring, this can be written

$$\check{H}_{S^1}^* \cong \left(\mathbb{Z} \oplus (\mathbb{R}/\mathbb{Z})\theta \right)[t],$$

where $|\theta| = 1$ and $|t| = 2$. The relations $\theta t = t\theta$ and $\theta^2 = 0$ are implied by the adjectives graded commutative.

To verify this is a ring isomorphism, first note that the characteristic class map gives a ring isomorphism $\check{H}_{S^1}^{2*} \cong \mathbb{Z}[t]$ in even degrees. We can use the product's definition to determine what happens when multiplying

by odd-degree elements. Choose some cocycle $\alpha \in Z^2(BS^1; \mathbb{Z})$ representing the generator of $H^2(BS^1; \mathbb{Z})$. An even element $nt^k \in \mathbb{Z}t^k \cong \check{H}_{S^1}^{2k}$ can be represented by a cocycle of the form (α^k, b, t^k) , and an odd element $[r]\theta t^k \in (\mathbb{R}/\mathbb{Z})\theta t^k \cong \check{H}_{S^1}^{2k+1}$ can be represented by $(0, r\alpha^k, 0)$. The cochain-level product yields

$$\begin{aligned} (n\alpha^{k_1}, b, nt^{k_1}) \cdot (0, r\alpha^{k_2}, 0) &= (0, nr\alpha^{k_1+k_2}, 0), \\ (0, r_1\alpha^{k_1}, 0) \cdot (0, r_2\alpha^{k_2}, 0) &= (0, 0, 0). \end{aligned}$$

So, the product of two odd elements is 0, and $(nt^{k_1}) \cdot ([r]\theta t^{k_2}) = [nr]\theta t^{k_1+k_2}$.

Example 2.12. Similarly, the odd cohomology groups of $BU(n)$ vanish, and

$$H^*(BU(n); \mathbb{Z}) \cong H^*(BU(n); \mathbb{R})_{\mathbb{Z}} \cong \left(S^{*/2} \mathfrak{u}(n)^* \right)_{\mathbb{Z}}^{U(n)} \cong \mathbb{Z}[c_1, \dots, c_n].$$

The same argument given in the previous example shows that

$$\begin{aligned} \check{H}_{U(n)}^{2k} &\cong H^{2k}(BU(n); \mathbb{Z}), \\ \check{H}_{U(n)}^{2k+1} &\cong H^{2k}(BU(n); \mathbb{R}) \otimes \mathbb{R}/\mathbb{Z}, \end{aligned}$$

and as graded commutative rings with $|c_i| = 2i$ and $|\theta| = 1$,

$$\check{H}_{U(n)}^* \cong \left(\mathbb{Z} \oplus (\mathbb{R}/\mathbb{Z})\theta \right)[c_1, \dots, c_n].$$

3. Differential cohomology of the differential quotient stack

From the description of the Weil algebra $W(\mathfrak{g})$ in Appendix A, it is clear that equivariant forms $\Omega_G(M)$ most naturally arise when considering principal bundles *with connection*. While the topological space $EG \rightarrow BG$ can be viewed as a universal bundle with connection, it does not naturally represent the *category* of principal bundles with connection on general manifolds. For this reason, it will be more convenient to replace $EG \times_G M$ with the stack $\mathcal{E}_{\nabla}G \times_G M$ and study its differential cohomology. While we are not aware of any work specifically on $\check{H}^*(\mathcal{E}_{\nabla}G \times_G M)$, the general idea of defining differential cohomology via sheaves on manifolds has been widely used. This includes, but is not limited to, [Bun, BunNV, HQ, FSS].

3.1. Background. First, remember that a *groupoid* is a category in which every morphism is invertible. Any *set* S can be viewed as a category, where the only morphisms are the identity morphisms. We denote this fully faithful embedding of the category of sets into the category of groupoids by $\text{Set} \subset \text{Gpd}$.

A *stack* \mathcal{M} (on the site of smooth manifolds) is a contravariant functor from the category of manifolds to the category of groupoids satisfying a descent condition; we denote the category by

$$\text{Stack} = \text{Shv}_{\text{Gpd}} \subset \text{Fun}(\text{Man}^{\text{op}}, \text{Gpd}).$$

This means that associated to every manifold X is a groupoid $\mathcal{M}(X)$, and a smooth map $X_1 \xrightarrow{f} X_2$ induces a functor

$$\mathcal{M}(X_2) \xrightarrow{f^*} \mathcal{M}(X_1).$$

There are associative natural transformations $g^*f^* \cong (g \circ f)^*$ to deal with composition of functions. To be a stack, as opposed to a prestack, \mathcal{M} must also satisfy a sheaf/descent condition that all of our examples will satisfy (see [Hei] or Section 6 for more details).

Example 3.1. Any smooth manifold N defines a stack \underline{N} by associating to X the set

$$\underline{N}(X) = C^\infty(X, N) \in \mathbf{Set},$$

and to a smooth map $X_1 \xrightarrow{F} X_2$ the pullback $C^\infty(X_2, N) \xrightarrow{F^*} C^\infty(X_1, N)$.

Example 3.2. Principal G -bundles with connection form a stack, which we denote $\mathcal{B}_\nabla G$. To any manifold X , let $\mathcal{B}_\nabla G(X)$ be the groupoid whose objects are principal G -bundles $P \xrightarrow{\pi} X$ with connection $\Theta \in \Omega^1(P; \mathfrak{g})$. A *morphism* $(P_1, \Theta_1) \xrightarrow{\varphi} (P_2, \Theta_2)$ is a bundle map preserving the connection; i.e., it is a G -equivariant map φ

$$\begin{array}{ccc} P_1 & \xrightarrow{\varphi} & P_2 \\ & \searrow \pi_1 & \swarrow \pi_2 \\ & X & \end{array}$$

such that $\varphi^*(\Theta_2) = \Theta_1$. Such a φ must be a diffeomorphism, and hence all morphisms in $\mathcal{B}_\nabla G(X)$ are isomorphisms. Since bundles and connections pull back, a smooth map $f: X_1 \rightarrow X_2$ induces a functor

$$f^*: \mathcal{B}_\nabla G(X_2) \rightarrow \mathcal{B}_\nabla G(X_1).$$

The groupoid of principal G -bundles on X , without connection, is defined analogously and denoted $\mathcal{B}G$.

The collection of *morphisms between stacks* $\mathbf{Shv}_{\mathbf{Gpd}}(\mathcal{M}_1, \mathcal{M}_2)$ naturally forms a groupoid. An object $\mathcal{M}_1 \xrightarrow{\Psi} \mathcal{M}_2$ in this groupoid is a collection of functors $\mathcal{M}_1(X) \xrightarrow{\Psi(X)} \mathcal{M}_2(X)$ for all X , together with natural transformations $\Psi(f): \Psi(X) \circ f^* \rightarrow f^* \circ \Psi(Y)$ for all smooth $f: X \rightarrow Y$; morphisms between morphisms are given by natural transformations, which must be invertible since $\mathcal{M}_2(-)$ is always a groupoid.

The Yoneda Lemma states that there is a canonical equivalence of categories

$$\mathcal{M}(X) \cong \mathbf{Shv}_{\mathbf{Gpd}}(\underline{X}, \mathcal{M}),$$

and this defines a faithful embedding of the category of manifolds into the 2-category of stacks. For this reason, we will usually not distinguish between a manifold X and its associated stack \underline{X} . We can also view an object in

the category $\mathcal{M}(X)$ as a map $X \rightarrow \mathcal{M}$. Thus, maps between stacks are a generalization of smooth maps between manifolds; when M and X are both manifolds,

$$\text{Shv}_{\text{Gpd}}(\underline{X}, \underline{M}) \cong \underline{M}(X) = C^\infty(X, M).$$

Via Yoneda, any bundle $(P, \Theta) \rightarrow X$ is naturally equivalent to a map

$$X \xrightarrow{(P, \Theta)} \mathcal{B}_\nabla G,$$

and the groupoid of bundles with connection is naturally equivalent to the category of maps $X \rightarrow \mathcal{B}_\nabla G$. This makes $\mathcal{B}_\nabla G$ a more convenient classifying object for many purposes than the ordinary topological space BG .

Example 3.3. Differential forms of degree k define a stack Ω^k , where

$$\Omega^k(X) \in \text{Set}$$

is a set viewed as a groupoid with only identity morphisms. A differential form $\omega \in \Omega^k(X)$ is equivalent to a morphism $X \xrightarrow{\omega} \Omega^k$.

Example 3.4. If M is a (left) G -manifold, the differential quotient stack $\mathcal{E}_\nabla G \times_G M$ is defined as follows. An object of $(\mathcal{E}_\nabla G \times_G M)(X)$ is a principal G -bundle with connection $(P, \Theta) \rightarrow X$, together with a G -equivariant map $f: P \rightarrow M$. The map f is equivalent to a section F of the associated fiber bundle $P \times_G M$, with the equivalence given by $F(x) = [p, f(p)] \in P \times_G M$.

$$X \xrightarrow{(P, \Theta, f)} \mathcal{E}_\nabla G \times_G M \quad \longleftrightarrow \quad \begin{array}{ccc} (P, \Theta) & \xrightarrow{f} & M \\ \downarrow & & \\ X & & \end{array} \quad \longleftrightarrow \quad \begin{array}{ccc} (P, \Theta) \times_G M & & \\ \downarrow \Big) F & & \\ X & & \end{array}$$

A morphism

$$\left(X \leftarrow (P_1, \Theta_1) \xrightarrow{f_1} M \right) \xrightarrow{\varphi} \left(X \leftarrow (P_2, \Theta_2) \xrightarrow{f_2} M \right)$$

is a connection-preserving bundle isomorphism that covers the maps to M ; i.e.,

$$\begin{array}{ccccc} & & (P_1, \Theta_1) & & \\ & \swarrow & \downarrow \varphi \cong & \searrow f_1 & \\ X & & & & M \\ & \swarrow & \downarrow \varphi \cong & \searrow f_2 & \\ & & (P_2, \Theta_2) & & \end{array}$$

with $\varphi^*(\Theta_2) = \Theta_1$.

In the case where $M = \text{pt}$, there is only one map $P \xrightarrow{f} \text{pt}$, leading to the natural isomorphism $\mathcal{E}_\nabla G \times_G \text{pt} \cong \mathcal{B}_\nabla G$. We also let $\mathcal{E}G \times_G M$ denote the stack defined analogously via bundles without connection.

Remark 3.5. Other possible notations for this quotient stack with connections could include $M //^\nabla G$ and $(M_G)_\nabla$. We choose to use $\mathcal{E}_\nabla G \times_G M$ to emphasize the correspondence between maps of stacks and maps of topological spaces.

3.2. Differential forms on the quotient stack. We want to study the differential cohomology of $\mathcal{E}_\nabla G \times_G M$, but it is not immediately clear what that means. As a first step, let's understand differential forms. Since $\Omega^n(X) \cong \text{Shv}_{\text{Gpd}}(X, \Omega^n)$, it is natural to define

$$(3.6) \quad \Omega^n(\mathcal{E}_\nabla G \times_G M) := \text{Shv}_{\text{Gpd}}(\mathcal{E}_\nabla G \times_G M, \Omega^n).$$

An element of $\Omega^n(\mathcal{E}_\nabla G \times_G M)$ is equivalent to functorially assigning, for all manifolds X , an element of $\Omega^n(X)$ to every object in $(\mathcal{E}_\nabla G \times_G M)(X)$. Since $\Omega^n(X)$ has only identity morphisms, isomorphic objects in $(\mathcal{E}_\nabla G \times_G M)(X)$ must give equal elements in $\Omega^n(X)$.

Based on the description of $\Omega_G(M)$ in Appendix A, there is an obvious homomorphism

$$(3.7) \quad \Omega_G^n(M) \xrightarrow{\cong} \Omega^n(\mathcal{E}_\nabla G \times_G M).$$

While it is not difficult to show this is injective, Freed–Hopkins prove it is in fact an isomorphism of abelian groups [FreH, Theorem 7.28]. Though $\Omega_G(M)$ is usually regarded as an algebraic replacement for $\Omega(EG \times_G M)$, it is the actual de Rham complex of the stack $\mathcal{E}_\nabla G \times_G M$.

We now explain the homomorphism (3.7). To a map $X \rightarrow \mathcal{E}_\nabla G \times_G M$, which is an object of $(\mathcal{E}_\nabla G \times_G M)(X)$, the composition $\Theta^* \circ f_G^* = \Theta^* \otimes f^*$ defines a homomorphism $\Omega_G^n(M) \rightarrow \Omega^n(X)$, as indicated in the following diagram.

$$\begin{array}{ccc} (P, \Theta) \xrightarrow{f} M & & \Omega_G^n(P) \xleftarrow{f_G^*} \Omega_G^n(M) \\ \downarrow & \mapsto & \downarrow \Theta^* \\ X & & \Omega^n(P)_{\text{basic}} \cong \Omega^n(X) \end{array}$$

More explicitly, for $\omega \otimes \eta \in \mathbb{W}^{2i,j}(\mathfrak{g})$ and $\psi \in \Omega^k(M)$, then

$$\omega \otimes \eta \otimes \psi \mapsto \omega(\Omega^{\wedge i}) \wedge \eta(\Theta^{\wedge j}) \wedge f^* \psi \in \pi^* \Omega^{2i+j+k}(X).$$

This construction is invariant under morphisms in $(\mathcal{E}_\nabla G \times_G M)(X)$. If

$$\begin{array}{ccccc} & & (P_1, \Theta_{P_1}) & & \\ & \swarrow \pi_1 & \downarrow \phi \cong & \searrow f_1 & \\ X & & & & M \\ & \swarrow \pi_2 & \downarrow \phi \cong & \searrow f_2 & \\ & & (P_2, \Theta_{P_2}) & & \end{array}$$

with $\phi^*(\Theta_2) = \Theta_1$, it immediately follows that

$$\phi^*(\Theta_2^* \otimes f_2^*) = \phi^* \Theta_2^* \otimes (f_2 \circ \phi)^* = \Theta_1^* \otimes f_1^*.$$

Since ϕ covers the identity on X , the two homomorphisms $\Omega_G^n(M) \rightarrow \Omega^n(X)$ are equal.

The de Rham differential induces a universal map of stacks $d: \Omega^n \rightarrow \Omega^{n+1}$. This gives a differential $\Omega^n(\mathcal{E}_{\nabla}G \times_G M) \xrightarrow{d} \Omega^{n+1}(\mathcal{E}_{\nabla}G \times_G M)$, and $(\Omega^*(\mathcal{E}_{\nabla}G \times_G M), d)$ is naturally isomorphic to $(\Omega_G^*(M), d_G)$ as a cochain complex.

Remark 3.8. Note that the cochain complex $\Omega(\mathcal{E}_{\nabla}G \times_G M)$ is given by

$$\cdots \xrightarrow{d} \text{Shv}_{\text{Gpd}}(\mathcal{E}_{\nabla}G \times_G M, \Omega^n) \xrightarrow{d} \text{Shv}_{\text{Gpd}}(\mathcal{E}_{\nabla}G \times_G M, \Omega^{n+1}) \xrightarrow{d} \cdots$$

Each degree is computed individually, with Ω^n being set valued, as opposed to some version of maps to the complex Ω^* , where there are nontrivial morphisms.

Example 3.9. In the case where $M = \text{pt}$, we have that $(S\mathfrak{g}^*)^G \cong \Omega(\mathcal{B}_{\nabla}G)$, which is concentrated in even degrees. A bundle with connection $(P, \Theta) \rightarrow X$ is naturally viewed as a map $(P, \Theta): X \rightarrow \mathcal{B}_{\nabla}G$; the induced map on differential forms is the usual map from Chern–Weil theory

$$\begin{array}{ccc} (S^n \mathfrak{g}^*)^G & \xrightarrow{\cong} & \Omega^{2n}(\mathcal{B}_{\nabla}G) \xrightarrow{(P, \Theta)^*} \Omega^{2n}(X) \\ \omega \longmapsto & & \omega(\Omega^{\wedge n}). \end{array}$$

3.3. Differential cohomology via sheaves. We now briefly outline the construction of $\check{H}^n(\mathcal{E}_{\nabla}G \times_G M)$ and state the main properties. Because the details are not important to understanding the constructions in Sections 4 and 5, we postpone them until Section 6.

Amongst its many constructions, ordinary cohomology can be represented by homotopy classes of maps to a space. For A an abelian group, there exists an Eilenberg–MacLane space $K(A, n) \in \text{Top}$ with

$$H^n(X; A) \cong [X, K(A, n)] = \text{hoTop}(X, K(A, n)).$$

While differential cohomology cannot be represented by a fixed topological space, it can be represented by a sheaf of “spaces” on the site of manifolds [FSS, Bun, HQ, BunNV]. We work with simplicial sets for convenience, as they form a natural home for groupoids, topological spaces, and chain complexes. Letting ∞Gpd denote the $(\infty, 1)$ -category of simplicial sets spanned by Kan complexes, define $\check{\mathcal{K}}(\mathbb{Z}, n) \in \text{Shv}_{\infty\text{Gpd}}$ to be a sheaf fitting into a homotopy-commutative diagram of the form

$$(3.10) \quad \begin{array}{ccc} \check{\mathcal{K}}(\mathbb{Z}, n) & \longrightarrow & \mathcal{K}(\mathbb{Z}, n) \\ \downarrow & & \downarrow \\ \mathbb{N}(\Omega_{\text{cl}}^n) & \longrightarrow & \mathcal{K}(\mathbb{R}, n). \end{array}$$

Here, $\mathcal{K}(A, n)$ is an object in $\text{Shv}_{\infty\text{Gpd}}$ representing ordinary cohomology.

We can consider the sheaf of groupoids $\mathcal{E}_{\nabla}G \times_G M$ as a sheaf of ∞ -groupoids via the nerve construction N . The cohomology groups and differential cohomology groups are defined by considering maps between sheaves in the homotopy category:

$$\begin{aligned} H^n(\mathcal{E}_{\nabla}G \times_G M; A) &:= \text{hoShv}_{\infty\text{Gpd}}(N(\mathcal{E}_{\nabla}G \times_G M), \mathcal{K}(A, n)), \\ \check{H}^n(\mathcal{E}_{\nabla}G \times_G M) &:= \text{hoShv}_{\infty\text{Gpd}}(N(\mathcal{E}_{\nabla}G \times_G M), \check{\mathcal{K}}(\mathbb{Z}, n)). \end{aligned}$$

The following theorem is essentially a consequence of the Freed–Hopkins isomorphism (3.7) and the work of Bunke–Nikolaus–Völkl on homotopy-invariant sheaves [BunNV]. The proof is given at the end of Section 6.

Theorem 3.11. *The two definitions of $\check{H}_G^*(M)$ are naturally equivalent,*

$$\check{H}^n(\mathcal{E}_{\nabla}G \times_G M) \cong H^n(\check{C}_G(n)^*(M; \mathbb{Z})).$$

This leads to the following commutative square.

(3.12)

$$\begin{array}{ccc} \check{H}^n(\mathcal{E}_{\nabla}G \times_G M) & \longrightarrow & H^n(\mathcal{E}_{\nabla}G \times_G M; \mathbb{Z}) \cong H_G^n(M; \mathbb{Z}) \\ \downarrow & & \downarrow \\ \Omega_G^n(M)_{\text{cl}} \cong \Omega_{\text{cl}}^n(\mathcal{E}_{\nabla}G \times_G M) & \longrightarrow & H^n(\mathcal{E}_{\nabla}G \times_G M; \mathbb{R}) \cong H_G^n(M; \mathbb{R}). \end{array}$$

The isomorphism in the bottom-left corner is given by the Freed–Hopkins isomorphism (3.7). The two isomorphisms on the right are induced by the map $EG \times_G M \rightarrow \mathcal{E}_{\nabla}G \times_G M$, which induces an isomorphism in cohomology (Proposition 6.7). In the following sections, we will use these two properties to check that maps of stacks induce the desired maps at the level of differential forms and cohomology.

Example 3.13. Since $\Omega^{2k-1}(\mathcal{B}_{\nabla}G) = 0$, the short exact sequence (SES 2) gives an isomorphism

$$\check{H}^{2k}(\mathcal{B}_{\nabla}G) \xrightarrow{\cong} H^{2k}(BG; \mathbb{Z}).$$

The Cheeger–Chern–Simons refinement of Chern–Weil theory [CheeS] can then be repackaged in the following way. A principal G -bundle with connection $(P, \Theta) \rightarrow X$ is equivalent to a map

$$X \xrightarrow{(P, \Theta)} \mathcal{B}_{\nabla}G,$$

and this induces a homomorphism

$$\check{H}^{2k}(X) \xleftarrow{(P, \Theta)^*} \check{H}^{2k}(\mathcal{B}_{\nabla}G) \cong H^{2k}(BG; \mathbb{Z}).$$

Hence, any universal characteristic class in $H^{2k}(BG; \mathbb{Z})$ has a canonical differential refinement.

4. Constructions of maps between stacks

We now explain how some important constructions, frequently described using classifying spaces or equivariant forms, are naturally given by explicit geometric constructions involving bundles with connection.

4.1. Associated bundles. Let $\phi: G_1 \rightarrow G_2$ be any Lie group homomorphism, with $\phi_*: \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ the associated Lie algebra homomorphism. The associated bundle construction, which functorially makes any G_1 -bundle with connection (P, Θ) into a G_2 -bundle with connection, induces morphisms

$$\begin{array}{ccc} \mathcal{E}_\nabla G_1 & \longrightarrow & \mathcal{E}_\nabla G_2 \\ \downarrow & & \downarrow \\ \mathcal{B}_\nabla G_1 & \longrightarrow & \mathcal{B}_\nabla G_2. \end{array}$$

The most common example is when ϕ is an inclusion, which is usually referred to as extending the structure group. For more details on the associated bundle construction, see Chapter II.6 of [KN] or Section 1 of [Fre].

If $(P, \Theta) \in \mathcal{B}_\nabla G_1(X)$, the associated G_2 -bundle is defined

$$(4.1) \quad P_\phi := P \times_\phi G_2 = (P \times G_2)/G_1 = (P \times G_2) / ((pg, g') \sim (p, \phi(g)g')).$$

To define the induced connection Θ_ϕ on P_ϕ , first note that the natural map $P \xrightarrow{\varphi} P_\phi$, given by $p \mapsto [(p, 1)]$, is ϕ -equivariant. Hence, the horizontal subspaces of TP are mapped equivariantly into TP_ϕ , and the image extends uniquely to an equivariant horizontal distribution in TP_ϕ . The induced connection Θ_ϕ can also be described as the unique connection on P_ϕ that is compatible with Θ in the sense that

$$(4.2) \quad \varphi^*(\Theta_\phi) = \phi_*(\Theta) \in \Omega^1(P; \mathfrak{g}_2).$$

Therefore, the associated bundle construction is compatible with the Weil homomorphism, giving the commutative diagram

$$(4.3) \quad \begin{array}{ccc} W(\mathfrak{g}_1) & \xrightarrow{\Theta^*} & \Omega(P) \\ \uparrow \phi^* & & \uparrow \varphi^* \\ W(\mathfrak{g}_2) & \xrightarrow{\Theta_\phi^*} & \Omega(P_\phi). \end{array}$$

Proposition 4.4. *Let $F: M \rightarrow N$ be a ϕ -equivariant map, where*

$$\phi: G_1 \rightarrow G_2$$

is a Lie group homomorphism.

- (1) *The associated bundle construction gives a natural morphism*

$$\mathcal{E}_\nabla G_1 \times_{G_1} M \xrightarrow{F_\phi} \mathcal{E}_\nabla G_2 \times_{G_2} N.$$

- (2) *The homomorphism induced by F_ϕ on differential forms is naturally isomorphic to $\phi^* \otimes F^*$ in the Weil model.*

$$\begin{array}{ccc} \Omega_{G_2}(N) \xrightarrow{\cong} \Omega(\mathcal{E}_\nabla G_2 \times_{G_2} N) \xrightarrow{F_\phi^*} \Omega(\mathcal{E}_\nabla G_1 \times_{G_1} M) \xleftarrow{\cong} \Omega_{G_1}(M) \\ \square \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \square \\ \mathbb{W}(\mathfrak{g}_2) \otimes \Omega(N) \xrightarrow{\phi^* \otimes F^*} \mathbb{W}(\mathfrak{g}_1) \otimes \Omega(M) \end{array}$$

- (3) *The homomorphism induced by F_ϕ on ordinary cohomology is naturally isomorphic to the homomorphism induced by the maps of classifying spaces*

$$EG_1 \times_{G_1} M \rightarrow (EG_1 \times_\phi G_2) \times_{G_2} N \rightarrow EG_2 \times_{G_2} N.$$

- (4) *The induced homomorphism*

$$\check{H}_{G_2}^*(N) \xrightarrow{F_\phi^*} \check{H}_{G_1}^*(M)$$

recovers the expected homomorphisms in equivariant cohomology and equivariant forms.

Before giving the proof of Proposition 4.4, we wish to first introduce notation for three special cases. In two of the cases, the general construction simplifies, so we also show what happens when the stack is evaluated on a general manifold X .

Example 4.5. Suppose that M and N are G -manifolds, and $M \xrightarrow{F} N$ is G -equivariant. The morphism of stacks

$$\begin{array}{ccc} \mathcal{E}_\nabla G \times_G M & \xrightarrow{F_G} & \mathcal{E}_\nabla G \times_G N \\ (P, \Theta) \xrightarrow{f} M & \mapsto & (P, \Theta) \xrightarrow{f} M \xrightarrow{F} N \\ \downarrow & & \downarrow \\ X & & X \end{array}$$

induces the homomorphism of abelian groups

$$\check{H}_G^*(N) \xrightarrow{F_G^*} \check{H}_G^*(M).$$

Example 4.6. Let M be a G_2 -manifold. A homomorphism $G_1 \xrightarrow{\phi} G_2$ naturally defines a G_1 -action on M via $g_1 \cdot M := \phi(g_1) \cdot M$, and the identity map $M \xrightarrow{\mathbb{1}} M$ is ϕ -equivariant. The associated bundle construction then induces

$$\begin{array}{c} \mathcal{E}_\nabla G_1 \times_{G_1} M \xrightarrow{\mathbb{1}_\phi} \mathcal{E}_\nabla G_2 \times_{G_2} M, \\ \check{H}_{G_1}^*(M) \xleftarrow{\mathbb{1}_\phi^*} \check{H}_{G_2}^*(M). \end{array}$$

Example 4.7. Suppose that $K \triangleleft G$ is a normal subgroup, and suppose the quotient map $M \xrightarrow{q} M/K$ is a smooth map between manifolds. Then, q is equivariant with respect to the quotient group homomorphism $G \xrightarrow{/K} G/K$. Quotienting M and G by K induces the stack morphism

$$\begin{array}{ccc}
 \mathcal{E}_{\nabla}G \times_G M & \xrightarrow{q/K} & \mathcal{E}_{\nabla}(G/K) \times_{G/K} M/K \\
 (P, \Theta) \xrightarrow{f} M & \longmapsto & \begin{array}{ccc} (P, \Theta) & \xrightarrow{f} & M \\ \vdots \downarrow & & \vdots \downarrow q \\ (P/K, \Theta/K) & \xrightarrow{f/K} & M/K \\ \downarrow & & \downarrow \\ X & & X \end{array}
 \end{array}$$

(with dotted arrows indicating maps no longer being used) and a homomorphism

$$\check{H}_{G/K}^*(M/K) \xrightarrow{q^*/K} \check{H}_G^*(M).$$

If K acts freely, this induces the usual isomorphism

$$(4.8) \quad H_{G/K}^*(M/K; -) \xrightarrow[\cong]{q^*/K} H_G^*(M; -)$$

in equivariant cohomology.

Proof of Proposition 4.4. Part (1) is given by constructing, for every X and natural with respect to maps $X \rightarrow Y$, the following functor.

$$\begin{array}{ccc}
 (\mathcal{E}_{\nabla}G_1 \times_{G_1} M)(X) & \xrightarrow{F_\phi(X)} & (\mathcal{E}_{\nabla}G_2 \times_{G_2} N)(X) \\
 (P, \Theta) \times_{G_1} M & \longmapsto & (P_\phi, \Theta_\phi) \times_{G_2} N \\
 \downarrow \Big) s & & \downarrow \Big) (\overline{\varphi \times F}) \circ s \\
 X & & X
 \end{array}$$

To explain in a bit more detail, the associated bundle construction associates to (P, Θ) the bundle (P_ϕ, Θ_ϕ) , along with a ϕ -equivariant map $P \xrightarrow{\varphi} P_\phi$. Since $M \xrightarrow{F} N$ is also ϕ -equivariant, the map

$$P \times M \xrightarrow{\varphi \times F} P_\phi \times N$$

descends to the quotient

$$P \times_{G_1} M \xrightarrow{\overline{\varphi \times F}} P_\phi \times_{G_2} N,$$

and we precompose with original section s . The associated bundle construction is functorial, so any morphism Φ in $(\mathcal{E}_{\nabla}G_1 \times_{G_1} M)(X)$ will induce a

morphism $F_\phi(\Phi)$ in $(\mathcal{E}_\nabla G_1 \times_{G_1} M)(X)$, as evidenced by the following commutative diagram.

$$\begin{array}{ccccc}
 & & (P, \Theta) \times_{G_1} M & \xrightarrow{\overline{\varphi \times F}} & (P_\phi, \Theta_\phi) \times_{G_2} N \\
 & \nearrow s & \downarrow \cong \Phi & & \downarrow \cong \Phi_\phi \\
 X & & & & \\
 & \searrow s' & (P', \Theta') \times_{G_1} M & \xrightarrow{\overline{\varphi' \times F}} & (P'_\phi, \Theta'_\phi) \times_{G_2} N
 \end{array}$$

Hence, the associated bundle construction induces a morphism of stacks.

For part (2), we simply trace through the construction of F_ϕ and the Weil homomorphism. Let $\omega \in \Omega(\mathcal{E}_\nabla G_2 \times_{G_2} N)$, which may be written as a sum of homogeneous elements $\alpha \otimes \psi \in (\mathbb{W}(\mathfrak{g}_2) \otimes \Omega(N))_{\text{basic}}$. Then, calculate $F_\phi^* \omega \in \Omega(\mathcal{E}_\nabla G_1 \times_{G_1} M)$ by evaluating on a test manifold X . By definition,

$$\begin{array}{ccc}
 (\mathcal{E}_\nabla G_1 \times_{G_1} M)(X) & \xrightarrow{F_\phi} & (\mathcal{E}_\nabla G_2 \times_{G_2} N)(X) \xrightarrow{\omega} \Omega(X) \\
 (P, \Theta) \times_{G_1} M & & (P_\phi, \Theta_\phi) \times_{G_2} N \\
 \downarrow \Big) s & \mapsto & \downarrow \Big)_{(\varphi \times F) \circ s} \mapsto s^* \left((\overline{\varphi \times F})^* (\alpha(\Theta_\phi) \otimes \psi) \right) \\
 X & & X
 \end{array}$$

As noted in (4.3), the associated bundle construction is compatible with the Weil homomorphism, and therefore

$$(\varphi \times F)^* (\alpha(\Theta_\phi) \otimes \psi) = \alpha(\varphi^* \Theta_\phi) \otimes F^* \psi = \alpha(\phi_* \Theta) \otimes F^* \psi = (\phi^* \alpha)(\Theta) \otimes F^* \psi.$$

Consequently, $F_\phi^* \omega$ gets mapped to $s^* ((\phi^* \alpha)(\Theta) \otimes F^* \psi) \in \Omega(X)$, and this is the same element that $\phi^* \alpha \otimes F^* \psi \in \Omega_{G_1}(M)$ maps to.

To see part (3), choose a classifying map for the G_2 -bundle $EG_1 \times_\phi G_2$, which gives the following diagram. We do not need to assume the classifying map is connection preserving.

$$\begin{array}{ccccc}
 & & \Psi & & \\
 EG_1 \times_{G_1} M & \longrightarrow & EG_1 \times_\phi G_2 & \longrightarrow & EG_2 \\
 \downarrow & & \downarrow & & \downarrow \\
 BG_1 & \xrightarrow{=} & BG_1 & \longrightarrow & BG_2
 \end{array}$$

For any $X \xrightarrow{f} BG_1$, there is an isomorphism of associated bundles

$$f^*(\Psi^* EG_2) \cong (f^* EG_1)_\phi.$$

This implies the diagram

$$\begin{array}{ccc}
 EG_1 \times_{G_1} M & \longrightarrow & EG_2 \times_{G_2} M \\
 \downarrow & & \downarrow \\
 \mathcal{E}G_1 \times_{G_1} M & \longrightarrow & \mathcal{E}G_2 \times_{G_2} N
 \end{array}$$

is commutative up to isomorphism, or homotopy-commutative. As shown in Proposition 6.7, the vertical maps induce isomorphisms in ordinary cohomology. Therefore, the induced map on the cohomology of the differential quotient stacks

$$H^*(\mathcal{E}_{\nabla}G_2 \times_{G_2} N; A) \xrightarrow{F_\phi^*} H^*(\mathcal{E}_{\nabla}G_1 \times_{G_1} M; A)$$

is given by the usual map between the homotopy quotient spaces.

Part (4) follows immediately from parts (2) and (3), and (3.12). □

4.2. Pulling back equivariant bundles. As discussed in the previous section, if G acts freely on a manifold P , there is a natural homomorphism

$$\check{H}^*(P/G) \xrightarrow{q_{/G}^*} \check{H}_G^*(P)$$

inducing the standard isomorphism in cohomology. We now describe the one-sided inverse to this map. First we outline the result, then we give the construction, and then we check the details.

To obtain a map in the opposite direction, one must choose a connection Θ on P . Using this and denoting $M = P/G$, one naturally obtains a morphism of stacks

$$M \xrightarrow{\Theta} \mathcal{E}_{\nabla}G \times_G P$$

by pulling back the bundle (P, Θ) , as indicated below when the stack is evaluated on a manifold X .

$$\begin{array}{ccc}
 X \xrightarrow{f} M & \mapsto & f^*(P, \Theta) \xrightarrow{\tilde{f}} (P, \Theta) \\
 & & \downarrow \qquad \qquad \qquad \downarrow \text{dotted} \\
 & & X \xrightarrow{\text{dotted } f} M
 \end{array}$$

Proposition 4.16 will imply that the induced map

$$\check{H}_G^*(P) = \check{H}^*(\mathcal{E}_{\nabla}G \times_G P) \xrightarrow{\Theta^*} \check{H}^*(M)$$

recovers the isomorphism inverse to $q_{/G}^*$ at the level of cohomology, and it recovers the Weil homomorphism

$$\begin{array}{ccc}
 \Omega_G(P) = (W(\mathfrak{g}) \otimes \Omega(P))_{\text{basic}} & \xrightarrow{\Theta^*} & \Omega(P)_{\text{basic}} \cong \Omega(M) \\
 \alpha \otimes \psi & \mapsto & \alpha(\Theta) \otimes \psi
 \end{array}$$

at the level of differential forms.

The above is actually a special case of what happens when we have an equivariant principal bundle with connection. Suppose that $(Q, \Theta) \rightarrow M$ is a G -equivariant principal K -bundle with G -invariant connection. This implies that Q is a $(G \times K)$ -manifold, M is a G -manifold, and the map $Q \rightarrow M$ is equivariant. (Note that in the following discussion, G will always act on the

base manifold M , and K will be the structure group for a principal bundle.) The above construction generalizes to give a stack morphism

$$\mathcal{E}_{\nabla}G \times_G M \xrightarrow{\Theta_G} \mathcal{E}_{\nabla}(G \times K) \times_{G \times K} Q,$$

and this gives the expected classical maps in cohomology and equivariant forms. In particular, we get the following maps between short exact sequences.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \frac{\Omega_{G \times K}^{*-1}(Q)}{\Omega_{G \times K}^{*-1}(Q)_{\mathbb{Z}}} & \longrightarrow & \check{H}_{G \times K}^*(Q) & \longrightarrow & H_{G \times K}^*(Q; \mathbb{Z}) \longrightarrow 0 \\ & & \Theta_G \downarrow & \uparrow q_{/K} & \Theta_G \downarrow & \uparrow q_{/K} & \uparrow \cong \\ 0 & \longrightarrow & \frac{\Omega_G^{*-1}(M)}{\Omega_G^{*-1}(M)_{\mathbb{Z}}} & \longrightarrow & \check{H}_G^*(M) & \longrightarrow & H_G^*(M; \mathbb{Z}) \longrightarrow 0 \\ \\ 0 & \longrightarrow & H_{G \times K}^{*-1}(Q; \mathbb{R}/\mathbb{Z}) & \longrightarrow & \check{H}_{G \times K}^*(Q) & \longrightarrow & \Omega_{G \times K}^*(Q)_{\mathbb{Z}} \longrightarrow 0 \\ & & \uparrow \cong & & \Theta_G \downarrow & \uparrow q_{/K} & \Theta_G \downarrow & \uparrow q_{/K} \\ 0 & \longrightarrow & H_G^{*-1}(M; \mathbb{R}/\mathbb{Z}) & \longrightarrow & \check{H}_G^*(M) & \longrightarrow & \Omega_G^*(M)_{\mathbb{Z}} \longrightarrow 0 \end{array}$$

Let us now give the precise construction in slightly greater generality. Assume that K is a normal subgroup of \tilde{G} with quotient $\tilde{G}/K \cong G$, and furthermore assume that we have fixed a splitting of the Lie algebra, i.e.,

$$(4.9) \quad \begin{aligned} 1 &\rightarrow K \hookrightarrow \tilde{G} \rightarrow G \rightarrow 1 \\ \tilde{\mathfrak{g}} &\cong \mathfrak{g} \oplus \mathfrak{k}. \end{aligned}$$

Such a decomposition of Lie algebras must exist since our groups are compact. Of primary interest is when $\tilde{G} = G \times K$, but we wish to also allow examples such as $SU(n) \hookrightarrow U(n) \rightarrow U(1)$ or $U(1) \hookrightarrow \text{Spin}^c(n) \rightarrow SO(n)$.

Definition 4.10. For $M \in G\text{-Man}$, let $\tilde{G}\text{-Bun}_{K, \nabla}(M)$ be the groupoid of \tilde{G} -equivariant principal K -bundles on M with invariant connection. An object $(Q, \Theta) \in \tilde{G}\text{-Bun}_{K, \nabla}(M)$ is a principal K -bundle with connection on M such that:

- $Q \in \tilde{G}\text{-Man}$ and $Q \xrightarrow{\pi} M$ is equivariant.
- $\Theta \in (\Omega^1(Q) \otimes \mathfrak{k})^{\tilde{G}}$.

For $(Q_i, \Theta_i) \in \tilde{G}\text{-Bun}_{K, \nabla}(M)$, a morphism $(Q_1, \Theta_1) \xrightarrow{\phi} (Q_2, \Theta_2)$ is a \tilde{G} -equivariant map $Q_1 \xrightarrow{\phi} Q_2$ which is an isomorphism of K -bundles with connection.

Remark 4.11. For $\tilde{G} = G \times K$, the condition $\Theta \in (\Omega^1(Q) \otimes \mathfrak{k})^{G \times K}$ can be rewritten as $k^*\Theta = \text{Ad}_{k^{-1}} \Theta$ and $g^*\Theta = \Theta$. Also, in this case we may refer to \tilde{G} -equivariant K -bundles as G -equivariant K -bundles. This is a standard convention, and we hope it does not cause any confusion.

Given an element $(Q, \Theta) \in \tilde{G}\text{-Bun}_{K,\nabla}(M)$, we want to define a natural map Θ_G

$$\mathcal{E}_{\nabla}\tilde{G} \times_{\tilde{G}} Q \begin{array}{c} \xleftarrow{\Theta_G} \\ \xrightarrow{q_{/K}} \end{array} \mathcal{E}_{\nabla}G \times_G M$$

such that the composition $q_K \circ \Theta_G$ is naturally isomorphic to the identity. We first describe the construction, hopefully emphasizing the geometric nature and naturalness. The details, which are relatively straightforward, are then checked in Proposition 4.16. While the induced maps in cohomology and equivariant forms that we recover are well-known, and the construction of Θ_G appears implicitly in [BoT], we do not know any references where this construction is done in general at the level of principal bundles with connection.

We now give a functor between the two groupoids produced when the stacks are evaluated on a test manifold X . Denoting $\Theta = \Theta_Q$ for added clarity, the map is given by the following construction.

$$(\mathcal{E}_{\nabla}G \times_G M)(X) \xrightarrow{\Theta_G(X)} (\mathcal{E}_{\nabla}\tilde{G} \times_{\tilde{G}} Q)(X)$$

$$\begin{array}{ccc} (Q, \Theta_Q) & \xrightarrow{\left(f^*Q, \pi_P^*\Theta_P \oplus (\tilde{f}^*\Theta_Q - \iota_{\pi_P^*\Theta_P}\tilde{f}^*\Theta_Q) \right)} & (Q, \Theta_Q) \\ \downarrow \pi_M & & \downarrow \pi_M \\ (P, \Theta_P) \xrightarrow{f} M & \xrightarrow{\pi'_X} & \begin{array}{ccc} \begin{array}{c} \downarrow \pi_P \\ (P, \Theta_P) \end{array} & \xrightarrow{\dots f \dots} & \begin{array}{c} \downarrow \pi_M \\ M \end{array} \\ \downarrow \pi_X & \swarrow & \downarrow \\ X & & X \end{array} \end{array}$$

Our new \tilde{G} -bundle is given by pulling back Q . The connection on f^*Q is given by the pulling back the connections on P and Q and subtracting a correction term. This extra term is defined to be the image

$$\begin{aligned} \Omega^1(\tilde{f}^*Q) \otimes \mathfrak{g} \otimes \Omega^1(\tilde{f}^*Q) \otimes \mathfrak{k} &\longrightarrow \Omega^1(\tilde{f}^*Q) \otimes \mathfrak{k} \\ \pi_P^*\Theta_P \otimes \tilde{f}^*\Theta_Q &\longmapsto \iota_{\pi_P^*\Theta_P}\tilde{f}^*\Theta_Q \end{aligned}$$

under the natural contraction $\mathfrak{g} \otimes \Omega^1(\tilde{f}^*Q) \xrightarrow{\iota} \Omega^0(\tilde{f}^*Q)$. While this term may initially seem obscure, it is necessary to ensure that we produce a connection on f^*Q , and it has a simple description in the Weil model.

Define the equivariant extension of the connection and curvature forms by

$$(4.12) \quad \Theta_G := \Theta - \iota_{\theta_{\mathfrak{g}}}\Theta \in (\mathbb{W}(\mathfrak{g}) \otimes \Omega(Q))^1 \otimes \mathfrak{k},$$

$$(4.13) \quad \Omega_G := d_G\Theta_G + \frac{1}{2}[\Theta_G \wedge \Theta_G] \in (\mathbb{W}(\mathfrak{g}) \otimes \Omega(Q))^2 \otimes \mathfrak{k}.$$

To clarify, $\iota_{\theta_{\mathfrak{g}}}\Theta \in \Lambda^1\mathfrak{g}^* \otimes \Omega^0(Q) \otimes \mathfrak{k}$ and it is evaluated on elements $\xi \in \mathfrak{g}$ by contracting the connection along the vector field in Q generated by the

G -action; i.e.,

$$\langle \iota_{\theta_{\mathfrak{g}}}\Theta, \xi \rangle := \iota_{\xi}\Theta \in \Omega^0(Q; \mathfrak{k}).$$

The \mathfrak{k} portion of the connection on \tilde{f}^*Q is simply the image of the equivariant connection Θ_G under the Weil homomorphism

$$\begin{aligned} W(\mathfrak{g}) \otimes \Omega(Q) \otimes \mathfrak{k} &\xrightarrow{(\pi_P^* \Theta_P)^* \otimes \tilde{f}^* \otimes 1} \Omega(\tilde{f}^*Q) \otimes \mathfrak{k} \\ \Theta_G = \Theta - \iota_{\theta_{\mathfrak{g}}}\Theta &\longmapsto \tilde{f}^*\Theta_Q - \iota_{\pi_P^* \Theta_P} \tilde{f}^*\Theta_Q. \end{aligned}$$

Note that the additional term $\iota_{\pi_P^* \Theta_P} \tilde{f}^*\Theta_Q$ uses the connection on the G -bundle P to detect the G -action on Q . This is key to understanding the equivariant Chern–Weil homomorphism in Section 5.

Remark 4.14. In the Cartan model for equivariant forms (see Section A.5), the equivariant connection and curvature simplify to

$$\begin{aligned} \Theta_G &\longleftrightarrow \Theta && \in S^0 \mathfrak{g}^* \otimes \Omega^1(Q) \otimes \mathfrak{k}, \\ \Omega_G &\longleftrightarrow \Omega - \iota_{\Omega_{\mathfrak{g}}}\Theta && \in \left(S^0 \mathfrak{g}^* \otimes \Omega^1(Q) + S^2 \mathfrak{g}^* \otimes \Omega^0(Q) \right) \otimes \mathfrak{k}, \end{aligned}$$

with Ω_G becoming the familiar equivariant curvature defined in [BeV, §2].

We proceed to check that Θ_G and Ω_G are equivariant forms and the above construction satisfies the desired properties.

Lemma 4.15. *For $(Q, \Theta) \in \tilde{G}\text{-Bun}_{K, \nabla}(M)$, the forms Θ_G and Ω_G are \tilde{G} -invariant and \mathfrak{g} -horizontal. In other words, they live in the subcomplex $(W(\mathfrak{g}) \otimes \Omega(Q) \otimes \mathfrak{k})_{\mathfrak{g}\text{-hor}}^{\tilde{G}}$. When $\tilde{G} = G \times K$, this can be written*

$$\Theta_G \in \Omega_G^1(Q; \mathfrak{k}), \quad \Omega_G \in \Omega_G^2(Q; \mathfrak{k}).$$

Proof. To see that Θ_G is \mathfrak{g} -horizontal, let $\xi \in \mathfrak{g}$. Then

$$\iota_{\xi}\Theta_G = \iota_{\xi}\Theta - \iota_{\theta_{\mathfrak{g}}(\xi)}\Theta = \iota_{\xi}\Theta - \iota_{\xi}\Theta = 0.$$

We show both terms in Θ_G are \tilde{G} -invariant. First note $\Theta \in (\Omega^1(Q) \otimes \mathfrak{k})^{\tilde{G}}$.

To check the term $\iota_{\theta_{\mathfrak{g}}}\Theta$, observe that $\theta_{\mathfrak{g}} \in (\Lambda^1 \mathfrak{g}^* \otimes \mathfrak{g})^G = (\Lambda^1 \mathfrak{g}^* \otimes \mathfrak{g})^{\tilde{G}}$, and the contraction map

$$\begin{aligned} \mathfrak{g} \otimes \Omega^1(Q) &\xrightarrow{\iota} \Omega^0(Q) \\ \xi \otimes \omega &\longmapsto \iota_{\xi}\omega \end{aligned}$$

is \tilde{G} -equivariant. Hence,

$$\iota_{\theta_{\mathfrak{g}}}\Theta \in (\Lambda^1 \mathfrak{g}^* \otimes \Omega^0(Q) \otimes \mathfrak{k})^{\tilde{G}},$$

so Θ_G is \mathfrak{g} -horizontal and \tilde{G} -invariant.

From the Leibniz rule for ι_{ξ} and the definition of tensor product of representations, it immediately follows that $[\Theta_G \wedge \Theta_G]$ is also \mathfrak{g} -horizontal and \tilde{G} -invariant. The standard identities (A.3) and (A.4) imply that $d_G \Theta_G$ is

also \tilde{G} -invariant and \mathfrak{g} -horizontal. Consequently, Ω_G is invariant and horizontal. \square

Proposition 4.16. *Assume $\tilde{G}/K \cong G$ satisfying (4.9). Let $(Q, \Theta) \in \tilde{G}\text{-Bun}_{K, \nabla}(M)$.*

- (1) *The above construction defines a natural morphism Θ_G*

$$\mathcal{E}_{\nabla} \tilde{G} \times_{\tilde{G}} Q \begin{array}{c} \xleftarrow{\Theta_G} \\ \xrightarrow{q/K} \end{array} \mathcal{E}_{\nabla} G \times_G M$$

such that the composition $q/K \circ \Theta_G$ is naturally isomorphic to the identity.

- (2) *The induced map*

$$\Omega(\mathcal{E}_{\nabla} \tilde{G} \times_{\tilde{G}} Q) \xrightarrow{\Theta_G^*} \Omega(\mathcal{E}_{\nabla} G \times_G M)$$

is equivalent to the homomorphism $\Theta_G^: \Omega_{\tilde{G}}(Q) \rightarrow \Omega_G(M)$ induced from*

$$\begin{aligned} \mathbb{W}(\mathfrak{g}) \otimes \mathbb{W}(\mathfrak{k}) \otimes \Omega(Q) &\xrightarrow{\Theta_G^*} \mathbb{W}(\mathfrak{g}) \otimes \Omega(Q) \\ \alpha \cdot \beta \cdot \gamma &\longmapsto \alpha \cdot \beta(\Theta_G) \cdot \gamma. \end{aligned}$$

- (3) *The induced map in cohomology is the standard isomorphism*

$$H_{\tilde{G}}^*(Q; -) \xrightarrow{\cong} H_G^*(M; -).$$

Proof. For part (1), we must check that our construction defines a principal \tilde{G} -bundle with connection. Since $P \xrightarrow{f} M \xleftarrow{\pi_M} Q$ are equivariant maps between \tilde{G} -manifolds, the pullback f^*Q is naturally a \tilde{G} -manifold. The freeness of the \tilde{G} -action follows easily by the following argument. Suppose $\tilde{g} \in \tilde{G}$ satisfies $\tilde{g}(p, q) = (p, q)$ for some point in $f^*Q \subset P \times Q$; then $gp = p \in P$ implies $g = 1$ and hence $\tilde{g} \in K$. Since K acts freely on Q , then $\tilde{g} = 1$. Hence, $f^*Q \rightarrow X$ is a principal \tilde{G} -bundle. To show that

$$(4.17) \quad \Theta_{f^*Q} = \pi_P^* \Theta_P \oplus (\tilde{f}^* \Theta_Q - \iota_{\pi_P^* \Theta_P} \tilde{f}^* \Theta_Q) \in \Omega^1(f^*Q) \otimes (\mathfrak{g} \oplus \mathfrak{k})$$

is a connection, we must check it is \tilde{G} -equivariant and restricts fiberwise to the Maurer–Cartan 1-form on \tilde{G} . By Lemma 4.15, we see that $\tilde{f}^* \Theta_Q - \iota_{\pi_P^* \Theta_P} \tilde{f}^* \Theta_Q$ must be \tilde{G} -invariant and \mathfrak{g} -horizontal, since it is the image of Θ_G under the map

$$(\mathbb{W}(\mathfrak{g}) \otimes \Omega(Q) \otimes \mathfrak{k})_{\mathfrak{g}\text{-hor}}^{\tilde{G}} \longrightarrow (\Omega(f^*Q) \otimes \mathfrak{k})_{\mathfrak{g}\text{-hor}}^{\tilde{G}}.$$

This, combined with the fact that $\Theta_P \in (\Omega^1(P) \otimes \mathfrak{g})^G$, implies that (4.17) is an element of $(\Omega^1(f^*Q) \otimes (\mathfrak{g} \oplus \mathfrak{k}))^{\tilde{G}}$. Similarly, if $\xi_1 \oplus \xi_2 \in \mathfrak{g} \oplus \mathfrak{k}$, then

$$\begin{aligned} & \iota_{\xi_1 \oplus \xi_2} (\pi_P^* \Theta_P \oplus (\tilde{f}^* \Theta_Q - \iota_{\pi_P^* \Theta_P} \tilde{f}^* \Theta_Q)) \\ &= (\iota_{\xi_1} \Theta_P) \oplus (\iota_{\xi_2} \Theta_Q) + 0 \oplus \iota_{\xi_1} (\tilde{f}^* \Theta_Q - \iota_{\pi_P^* \Theta_P} \tilde{f}^* \Theta_Q) \\ &= \xi_1 \oplus \xi_2. \end{aligned}$$

Therefore, the morphism of stacks Θ_G is well-defined. The map π_P gives a natural isomorphism $(f^*Q)/K \cong P$ compatible with the connections. Hence, $q/K \circ \Theta_G$ is naturally isomorphic to the identity on $\mathcal{E}_{\nabla} G \times_G M$.

The proof of part (2) is given by tracing through the Freed–Hopkins isomorphism, discussed in Section 3.2, and the definition of our map. Essentially, we must show that the diagram

$$\begin{array}{ccc} \Omega_{G \times K}(Q) & \xrightarrow{\cong} & \Omega(\mathcal{E}_{\nabla}(G \times K) \times_{G \times K} Q) \\ \downarrow \Theta_G^* & & \downarrow \Theta_G^* \\ \Omega_G(M) & \xrightarrow{\cong} & \Omega(\mathcal{E}_{\nabla} G \times_G M) \end{array}$$

commutes by evaluating on a test manifold X . Consider a homogeneous element $\alpha\beta\gamma \in \Omega_{G \times K}(Q) \subset W(\mathfrak{g}) \otimes W(\mathfrak{k}) \otimes \Omega(Q)$. We first check the clockwise direction. Given $X \rightarrow \mathcal{E}_{\nabla} G \times_G M$, we have

$$\begin{array}{ccc} (P, \Theta_P) & \xrightarrow{f} & M \\ \downarrow & & \\ X & & \end{array} \quad \mapsto \quad \begin{array}{ccc} (f^*Q, \Theta_{f^*Q}) & \xrightarrow{\tilde{f}} & (Q, \Theta) \\ \downarrow \pi_P & & \downarrow \\ (P, \Theta_P) & \xrightarrow{f} & M \\ \downarrow & & \\ X & & \end{array}$$

where

$$\Theta_{f^*Q} = \pi_P^* \Theta_P \oplus (\tilde{f}^* \Theta - \iota_{\pi_P^* \Theta_P} \tilde{f}^* \Theta).$$

Therefore, $\alpha\beta\gamma \in \Omega_{G \times K}(Q)$ gets mapped to

$$\alpha(\pi_P^* \Theta_P) \wedge \beta(\tilde{f}^* \Theta - \iota_{\pi_P^* \Theta_P} \tilde{f}^* \Theta) \wedge \tilde{f}^* \gamma \in \Omega(f^*Q)_{G \times K\text{-basic}} \cong \Omega(X).$$

Going counter-clockwise, we first have that $\alpha\beta\gamma \mapsto \alpha\beta(\Theta_G)\gamma$, which lives in

$$(W(\mathfrak{g}) \otimes \Omega(Q))_{G \times K\text{-basic}} \cong (W(\mathfrak{g}) \otimes \Omega(M))_{G\text{-basic}}.$$

When we evaluate on $X \leftarrow (P, \Theta) \xrightarrow{f} M$,

$$\alpha\beta(\Theta_G)\gamma \quad \mapsto \quad \alpha(\Theta_P)\beta(f^* \Theta - \iota_{\Theta_P} f^* \Theta) f^* \gamma \in \Omega(P)_{G\text{-basic}} \cong \Omega(X).$$

Pulling this back up to f^*Q via π_P^* gives us the form from the other direction. Hence, they restrict to the same form in $\Omega(X)$.

Finally, part (1) implies that $\Theta_G^* \circ q_K^* = \text{Id}$, and we know from (4.8) that $q_{/K}^*$ is an isomorphism in ordinary cohomology. Therefore, part (3) follows immediately. \square

4.3. Uniqueness of \check{H}_G^* . A natural question is whether one can give an axiomatic characterization of the functor \check{H}_G^* . We now show that the character diagram, combined with the maps $\check{H}_G^*(P) \xrightarrow{\Theta^*} \check{H}^*(M)$ for principal G -bundles with connection, uniquely characterize our differential extension of Borel equivariant cohomology. The idea is to use Θ^* to regard $\check{H}_G^*(M)$ a subgroup of $\check{H}^*(EG \times_G M)$ and invoke the Simons–Sullivan axiomatic characterization of differential cohomology. It is unknown to the author if the uniqueness result holds without assuming the additional structure maps Θ^* .

Proposition 4.18. *If \mathbb{H}_G^* is a functor from G -manifolds to graded abelian groups satisfying parts A and C of Theorem 1.1, then there is a natural equivalence $\mathbb{H}_G^* \rightarrow \check{H}_G^*$ that commutes with the identity map on all other functors in the character diagram.*

Proof. Assume that \mathbb{H}_G^* satisfies Theorem 1.1 parts A and C. As described in Section A.4, let $(EG, \Theta_{EG}) \rightarrow BG$ be the universal bundle with connection obtained as a direct limit of smooth manifolds. The projection map $EG \times M \rightarrow M$ is G -equivariant and can be combined with the universal connection to produce

$$\mathbb{H}_G^*(M) \rightarrow \mathbb{H}_G^*(EG \times M) \xrightarrow{\Theta_{EG}^*} \mathbb{H}^*(EG \times_G M).$$

On ordinary cohomology, this map is an isomorphism. Since the Weil homomorphism $\Omega_G(M) \rightarrow \Omega(EG \times_G M)$ is injective (Lemma A.17), it follows from the short exact sequence (SES 2) that $\mathbb{H}_G^*(M)$ is naturally an abelian subgroup of $\mathbb{H}^*(EG \times_G M)$. The same argument shows $\check{H}_G^*(M)$ is a subgroup of $\check{H}^*(EG \times_G M)$.

In [SS], Simons–Sullivan show the character diagram uniquely characterizes ordinary differential cohomology, and there exists a unique natural transformation $\mathbb{H}^* \xrightarrow{\cong} \check{H}^*$. This isomorphism combines with (SES 2) to give the following.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \frac{\Omega_G^{*-1}(M)}{\Omega_G^{*-1}(M)_{\mathbb{Z}}} & \longrightarrow & \mathbb{H}_G^*(M) & \longrightarrow & H_G^*(M; \mathbb{Z}) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & \frac{\Omega^{*-1}(EG \times_G M)}{\Omega^{*-1}(EG \times_G M)_{\mathbb{Z}}} & \longrightarrow & \check{H}^*(EG \times_G M) & \longrightarrow & H^*(EG \times_G M; \mathbb{Z}) \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \cong \\ 0 & \longrightarrow & \frac{\Omega_G^{*-1}(M)}{\Omega_G^{*-1}(M)_{\mathbb{Z}}} & \longrightarrow & \check{H}_G^*(M) & \longrightarrow & H_G^*(M; \mathbb{Z}) \longrightarrow 0. \end{array}$$

It immediately follows that $\mathbb{H}_G^*(M)$ and $\check{H}_G^*(M)$ can be naturally considered as the same abelian subgroup of $\check{H}^*(EG \times_G M)$, and this gives us our desired natural isomorphism. \square

5. Equivariant Chern–Weil theory

The constructions in Propositions 4.4 and 4.16 combine to give a geometric interpretation of equivariant Chern–Weil theory. For simplicity, let $\tilde{G} = G \times K$ in the following preliminary discussion.

Suppose that M is a G -manifold and $(Q, \Theta) \rightarrow M$ is a G -equivariant principal K -bundle with G -invariant connection; i.e., $(Q, \Theta) \in G\text{-Bun}_{K, \nabla}(M)$. We briefly explain two classical ways to construct equivariant characteristic classes. The first construction is purely topological and does not use the connection. Consider the principal K -bundle $EG \times_G Q \rightarrow EG \times_G M$. The choice of classifying map

$$(5.1) \quad EG \times_G M \longrightarrow BK.$$

induces a homomorphism

$$H^*(BK; A) \longrightarrow H_G^*(M; A)$$

for any abelian group A . The second method, due originally to Berline–Vergne [BeV], uses the connection to construct an equivariant differential form. Given $\omega \in (S^n \mathfrak{k}^*)^K \cong H^{2n}(BK; \mathbb{R})$, define the equivariant Chern–Weil form

$$\omega(\Theta_G) := \omega(\Omega_G^{\wedge n}) \in \Omega_G^{2n}(M).$$

As noted in Remark 4.14, this becomes

$$\omega((\Omega - \iota_{\Omega_{\mathfrak{g}}} \Theta)^{\wedge n}) \in (S\mathfrak{g}^* \otimes \Omega(M))^G$$

in the more commonly used Cartan model. While the form $\omega(\Theta_G)$ depends on the connection Θ , its class in $H_G^{2n}(M; \mathbb{R})$ does not. These give equivariant Chern classes and forms when $K = U(n)$, and equivariant Pontryagin classes and forms when $K = SO(n)$. In [BoT], Bott–Tu use the universal connection on EG to show these two constructions give the same classes in $H_G^*(M; \mathbb{R})$. The following construction is similar to theirs, but we replace EG with $\mathcal{E}_{\nabla}G$ and view everything in the world of stacks.

Suppose that $(Q, \Theta) \in G\text{-Bun}_{K, \nabla}(M)$. The following maps compose to give a refinement of (5.1) to stacks.

$$(5.2) \quad \mathcal{E}_{\nabla}G \times_G M \xrightarrow{\Theta_G} \mathcal{E}_{\nabla}(G \times K) \times_{G \times K} Q \longrightarrow \mathcal{B}_{\nabla}(G \times K) \xrightarrow{q/G} \mathcal{B}_{\nabla}K.$$

The first map is given by the bundle pullback construction of Section 4.2, the second is induced by the equivariant map $Q \rightarrow \text{pt}$, and the third map is induced by the quotient $G \times K \xrightarrow{/G} K$. When applied to a test manifold X ,

the composition performs the following construction.

$$\begin{array}{ccc}
 & (Q, \Theta_Q) & \\
 & \downarrow \pi_M & \\
 (P, \Theta_P) \xrightarrow{f} & M & \mapsto (f^*Q, \pi_P^*\Theta_P \oplus (\tilde{f}^*\Theta_Q - \iota_{\Theta_P}\tilde{f}^*\Theta_Q))/G \\
 \downarrow \pi_X & & \downarrow \\
 & & X
 \end{array}$$

It pulls back Q , modifies the connection to create a $G \times K$ -bundle with connection, and then quotients by G .

Theorem 5.3. *Associated to $(Q, \Theta) \in G\text{-Bun}_{K,\nabla}(M)$ is a natural morphism*

$$\mathcal{E}_{\nabla}G \times_G M \xrightarrow{(Q_G, \Theta_G)} \mathcal{B}_{\nabla}K,$$

such that the induced map

$$\check{H}_K^* \xrightarrow{(Q_G, \Theta_G)^*} \check{H}_G^*(M)$$

refines the traditional equivariant characteristic classes and forms as described above.

Proof. We know the morphism $\mathcal{E}_{\nabla}G \times_G M \rightarrow \mathcal{B}_{\nabla}K$ is well-defined, since it is defined in (5.2) as the composition of three morphisms.

We now must show that at the level of cohomology, $(Q_G, \Theta_G)^*$ is the homomorphism ψ^* , where $EG \times_G M \xrightarrow{\psi} BK$ is a classifying map for $EG \times_G Q \rightarrow EG \times_G M$. By Proposition 6.7, the maps

$$EG \times_G M \rightarrow \mathcal{E}_{\nabla}G \times_G M \rightarrow \mathcal{E}G \times_G M \quad \text{and} \quad BK \rightarrow \mathcal{B}_{\nabla}K \rightarrow \mathcal{B}K$$

induce isomorphisms in cohomology. Therefore, we must check that the diagram

$$(5.4) \quad \begin{array}{ccc}
 EG \times_G M & \longrightarrow & \mathcal{E}G \times_G M \\
 \psi \downarrow & & \downarrow Q_G \\
 BK & \longrightarrow & \mathcal{B}K,
 \end{array}$$

where the two horizontal morphisms given by viewing $EG \rightarrow BG$ and $EK \rightarrow BK$ as principal bundles, is commutative up to isomorphism.

We evaluate on a test manifold X and trace through where $X \xrightarrow{s} EG \times_G M$ is sent, using the following diagram to identify maps.

$$\begin{array}{ccccc}
 \tilde{s}^*(\Pi_M^* Q) & \longrightarrow & \Pi_M^* Q & \longrightarrow & Q \\
 \downarrow & & \downarrow & & \downarrow \\
 s^*(EG \times M) & \xrightarrow{\tilde{s}} & EG \times M & \xrightarrow{\Pi_M} & M \\
 \downarrow & & \downarrow & & \downarrow \\
 X & \xrightarrow{s} & EG \times_G M & &
 \end{array}$$

Going clockwise in (5.4), we have

$$X \xrightarrow{s} EG \times_G M \mapsto \begin{array}{ccc} s^*(EG \times M) & \xrightarrow{\Pi_M \circ \tilde{s}} & M \\ \downarrow & & \downarrow \\ X & & X \end{array} \mapsto \begin{array}{ccc} ((\Pi_M \circ \tilde{s})^* Q)/G & & \\ \downarrow & & \\ X & & \end{array},$$

and the evident isomorphism $\Pi_M^* Q \cong EG \times Q$ gives

$$((\Pi_M \circ \tilde{s})^* Q)/G \cong (\tilde{s}^*(EG \times Q))/G \cong s^*(EG \times_G Q).$$

Going counter-clockwise in (5.4), we have

$$X \xrightarrow{s} EG \times_G M \mapsto X \xrightarrow{\psi \circ s} BK \mapsto \begin{array}{ccc} (\psi \circ s)^* EK & & \\ \downarrow & & \\ X & & \end{array}.$$

Since

$$(\psi \circ s)^* EK \cong s^*(\psi^* EK) \cong s^*(EG \times_G Q),$$

the diagram (5.4) commutes up to isomorphism.

For differential forms, we trace through the composition (5.2), which is given by the restriction of the following to the basic sub-complexes

$$W(\mathfrak{k}) \rightarrow W(\mathfrak{g}) \otimes W(\mathfrak{k}) \rightarrow W(\mathfrak{g}) \otimes W(\mathfrak{k}) \otimes \Omega(Q) \xrightarrow{\Theta_G^*} W(\mathfrak{g}) \otimes \Omega(M).$$

The first two maps are given by the mapping $W(\mathfrak{k})$ into its factor in the tensor product, so the composition is simply

$$\begin{aligned}
 (S\mathfrak{k}^*)^K = W(\mathfrak{k})_{\text{basic}} &\longrightarrow (W(\mathfrak{g}) \otimes \Omega(M))_{\text{basic}} = \Omega_G(M) \\
 \omega &\longmapsto \omega(\Theta_G),
 \end{aligned}$$

which is precisely the equivariant Chern–Weil form in the Weil model. \square

One can also construct equivariant extensions of the Chern–Simons forms. As explained in (A.13), to $\omega \in (S^k \mathfrak{k}^*)^K$ is naturally associated a Chern–Simons form $CS_\omega \in W^{2k-1}(\mathfrak{k})^K$ satisfying $d_W CS_\omega = \omega$. Given

$$(Q, \Theta) \in G\text{-Bun}_{K, \nabla}(M),$$

we define the equivariant Chern–Simons form to be the image of CS_ω under the equivariant equivariant Weil homomorphism

$$(5.5) \quad \begin{aligned} \mathbb{W}^{2k-1} &\xrightarrow{\Theta_G^*} \Omega_G^{2k-1}(Q) \\ \text{CS}_\omega &\mapsto \text{CS}_\omega(\Theta_G). \end{aligned}$$

As in the case of differential characters, the Chern–Simons forms are closely related to the differential refinements of characteristic classes.

Proposition 5.6. *Let $\check{\omega} \in \check{H}^{2k}(\mathcal{B}_\nabla K) \cong H^{2k}(BK; \mathbb{Z})$ be a differential refinement of $\omega \in (S^k \mathfrak{k}^*)^K$. Suppose that $(Q, \Theta) \in G\text{-Bun}_{K, \nabla}(M)$ admits a G -equivariant section $s : M \rightarrow Q$. Then*

$$[s_G^* \text{CS}_\omega(\Theta_G)] = \check{\omega}(\Theta_G) \in \check{H}_G^{2k}(M).$$

Proof. First note that because EG is contractible, the character diagram for $\mathcal{E}_\nabla G$ gives isomorphisms

$$\begin{array}{ccc} & \check{H}^{2k}(\mathcal{E}_\nabla G) & \\ \cong \nearrow & & \searrow \cong \\ \frac{\mathbb{W}^{2k-1}(\mathfrak{k})}{d_W \mathbb{W}^{2k-2}(\mathfrak{k})} & \xrightarrow{d_W} & \mathbb{W}^{2k}(\mathfrak{k})_0. \end{array}$$

This gives the universal property $\pi^* \check{\omega} = [\text{CS}_\omega] \in \check{H}^{2k}(\mathcal{E}_\nabla K)$.

The construction (5.2) readily generalizes to the following diagram,

$$\begin{array}{ccc} \mathcal{E}_\nabla G \times_G Q & \longrightarrow & \mathcal{E}_\nabla K \\ \downarrow \Big) s_G & & \downarrow \pi \\ \mathcal{E}_\nabla G \times_G M & \xrightarrow{(Q_G, \Theta_G)} & \mathcal{B}_\nabla K \end{array}$$

giving us two ways of writing the morphism $\mathcal{E}_\nabla G \times_G M \rightarrow \mathcal{B}_\nabla K$. The bottom morphism induces $\check{\omega}(\Theta_G)$; the other one gives

$$\begin{aligned} \check{H}(\mathcal{B}_\nabla K) &\longrightarrow \check{H}(\mathcal{E}_\nabla G) \longrightarrow \check{H}(\mathcal{E}_\nabla G \times_G Q) \longrightarrow \check{H}(\mathcal{E}_\nabla G \times_G M) \\ \check{\omega} &\longmapsto [\text{CS}_\omega] \longmapsto [\text{CS}_\omega(\Theta_G)] \longmapsto [s_G^* \text{CS}_\omega(\Theta_G)]. \end{aligned}$$

□

Remark 5.7. In the statement of Theorem 5.3, we assumed that the total space of the bundle Q had an action of $G \times K$. Suppose instead that $(Q, \Theta) \in G\text{-Bun}_{\tilde{G}/K, \nabla}(M)$, where $\tilde{G}/K \cong G$ with a fixed Lie algebra splitting (4.9). It no longer makes sense to quotient by G at the end, but we still have the composition

$$\mathcal{E}_\nabla G \times_G M \xrightarrow{\Theta_G} \mathcal{E}_\nabla \tilde{G} \times_{\tilde{G}} Q \longrightarrow \mathcal{B}_\nabla \tilde{G}.$$

This induces differential equivariant characteristic classes via the homomorphism

$$\check{H}_{\tilde{G}}^* \longrightarrow \check{H}_G^*(M).$$

Note that Proposition 5.6 would not be applicable unless there exists an isomorphism $\tilde{G} \cong G \times K$.

5.1. Equivariant differential characters. One convenient description of differential cohomology is via differential characters [CheeS]. An element $\check{x} \in \check{H}^n(M)$, with curvature $R(\check{x}) = \omega$, associates to every smooth singular $(n-1)$ -cycle $X_{n-1} \xrightarrow{z} M$ an element $\langle \check{x}, z \rangle \in \mathbb{R}/\mathbb{Z}$; if $z = \partial \tilde{z}$ for some smooth n -chain $Y_n \xrightarrow{\tilde{z}} M$, then

$$\langle \check{x}, z \rangle = \int_{Y_n} \tilde{z}^* \omega \pmod{\mathbb{Z}}.$$

In degree 2, an \mathbb{R}/\mathbb{Z} -bundle with connection on M naturally defines an element of $\check{H}^2(M)$ via its holonomy. More generally, evaluating a differential character on a $(n-1)$ -cycle can be thought of as the holonomy for some higher gerbe or abelian gauge field.

This idea generalizes to the equivariant setting, with classes in $\check{H}_G^n(M)$ giving “differential characters” on the stack $\mathcal{E}_{\nabla}G \times_G M$. This type of structure naturally appears in the physics literature as WZW terms for gauged sigma models [FOS]. We state the main idea here, but we do not give enough details on singular cycles to consider this a definition of $\check{H}_G^*(M)$.

Assume that X_{n-1} is a closed smooth manifold. To any map

$$X_{n-1} \xrightarrow{z} \mathcal{E}_{\nabla}G \times_G M,$$

a class $\check{x} \in \check{H}_G^n(M)$ associates an “equivariant holonomy” in \mathbb{R}/\mathbb{Z} , as seen in the following diagram.

$$\begin{array}{ccc} (P, \Theta) \xrightarrow{f} M & & \check{H}_G^n(P) \xleftarrow{f_G^*} \check{H}_G^n(M) \\ \downarrow & \mapsto & \downarrow \Theta^* \\ X_{n-1} & & \check{H}^n(X_{n-1}) \cong \mathbb{R}/\mathbb{Z} \end{array}$$

If the cycle is a boundary, with $[(P, \Theta) \rightarrow X_{n-1}] = \partial [(P', \Theta') \rightarrow Y_n]$ and f extending to a map $P' \xrightarrow{\tilde{f}} M$, then the holonomy can be computed by integrating the equivariant curvature $\omega = R(\check{x}) \in \Omega_G^n(M)$

$$\langle \check{x}, X_{n-1} \leftarrow (P, \Theta) \xrightarrow{f} M \rangle = \int_{Y_n} (\Theta^* \otimes \tilde{f}^*) \omega \pmod{\mathbb{Z}}.$$

Furthermore, these equivariant holonomies are gauge-invariant; if two objects in the groupoid $\text{Map}(X_{n-1}, \mathcal{E}_{\nabla}G \times_G M)$ are isomorphic, they associate to $\check{x} \in \check{H}_G^n(M)$ equal elements in \mathbb{R}/\mathbb{Z} .

Remark 5.8. The entire above discussion concerning equivariant characters immediately generalizes from $\mathbb{Z} \subset \mathbb{R}$ to $\Lambda \subset V$, where elements of $\check{H}_G^*(M; \Lambda)$ give characters valued in V/Λ .

5.2. Geometric models in low degrees.

Proposition 5.9. *There are natural isomorphisms*

$$\begin{aligned} \check{H}_G^0(M) &\cong H^0(M/G; \mathbb{Z}), \\ \check{H}_G^1(M) &\cong \check{H}^1(M)^G \cong C^\infty(M, \mathbb{R}/\mathbb{Z})^G. \end{aligned}$$

Proof. The short exact sequence (SES 1) implies that $\check{H}_G^0(M) \cong \Omega_G^0(M)_{\mathbb{Z}}$. Unraveling this, we obtain the following chain of equalities and isomorphisms,

$$\check{H}_G^0(M) \cong \Omega_G^0(M)_{\mathbb{Z}} = C^\infty(M, \mathbb{Z})^G \cong C^0(M/G, \mathbb{Z}) \cong H^0(M/G; \mathbb{Z}).$$

In degree one, consider the homomorphism $\check{H}_G^1(M) \rightarrow C^\infty(M, \mathbb{R}/\mathbb{Z})^G$ defined as follows using the equivariant differential character construction of Section 5.1. Given $\check{x} \in \check{H}_G^1(M)$ and a point $m \in M$, define the element $F_{\check{x}}(M) \in \mathbb{R}/\mathbb{Z}$ by evaluating on the G -orbit of m ,

$$F_{\check{x}}(m) := \left\langle \check{x}, \text{pt} \leftarrow G \xrightarrow{G \cdot m} M \right\rangle.$$

The gauge invariance of the equivariant character implies that the function $F_{\check{x}}$ is G -invariant. Furthermore, $\Omega_G^1(M)_{\mathbb{Z}} \cong \Omega^1(M)_{\mathbb{Z}}^G$, and integration shows that $F_{\check{x}}$ is a smooth function with derivative $R(\check{x})$. Hence, $F_{\check{x}} \in C^\infty(M, \mathbb{R}/\mathbb{Z})^G$.

Note that the kernel of $C^\infty(M, \mathbb{R}/\mathbb{Z})^G \xrightarrow{d} \Omega^1(M)_{\mathbb{Z}}^G$ is given by the locally constant functions

$$\text{Ker}(d) = C^\infty(M, \mathbb{R}/\mathbb{Z})_{\text{cl}}^G \cong H^0(M/G; \mathbb{R}/\mathbb{Z}) \cong H_G^0(M; \mathbb{R}/\mathbb{Z}).$$

Therefore, the homomorphism of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_G^0(M; \mathbb{R}/\mathbb{Z}) & \longrightarrow & \check{H}_G^1(M) & \longrightarrow & \Omega_G^1(M)_{\mathbb{Z}} \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & \text{Ker}(d) & \longrightarrow & C^\infty(M, \mathbb{R}/\mathbb{Z})^G & \longrightarrow & \Omega^1(M)_{\mathbb{Z}}^G \longrightarrow 0 \end{array}$$

gives our desired isomorphism via the Five Lemma. □

Proposition 5.10. *In degree two, there is a natural isomorphism*

$$\check{H}_G^2(M) \cong \pi_0(G\text{-Bun}_{\mathbb{R}/\mathbb{Z}, \nabla}(M));$$

i.e., G -equivariant \mathbb{R}/\mathbb{Z} -bundles with connection on M , modulo equivariant isomorphisms that preserve connection, are in bijection with elements of $\check{H}_G^2(M)$.

Proof. Let $\check{c} \in \check{H}_{\mathbb{R}/\mathbb{Z}}^2 \cong H^2(B\mathbb{R}/\mathbb{Z}; \mathbb{Z}) \cong \mathbb{Z}$ be the standard generator. The equivariant Chern–Weil construction of Theorem 5.3 gives us a map

$$(5.11) \quad \begin{aligned} G\text{-Bun}_{\mathbb{R}/\mathbb{Z}, \nabla}(M) &\longrightarrow \check{H}_G^2(M) \\ (Q, \Theta) &\longmapsto \check{c}(Q_G, \Theta_G). \end{aligned}$$

Multiplication in the abelian group \mathbb{R}/\mathbb{Z} makes $G\text{-Bun}_{\mathbb{R}/\mathbb{Z}, \nabla}(M)$ into a Picard groupoid (this symmetric monoidal structure corresponds to the tensor product of line bundles), and (5.11) is symmetric monoidal. Furthermore, the map only depends on the isomorphism class of (Q, Θ) in $G\text{-Bun}_{\mathbb{R}/\mathbb{Z}, \nabla}(M)$.

Hence, the map (5.11) descends to a homomorphism of abelian groups $\pi_0(G\text{-Bun}_{\mathbb{R}/\mathbb{Z}, \nabla}(M)) \rightarrow \check{H}_G^2(M)$. And, the characteristic class $c(Q_G) \in H_G^2(M; \mathbb{Z})$ does not depend on the connection Θ . This gives the following homomorphism of short exact sequences.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker } \Pi & \longrightarrow & \pi_0 [G\text{-Bun}_{\mathbb{R}/\mathbb{Z}, \nabla}(M)] & \xrightarrow{\Pi} & \pi_0 [G\text{-Bun}_{\mathbb{R}/\mathbb{Z}}(M)] \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \frac{\Omega_G^1(M)}{\Omega_G^1(M)_{\mathbb{Z}}} & \longrightarrow & \check{H}_G^2(M) & \longrightarrow & H_G^2(M; \mathbb{Z}) \longrightarrow 0 \end{array}$$

We now show that each of the above vertical maps is an isomorphism.

It is a classical result [LMS] that, up to isomorphism, equivariant bundles with abelian structure group are classified by Borel cohomology; i.e., $\pi_0(G\text{-Bun}_A(M)) \cong [EG \times_G M, BA]$ when A is abelian. This implies the right vertical map is an isomorphism.

The G -invariant connections on the trivial bundle $M \times \mathbb{R}/\mathbb{Z}$ are in bijection with $\Omega^1(M)^G$. Such a connection form is gauge equivalent to the trivial connection if and only if it is the derivative of a gauge transformation $C^\infty(M, \mathbb{R}/\mathbb{Z})^G \cong \check{H}_G^1(M)$; i.e., if it lives in $\Omega^1(M)_{\mathbb{Z}}^G$. Hence,

$$\text{Ker } \Pi \cong \frac{\Omega^1(M)^G}{\Omega^1(M)_{\mathbb{Z}}^G} \xrightarrow{\cong} \frac{\Omega_G^1(M)}{\Omega_G^1(M)_{\mathbb{Z}}},$$

giving that the left vertical arrow is an isomorphism. The proposition now follows by the Five Lemma. \square

6. Details of the simplicial sheaf construction

This section contains a more detailed account of Section 3.3. The results essentially follow by combining the work of [FreH] and [BunNV]; the reader may refer to these works, along with [Lur, Chapter 5] for further details. Let us briefly explain our notation.

We will use simplicial sets in order to deal with groupoids, nonnegatively graded chain complexes, and topological spaces simultaneously. Let $\infty\mathbf{Gpd}$ be the $(\infty, 1)$ -category of “ ∞ -groupoids,” taken here to be the full simplicially enriched subcategory of simplicial sets spanned by Kan complexes. The category $\infty\mathbf{Gpd}$ is naturally equivalent, as an $(\infty, 1)$ -category, to the topologically enriched category \mathbf{Top} of topological spaces with the homotopy type of a CW complex. Though it is not strictly necessary, we use this equivalence $\infty\mathbf{Gpd} \simeq \mathbf{Top}$ to make certain statements easier to read. Any groupoid is naturally a Kan complex via the nerve construction N , and the Dold–Kan correspondence Γ makes a nonnegatively graded chain complex

into a simplicial abelian group. These structures, together with the singular functor \mathcal{S} and geometric realization $||$, fit into the following picture.

$$\begin{array}{ccc}
 & \text{Top} & \\
 & \uparrow \downarrow \mathcal{S} & \\
 & || \simeq & \\
 \text{Gpd} & \xrightarrow{\text{N}} \infty\text{Gpd} & \xleftarrow{\Gamma} \text{Ch}_{\geq 0}
 \end{array}$$

Let $\text{PShv}_{\infty\text{Gpd}} = \text{Fun}(\text{Man}^{\text{op}}, \infty\text{Gpd})$ be the $(\infty, 1)$ -category of ∞Gpd -valued presheaves on manifolds. A functor $\mathcal{F} \in \text{PShv}_{\infty\text{Gpd}}$ is a sheaf if it satisfies the following descent condition: for any covering $\mathcal{U} \rightarrow X$, the canonical map $\mathcal{F}(X) \xrightarrow{\simeq} \text{holim}_{\Delta} \mathcal{F}(\mathcal{U}^{\bullet})$ is an equivalence. The full subcategory of simplicial sheaves is denoted $\text{Shv}_{\infty\text{Gpd}}$, and there is a sheafification functor L forming the adjunction

$$L : \text{PShv}_{\infty\text{Gpd}} \rightleftarrows \text{Shv}_{\infty\text{Gpd}} : \text{inclusion.}$$

We continue to suppress the embedding of $\text{Set} \hookrightarrow \text{Gpd}$, but we use the nerve symbol to denote $\text{N} : \text{Shv}_{\text{Gpd}} \rightarrow \text{Shv}_{\infty\text{Gpd}}$. As in previous sections, we will use M to denote both a manifold M and the stack it represents, but we will begin using the notation \underline{M} within proofs for added clarity. For $Y \in \text{Top}$, we define $\text{Sing}_* Y \in \text{Shv}_{\infty\text{Gpd}}$ by $\text{Sing}_* Y(X) = \mathcal{S}(Y^X)$, the singular complex of the space of continuous maps from $X \rightarrow Y$. Let $\mathcal{K}(A, n) \in \text{Shv}_{\infty\text{Gpd}}$ denote any simplicial sheaf equivalent to $\text{Sing}_* K(A, n)$.

We now recount the results we need from the theory of homotopy-invariant sheaves developed by Bunke–Nikolaus–Völkl in [BunNV]. Their results hold for general \mathcal{C} -valued sheaves on manifolds, where \mathcal{C} is an $(\infty, 1)$ -category, but we have specialized them to the case of $\mathcal{C} = \infty\text{Gpd} \simeq \text{Top}$. A sheaf \mathcal{F} is homotopy-invariant if, for all manifolds X , the projection $X \times I \rightarrow X$ induces an equivalence

$$\mathcal{F}(X) \xrightarrow{\simeq} \mathcal{F}(X \times I).$$

Let $\text{Shv}_{\infty\text{Gpd}}^{\text{h}}$ denote the full subcategory of homotopy-invariant sheaves. The following results are found in Proposition 2.6, Equation (68), and Lemma 7.13 of [BunNV].

- The functor Sing_* gives an equivalence $\text{Top} \xrightarrow{\simeq} \text{Shv}_{\infty\text{Gpd}}^{\text{h}}$; the inverse is given by evaluating at a point and taking the geometric realization. There is also a homotopification functor \mathcal{H} , and these form the following adjunctions.

$$\text{Shv}_{\infty\text{Gpd}} \begin{array}{c} \xrightarrow{\mathcal{H}} \\ \xleftarrow{\text{Sing}_*} \end{array} \text{Shv}_{\infty\text{Gpd}}^{\text{h}} \begin{array}{c} \xrightarrow{|\text{ev}(\text{pt})|} \\ \xleftarrow{\simeq} \end{array} \text{Top}$$

- These adjunctions combine to give the following adjunction:

$$(6.1) \quad \mathbf{h} : \text{Shv}_{\infty\text{Gpd}} \rightleftarrows \text{Top} : \text{Sing}_*.$$

Because \mathbf{h} is a left adjoint, it automatically commutes with homotopy colimits. The functor \mathbf{h} also preserves finite products.

- Let M be a manifold and $\underline{M} \in \text{Shv}_{\text{Gpd}}$ the stack it represents. Then,

$$(6.2) \quad \mathfrak{h}(\mathbf{N}(\underline{M})) \simeq M \in \text{Top}.$$

- If $\mathcal{A} \in \text{Shv}_{\text{Ab}} \hookrightarrow \text{Shv}_{\text{Gpd}}$ is a sheaf of abelian groups that are C^∞ -modules, then

$$(6.3) \quad \mathfrak{h}(\mathbf{N}(\mathcal{A})) \simeq \text{pt} \in \text{Top}.$$

We now make the following definition/abbreviation.

Definition 6.4. Let A be a fixed abelian group. For a stack $\mathcal{M} \in \text{Shv}_{\text{Gpd}}$,

$$H^n(\mathcal{M}; A) := \text{hoShv}_{\infty\text{Gpd}}(\mathbf{N}(\mathcal{M}), \mathcal{K}(A, n)).$$

Example 6.5. For $M \in \text{Man}$, consider $\mathbf{N}(\underline{M}) \in \text{Shv}_{\infty\text{Gpd}}$, which is defined by considering the set $C^\infty(X, M)$ as a discrete simplicial set. The Yoneda lemma shows that Definition 6.4 agrees with our ordinary definition of cohomology,

$$\begin{aligned} H^n(\underline{M}, A) &= \text{hoShv}_{\infty\text{Gpd}}(\mathbf{N}(\underline{M}), \text{Sing}_* K(A, n)) \\ &\cong \text{hoTop}(M, K(A, n)) \cong H^n(M; A). \end{aligned}$$

The following proposition and proof are taken almost directly from Lemma 5.2 in [BunNV].

Proposition 6.6. *The natural map $EG \times_G M \rightarrow \mathcal{E}_\nabla G \times_G M$ induces equivalences*

$$EG \times_G M \xrightarrow{\simeq} \mathfrak{h}(\mathbf{N}(\mathcal{E}_\nabla G \times_G M)) \xrightarrow{\simeq} \mathfrak{h}(\mathbf{N}(EG \times_G M)).$$

Proof. We first show $\mathfrak{h}(\mathbf{N}(\mathcal{E}_\nabla G \times_G M)) \xrightarrow{\simeq} \mathfrak{h}(\mathbf{N}(EG \times_G M))$. Consider the following commutative diagram of prestacks

$$\begin{array}{ccc} \mathcal{E}_\nabla G \times_G M & \longrightarrow & EG \times_G M \\ \uparrow & & \uparrow \\ (\Omega^1 \otimes \mathfrak{g} \times M) // G & \longrightarrow & M // G, \end{array}$$

where we use the notation $\bullet // G$ to denote the action groupoid $[\bullet \rightrightarrows G \times \bullet]$ associated to a G -action. The group G acts on $\Omega^1 \otimes \mathfrak{g}$ via the usual change of connection formula, $\omega \cdot g = \text{Ad}_{g^{-1}} \omega + g^* \theta_{\mathfrak{g}}$. The right vertical map is given by associating to $X \rightarrow (M // G)$ the trivial bundle $X \leftarrow X \times G$, together with the map $X \times G \rightarrow M$ determined by the map $X \rightarrow M$. Likewise, the left vertical map is given by the same construction, but with the connection on $X \times G$ determined by the map $X \rightarrow \Omega^1 \otimes \mathfrak{g}$.

Since any bundle is locally trivializable, the two vertical maps induce equivalences of groupoids when evaluated on stalks (they are the stackification maps). Hence, upon taking the nerve and sheafifying, the vertical

maps induce equivalences in $\mathbf{Shv}_{\infty\mathbf{Gpd}}$ (see [BunNV, Lemma 5.2] and [FreH, (7.23)]).

$$\begin{array}{ccc} \mathbf{N}(\mathcal{E}_{\nabla}G \times_G M) & \longrightarrow & \mathbf{N}(\mathcal{E}G \times_G M) \\ \simeq \uparrow & & \simeq \uparrow \\ L(\mathbf{N}((\Omega^1 \otimes \mathfrak{g} \times M) // G)) & \longrightarrow & L(\mathbf{N}(M // G)) \end{array}$$

We now use the general fact that for a simplicial set B , the induced simplicial object in simplicial sets, given by the composition $\Delta^{\text{op}} \xrightarrow{B} \mathbf{Set} \xrightarrow{\mathbf{N}} \mathbf{sSet}$, gives a natural equivalence

$$\text{hocolim}_{\Delta^{\text{op}}} \mathbf{N}(B) \simeq B.$$

Rewriting the bottom entries as colimits, and applying the properties of \mathbf{h} described above, gives the following sequence of equivalences.

$$\begin{aligned} & \mathbf{h}(L(\mathbf{N}((\Omega^1 \otimes \mathfrak{g} \times M) // G))) \\ & \simeq \mathbf{h}(L(\Omega^1 \otimes \mathfrak{g} \times \underline{M} \rightrightarrows \underline{G} \times \Omega^1 \otimes \mathfrak{g} \times \underline{M} \xleftarrow{\cdot} \dots)) \\ & \simeq \mathbf{h}\left(L\left(\text{hocolim}_{\Delta^{\text{op}}} (\mathbf{N}(\Omega^1 \otimes \mathfrak{g} \times \underline{M}) \rightrightarrows \mathbf{N}(\underline{G} \times \Omega^1 \otimes \mathfrak{g} \times \underline{M}) \xleftarrow{\cdot} \dots)\right)\right) \\ & \simeq \text{hocolim}_{\Delta^{\text{op}}} (\mathbf{h}(\mathbf{N}(\Omega^1 \otimes \mathfrak{g})) \times \mathbf{h}(\mathbf{N}(\underline{M})) \rightrightarrows \mathbf{h}(\mathbf{N}(\underline{G})) \times \mathbf{h}(\mathbf{N}(\Omega^1 \otimes \mathfrak{g})) \times \mathbf{h}(\mathbf{N}(\underline{M})) \dots) \\ & \xrightarrow{\simeq} \text{hocolim}_{\Delta^{\text{op}}} (M \rightrightarrows G \times M \xleftarrow{\cdot} G \times G \times M \dots) \simeq \mathbf{h}(L(\mathbf{N}(M // G))). \end{aligned}$$

The equivalence $\mathbf{h}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M)) \xrightarrow{\simeq} \mathbf{h}(\mathbf{N}(\mathcal{E}G \times_G M))$ follows immediately. In the above argument, the third equivalence was given by fact that \mathbf{h} and L commute with colimits, $\mathbf{N}(\underline{G}^{\times k} \times \Omega^1 \otimes \mathfrak{g} \times \underline{M})$ is a sheaf, and \mathbf{h} preserves finite products. The fourth equivalence was given by the fact that $\mathbf{h}(\mathbf{N}(\underline{X})) \simeq X$, and $\mathbf{h}(\mathbf{N}(\Omega^1 \otimes \mathfrak{g})) \simeq \text{pt}$ since $\Omega^1 \otimes \mathfrak{g}$ is a sheaf of C^∞ -modules. Note that the middle entries were given as the homotopy colimit of a simplicial space (simplicial object in \mathbf{Top}). For proper simplicial spaces, the homotopy colimit is equivalent to the geometric realization; this is a common construction of BG when $M = \text{pt}$.

Now consider also the following commutative diagram of prestacks

$$\begin{array}{ccc} EG \times_G M & \longrightarrow & \mathcal{E}G \times_G M \\ \uparrow & & \uparrow \\ (EG \times M) // G & \longleftarrow & M // G, \end{array}$$

where the bottom left map is given by the homotopy equivalence $M \xrightarrow{\simeq} EG \times M$. The left vertical map is given by quotienting, and since G acts freely, it leads to an equivalence of groupoids when evaluated on any manifold X .

Hence, both vertical maps give equivalences in $\mathbf{Shv}_{\infty\mathbf{Gpd}}$.

$$\begin{array}{ccc} \mathbf{N}(EG \times_G M) & \longrightarrow & \mathbf{N}(\mathcal{E}G \times_G M) \\ \simeq \uparrow & & \simeq \uparrow \\ L(\mathbf{N}[EG \times M \rightrightarrows G \times (EG \times M)]) & \longleftarrow & L(\mathbf{N}[M \rightrightarrows G \times M]) \end{array}$$

By the same calculation as above

$$\begin{aligned} \mathbf{h}(L(\mathbf{N}(M // G))) &\simeq \operatorname{hocolim}_{\Delta^{\text{op}}} (M \rightrightarrows G \times M \overset{\leftarrow}{\rightrightarrows} G \times G \times M \cdots), \\ \mathbf{h}(L(\mathbf{N}((EG \times M) // G))) &\simeq \operatorname{hocolim}_{\Delta^{\text{op}}} (EG \times M \rightrightarrows G \times EG \times M \overset{\leftarrow}{\rightrightarrows} \cdots), \end{aligned}$$

and the homotopy equivalence $\text{pt} \xrightarrow{\simeq} EG$ induces a homotopy equivalence of the relevant simplicial spaces. This implies $\mathbf{h}(L(\mathbf{N}(M // G))) \xrightarrow{\simeq} \mathbf{h}(L(\mathbf{N}((EG \times M) // G)))$, which in turn gives the equivalence

$$EG \times_G M \simeq \mathbf{h}(L(\mathbf{N}(EG \times_G M))) \xrightarrow{\simeq} \mathbf{h}(\mathbf{N}(\mathcal{E}G \times_G M)). \quad \square$$

Proposition 6.7. *The map $EG \times_G M \rightarrow \mathcal{E}_{\nabla}G \times_G M$, defined by the connection on EG , induces an isomorphism in cohomology*

$$H^*(\mathcal{E}_{\nabla}G \times_G M; A) \xrightarrow{\cong} H^*(EG \times_G M; A).$$

Proof. This follows almost immediately from Definition 6.4, the adjunction (6.1), and the calculation of $\mathbf{h}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M))$ in Proposition 6.6. Together, these give the following sequence of isomorphisms:

$$\begin{aligned} H^n(\mathcal{E}_{\nabla}G \times_G M; A) &= \operatorname{hoShv}_{\infty\mathbf{Gpd}}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M), \mathcal{K}(A, n)) \\ &\cong \operatorname{hoTop}(\mathbf{h}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M)), K(A, n)) \\ &\xrightarrow{\cong} \operatorname{hoTop}(EG \times_G M, K(A, n)) \\ &\cong H^n(EG \times_G M; A). \end{aligned} \quad \square$$

Finally, we repeat an explicit construction of a simplicial sheaf representing differential cohomology. This construction, and minor variations, have already appeared in several places, including [HS] implicitly and [BunNV, Bun, FSS, HQ, Sch] explicitly. In order to obtain simplicial sets from cochain complexes, we use the standard trick of reversing the grading and shifting, and then using the Dold–Kan construction. The inclusions of

$$\mathbb{Z} \hookrightarrow \mathbb{R} \hookrightarrow \Omega^0$$

induce the following maps of presheaves of chain complexes, where degree 0 is written on the left.

$$(\Omega_{\text{cl}}^n \leftarrow 0 \leftarrow \cdots) \hookrightarrow (\Omega_{\text{cl}}^n \xleftarrow{d} \Omega^{n-1} \xleftarrow{d} \cdots \xleftarrow{d} \Omega^0) \hookrightarrow (0 \leftarrow \cdots \leftarrow \mathbb{Z} \leftarrow \cdots)$$

Applying Dold–Kan (and sheafifying where necessary), we then take the homotopy pullback in $\mathbf{Shv}_{\infty\mathbf{Gpd}}$, obtaining

$$(6.8) \quad \begin{array}{ccc} \check{\mathcal{K}}(\mathbb{Z}, n) & \longrightarrow & L(\Gamma(0 \leftarrow \cdots \leftarrow \mathbb{Z} \leftarrow \cdots)) \\ \downarrow & & \downarrow \\ \Gamma(\Omega_{\text{cl}}^n \leftarrow 0 \leftarrow \cdots) & \longrightarrow & \Gamma(\Omega_{\text{cl}}^n \xleftarrow{d} \Omega^{n-1} \xleftarrow{d} \cdots \xleftarrow{d} \Omega^0) \end{array}$$

The top right corner is equivalent to $\mathcal{K}(\mathbb{Z}, n)$, and the bottom right corner is equivalent to $\mathcal{K}(\mathbb{R}, n)$, so the construction of $\check{\mathcal{K}}(\mathbb{Z}, n)$ fits the description from (3.10).

We now have two potential definitions of $\check{H}_G^*(M)$. The first was given in Section 2 using the cochain complex $\check{C}_G(n)^*(M; \mathbb{Z})$, and the second is given in this section as

$$\text{hoShv}_{\infty\mathbf{Gpd}}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M), \check{\mathcal{K}}(\mathbb{Z}, n)).$$

We now prove these two possible definitions of $\check{H}_G^*(M)$ agree.

Proof of Theorem 3.11. We use Dold–Kan to consider $\check{C}_G(n)^*(M)$ as a homotopy pullback of simplicial abelian groups via the following.

$$(6.9) \quad \begin{array}{ccc} \Gamma(\check{Z}_G(n)^n(M) \xleftarrow{d} \check{C}_G(n)^{n-1}(M) \xleftarrow{d}) & \longrightarrow & \Gamma(Z^n(EG \times_G M; \mathbb{Z}) \xleftarrow{\delta} \cdots) \\ \downarrow & & \downarrow \\ \Gamma(\Omega_G^n(M)_{\text{cl}} \leftarrow 0) & \longrightarrow & \Gamma(Z^n(EG \times_G M; \mathbb{R}) \xleftarrow{\delta} \cdots) \end{array}$$

The Freed–Hopkins isomorphism [FreH, Propositions 6.22 and Theorem 7.28] gives us

$$\text{hoShv}_{\infty\mathbf{Gpd}}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M), \mathbf{N}(\Omega_{\text{cl}}^n)) \cong \text{Shv}_{\mathbf{Gpd}}(\mathcal{E}_{\nabla}G \times_G M, \Omega_{\text{cl}}^n) \cong \Omega_G^n(M)_{\text{cl}}.$$

Likewise Proposition 6.6 shows us that

$$\begin{aligned} & \text{Shv}_{\infty\mathbf{Gpd}}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M), L(\Gamma(0 \leftarrow \cdots \leftarrow \mathbb{Z}))) \\ & \simeq \text{Shv}_{\infty\mathbf{Gpd}}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M), \mathcal{K}(\mathbb{Z}, n)) \\ & \simeq \text{Shv}_{\infty\mathbf{Gpd}}^h(\mathbf{h}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M)), \mathcal{K}(\mathbb{Z}, n)) \simeq \mathcal{S}(K(\mathbb{Z}, n)^{EG \times_G M}), \\ & \text{Shv}_{\infty\mathbf{Gpd}}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M), \Gamma(\Omega_{\text{cl}}^n \xleftarrow{d} \Omega^{n-1} \xleftarrow{d} \cdots)) \\ & \simeq \text{Shv}_{\infty\mathbf{Gpd}}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M), \mathcal{K}(\mathbb{R}, n)) \\ & \simeq \text{Shv}_{\infty\mathbf{Gpd}}^h(\mathbf{h}(\mathbf{N}(\mathcal{E}_{\nabla}G \times_G M)), \mathcal{K}(\mathbb{R}, n)) \simeq \mathcal{S}(K(\mathbb{R}, n)^{EG \times_G M}). \end{aligned}$$

There is a natural homotopy equivalence

$$\mathcal{S}(K(A, n)^X) \xrightarrow{\simeq} \Gamma(Z^n(X; A) \xleftarrow{\delta} C^{n-1}(X; A) \xleftarrow{\delta} \cdots)$$

induced by pulling back a fundamental cocycle in $Z^n(K(A, n); A)$ and integrating over the simplex [HS, Proposition A.12 and Corollary D.13]. Thus,

we have equivalences from three corners of (6.8) evaluated on $\mathbf{N}(\mathcal{E}_{\check{\vee}}G \times_G M)$, and the corresponding three corners of (6.9). Therefore, the induced map between homotopy pullbacks

$$\mathrm{Shv}_{\infty\mathrm{Gpd}}(\mathbf{N}(\mathcal{E}_{\check{\vee}}G \times_G M), \check{\mathcal{K}}(\mathbb{Z}, n)) \xrightarrow{\sim} \Gamma(\check{Z}_G(n)^n \xleftarrow{d} \check{C}_G(n)^{n-1} \xleftarrow{d} \dots)$$

is also an equivalence in $\infty\mathrm{Gpd}$, which gives us the desired isomorphism in the homotopy category. \square

Appendix A. Equivariant de Rham theory

We briefly recall some standard facts about equivariant cohomology, principal bundles with connection, and equivariant differential forms. Our goal is to describe the Weil algebra $W(\mathfrak{g})$ from the perspective of differential forms on principal bundles. We hope this makes clear why having connections is so essential to using $\Omega_G(M)$. Among the many wonderful references, our treatment is most heavily influenced by: the introduction to principal bundles in Section 1 of [Fre], the summary of equivariant cohomology [Mei], and the standard textbook for G^* -algebras and equivariant de Rham cohomology [GS].

A.1. Equivariant cohomology. Let G be a reasonably nice topological group. Then there exists a contractible space EG on which G acts freely from the right, giving us a universal G -bundle $EG \rightarrow BG$. If M has a continuous left G -action, the *homotopy quotient* (or *Borel construction*) of M is defined

$$EG \times_G M := (EG \times M) / ((eg, x) \sim (e, gx)).$$

Borel's construction of the *equivariant cohomology* of M , with coefficients in an abelian group A , is simply the ordinary cohomology of the homotopy quotient

$$H_G^*(M; A) := H^*(EG \times_G M; A).$$

A G -equivariant map $M \xrightarrow{f} N$ induces a map on the Borel constructions, giving the desired homomorphisms

$$H_G^*(N; A) \xrightarrow{f_G^*} H_G^*(M; A).$$

There are important, though subtle, relationships between equivariant and ordinary cohomology. In particular, $H_G^*(\mathrm{pt}; A) = H^*(BG; A)$, and the natural map $M \rightarrow \mathrm{pt}$ makes $H_G^*(M; R)$ into a $H^*(BG; R)$ -module when R is a ring. When G acts freely on M , there is a natural isomorphism $H_G^*(M; A) \cong H^*(M/G; A)$. This follows from the fact that the projection $EG \times_G M \rightarrow M/G$ is a locally trivial fiber bundle with fiber EG , and hence it is a homotopy equivalence.

A.2. Principal G -bundles with connection. Let G be a Lie group, and let \mathfrak{g} be the Lie algebra of left-invariant vector fields on G .

A *principal G -bundle* on a manifold X is a manifold P equipped with a free right G -action and map to X such that

$$P \xrightarrow{\pi} X \cong P/G.$$

Note that on differential forms, $\pi^*: \Omega(X) \rightarrow \Omega(P)$ is an injective map, and the subspace $\pi^*\Omega(X)$ is naturally identified as the *basic* (i.e., invariant horizontal) forms

$$(A.1) \quad \Omega(X) \cong \pi^*\Omega(X) = \Omega(X)_{\text{hor}}^G \subset \Omega(P).$$

Here, the horizontal forms are defined by: $\omega \in \Omega(P)_{\text{hor}}$ if $\iota_X\omega = 0$ for all $X \in \mathfrak{g}$.

A *connection* on P is an Ad-equivariant \mathfrak{g} -valued 1-form $\Theta \in \Omega^1(P; \mathfrak{g})$ that restricts fiberwise to the Maurer–Cartan form $\theta_{\mathfrak{g}}$. More explicitly:

- $\iota_X\Theta = X \in \Omega^0(P; \mathfrak{g})$ for any vector field $X \in \mathfrak{g}$.
- $g^*(\Theta) = \text{Ad}_{g^{-1}}\Theta$, where $g: P \rightarrow P$ is right multiplication by g .

The Ad-equivariance can be rewritten as $\Theta \in (\Omega^1(P) \otimes \mathfrak{g})^G$, where the second factor \mathfrak{g} is the adjoint representation of G . The *curvature* is defined

$$\Omega := d\Theta + \frac{1}{2}[\Theta \wedge \Theta] \in \Omega^2(P; \mathfrak{g}).$$

There is also an equivalent, but more geometric, interpretation of a connection. The G -action defines the distribution T^VP of vertical tangent vectors in P by

$$\mathfrak{g} \xrightarrow{\cong} \text{Ker } \pi_* =: T^VP \subset TP.$$

A connection on P is the choice of an equivariant horizontal distribution T^HP ; i.e., a connection is equivalent to an equivariant splitting of the tangent bundle

$$TP = T^HP \oplus T^VP \cong \pi^*TM \oplus \mathfrak{g}.$$

The connection 1-form Θ gives a projection from TP onto $T^VP \cong \mathfrak{g}$, thus defining the horizontal distribution by

$$T^HP := \text{Ker } \Theta \subset TP.$$

The splitting of TP into horizontal and vertical subspaces induces a bi-grading on the differential forms

$$\Omega^{i,j}(P) := C^\infty(P, \Lambda^i T^HP^* \otimes \Lambda^j T^VP^*) \cong C^\infty(P, \pi^* \Lambda^i TX^* \otimes \Lambda^j \mathfrak{g}^*).$$

In this bi-grading, the exterior derivative decomposes as

$$d = d^{0,1} + d^{1,0} + d^{2,-1} \cong (-1)^i d_{\mathfrak{g}} + d_{\nabla} + (-1)^i \iota_{\Omega}.$$

Here, $d_{\mathfrak{g}}: \Lambda^j \mathfrak{g}^* \otimes C^\infty(P) \rightarrow \Lambda^{j+1} \mathfrak{g}^* \otimes C^\infty(P)$ is the Lie algebra (or Chevalley–Eilenberg) differential for the G -module $C^\infty(P)$ [ChevE]. If we restrict to $\Omega(P)^G$, then $(\Lambda \mathfrak{g}^*, d_{\mathfrak{g}})$ is naturally the de Rham complex of left-invariant forms on G . The connection Θ induces the covariant derivative d_{∇} , and ι_{Θ}

is the derivation induced by contracting along the vector-valued 2-form Ω . See Section 3 of [Red] for more details.

A.3. G^* -algebras. Let (\mathcal{A}, d) be a commutative differential graded algebra (DGA), where d is a derivation of degree +1 and commutative means $ab = (-1)^{|a||b|}ba$ for homogeneous elements. A DGA-automorphism of degree 0 is an algebra automorphism $\phi: \mathcal{A} \rightarrow \mathcal{A}$ that preserves grading and commutes with d .

Definition A.2 (Section 2.3 of [GS]). A G^* -algebra is a commutative DGA (\mathcal{A}, d) equipped with representations

$$G \xrightarrow{\rho} \text{Aut}_{\text{DGA}}(\mathcal{A}) \quad \text{and} \quad \mathfrak{g} \xrightarrow{\iota} \text{Der}(\mathcal{A})$$

of degree 0 and -1, respectively, such that ι is G -equivariant with respect to ρ and satisfies the Cartan equation; i.e., for all $X \in \mathfrak{g}$

$$(A.3) \quad \rho_g \iota_X \rho_{g^{-1}} = \iota_{\text{Ad}_g X},$$

$$(A.4) \quad \iota_X d + d \iota_X = L_X.$$

Here, $L: \mathfrak{g} \rightarrow \text{Der}(\mathcal{A})$ is the Lie algebra representation induced by ρ .

A map $\phi: \mathcal{A}_1 \rightarrow \mathcal{A}_2$ is a morphism of G^* -algebras if ϕ commutes with multiplication, ρ , d , and ι .

Example A.5. Suppose that a manifold M has a left G -action. Then the de Rham complex $(\Omega(M), d)$ is naturally a G^* -algebra. The G -action on $\Omega(M)$ is defined by

$$\rho_g \omega := (g^{-1})^* \omega,$$

and ι is defined by composing the usual interior derivative with the action of \mathfrak{g} on vector fields

$$\mathfrak{g} \rightarrow \mathfrak{X}(M) \xrightarrow{\iota} \text{Der}(\Omega(M)).$$

If $f: M_1 \rightarrow M_2$ is G -equivariant, then $f^*: \Omega(M_2) \rightarrow \Omega(M_1)$ is a morphism of G^* -algebras.

Remark A.6. We use the convention that manifolds M have a left G -action, while principal bundles P have a right G -action. However, we sometimes implicitly use the natural switch between left and right actions. Given a left G -action on a set Y , define the right G -action via the formula $y \cdot g := g^{-1} \cdot y$; similarly, a right G -action induces a left G -action.

Definition A.7. Let \mathcal{A} a G^* -algebra. An element a is *invariant* if $\rho_g a = a$ for all $g \in G$, and it is *horizontal* if $\iota_X a = 0$ for all $X \in \mathfrak{g}$. The *basic* sub-algebra is the intersection of the invariant and horizontal elements:

$$\mathcal{A}_{\text{basic}} := \mathcal{A}^G \cap \mathcal{A}_{\text{hor}} = \mathcal{A}_{\text{hor}}^G.$$

The definition of a G^* -algebra implies that $(\mathcal{A}_{\text{basic}}, d)$ is a sub-DGA of (\mathcal{A}, d) . Furthermore, if $\phi: \mathcal{A} \rightarrow \mathcal{B}$ is a morphism of G^* -algebras, then ϕ restricts to a DGA-morphism on the basic subcomplexes $\phi: \mathcal{A}_{\text{bas}} \rightarrow \mathcal{B}_{\text{bas}}$.

Definition A.8. A connection on a G^* -algebra \mathcal{A} is an element

$$\Theta \in (\mathcal{A}^1 \otimes \mathfrak{g})^G$$

such that $\iota_X \Theta = X$ for all $X \in \mathfrak{g}$. The curvature of Θ is defined

$$\Omega := d\Theta + \frac{1}{2}[\Theta, \Theta] \in (\mathcal{A}^2 \otimes \mathfrak{g})^G.$$

Example A.9. Suppose $P \xrightarrow{\pi} X$ is a principal G -bundle. Example A.5 and Remark A.6 show that $\Omega(P)$ is naturally a G^* -algebra, though we write $\rho_g \omega = g^* \omega$ due to the fact that G acts on the right. As noted in (A.1), the basic subcomplex $\Omega(P)_{\text{basic}}$ is naturally isomorphic to $\Omega(X)$. Furthermore, connections for the principal bundle P are equivalent to connections for the G^* -algebra $\Omega(P)$.

Example A.10. The *Weil algebra* $W(\mathfrak{g})$ is a G^* -algebra with connection, and it is constructed so that it canonically maps to any other G^* -algebra with connection. Explicitly,

$$\begin{aligned} W(\mathfrak{g}) &:= S\mathfrak{g}^* \otimes \Lambda\mathfrak{g}^*, \\ \deg S^1\mathfrak{g}^* &= 2, \quad \deg \Lambda^1\mathfrak{g}^* = 1, \\ d_W &:= d_{\mathfrak{g}} + d_K. \end{aligned}$$

Here S and Λ are the total symmetric and exterior powers, so the coadjoint representation \mathfrak{g}^* naturally makes $W(\mathfrak{g})$ into a G -representation. The differential $d_{\mathfrak{g}}$ is the Chevalley–Eilenberg differential for Lie algebra cohomology with values in the \mathfrak{g} -module $S\mathfrak{g}^*$

$$S^i\mathfrak{g}^* \otimes \Lambda^j\mathfrak{g}^* \xrightarrow{d_{\mathfrak{g}}} S^i\mathfrak{g}^* \otimes \Lambda^{j+1}\mathfrak{g}^*;$$

it has degree $(0, 1)$ under the bi-grading $W^{2i,j}(\mathfrak{g}) = S^i\mathfrak{g}^* \otimes \Lambda^j\mathfrak{g}^*$. The Koszul differential d_K has degree $(2, -1)$ and is defined by extending the natural isomorphism

$$\Lambda^1\mathfrak{g}^* \xrightarrow[\cong]{d_K} S^1(\mathfrak{g}^*)$$

to a derivation

$$S^i(\mathfrak{g}^*) \otimes \Lambda^j(\mathfrak{g}^*) \xrightarrow{d_K} S^{i+1}(\mathfrak{g}^*) \otimes \Lambda^{j-1}(\mathfrak{g}^*).$$

The derivation ι has degree $(0, -1)$ and is induced by the usual contraction

$$\mathfrak{g} \xrightarrow{\iota} \text{End}(\Lambda\mathfrak{g}^*).$$

The Weil algebra $W(\mathfrak{g})$ has the same cohomology as $EG \simeq \text{pt}$; i.e., $H^0(W(\mathfrak{g}), d_W) = \mathbb{R}$ and $H^i(W(\mathfrak{g}), d_W) = 0$ for $i > 0$. It has a natural connection

$$\theta_{\mathfrak{g}} \in \Lambda^1\mathfrak{g}^* \otimes \mathfrak{g} = W^{0,1}(\mathfrak{g}) \otimes \mathfrak{g}$$


given by the identity map $\mathfrak{g} \rightarrow \mathfrak{g}$; i.e., $\theta_{\mathfrak{g}}(X) = X$ for $X \in \mathfrak{g}$. When there is no risk of confusion, we will drop the subscript and write θ for $\theta_{\mathfrak{g}}$. Using the fact that $d_{\mathfrak{g}}\theta = -\frac{1}{2}[\theta \wedge \theta]$, we see that the curvature $\Omega = \Omega_{\mathfrak{g}}$ equals $d_K\theta_{\mathfrak{g}}$. In light of this, we rewrite the Koszul derivative d_K as $\iota_{\Omega_{\mathfrak{g}}}$.

Proposition A.11 (Weil homomorphism). *Let \mathcal{A} be a G^* -algebra with connection Θ . Then, there is natural morphism of G^* -algebras with connection $W(\mathfrak{g}) \xrightarrow{\Theta^*} \mathcal{A}$ induced by $\theta \mapsto \Theta$, $\Omega \mapsto \Omega$.*

Therefore, we see that $W(\mathfrak{g})$ serves as a natural algebraic model for differential forms on EG . It is acyclic, and to any G -bundle with connection $(P, \Theta) \rightarrow X$, the Weil homomorphism is a natural map

$$(A.12) \quad \begin{aligned} W^{2i,j}(\mathfrak{g}) &\xrightarrow{\Theta^*} \Omega^{2i,j}(P) \\ \omega \otimes \eta &\longmapsto \omega(\Omega^{\wedge i}) \wedge \eta(\Theta^{\wedge j}) \end{aligned}$$

that is compatible with the bi-grading, multiplication, G -action, derivative d , and contraction ι . Below is a diagram showing this. For generic $\alpha \in W(\mathfrak{g})$, we use the notation $\alpha(\Theta)$ for $\Theta^*(\alpha)$.



The universal Chern–Weil and Chern–Simons forms are naturally described via the Weil model. The usual Chern–Weil forms are given by simply restricting (A.12) to the basic subcomplex

$$\begin{aligned} (S^k \mathfrak{g}^*)^G &= W^{2k}(\mathfrak{g})_{\text{basic}} \xrightarrow{\Theta^*} \Omega^{2k}(P)_{\text{basic}} \cong \Omega^{2k}(M) \\ \omega &\longmapsto \omega(\Theta^{\wedge n}). \end{aligned}$$

That $W(\mathfrak{g})$ is acyclic implies any such $\omega \in (S\mathfrak{g}^*)^G$ is exact in $W(\mathfrak{g})$, and there is a standard way to pick out such a coboundary. Using the trivial G^* -algebra $\Omega([0, 1])$ with natural coordinate t , define

$$\begin{aligned} \theta_t &:= t\theta && \in (\Omega([0, 1]) \otimes W(\mathfrak{g}))^1 \otimes \mathfrak{g}, \\ \Omega_t &:= d(\theta_t) + \frac{1}{2}[\theta_t \wedge \theta_t] && \in (\Omega([0, 1]) \otimes W(\mathfrak{g}))^2 \otimes \mathfrak{g}, \end{aligned}$$

where one can rewrite the curvature to give $\Omega_t = dt\theta + t\Omega + \frac{1}{2}(t^2 - t)[\theta \wedge \theta]$.

For $\omega \in (S^k \mathfrak{g}^*)^K$, then $\omega(\Omega_t^{\wedge k}) \in (\Omega([0, 1]) \otimes W(\mathfrak{g}))_{\text{basic}}$, and the Chern–Simons form is defined using integration over the interval $[0, 1]$ by

$$(A.13) \quad \text{CS}_\omega := \int_{[0,1]} \omega(\Omega_t^{\wedge k}) \in W^{2k-1}(\mathfrak{g})^G.$$

Stokes Theorem implies that $d\text{CS}_\omega = \omega \in \mathbb{W}(\mathfrak{g})$, and the Weil homomorphism sends this to the standard Chern–Simons forms on the total space of principal bundles

$$\begin{aligned} \mathbb{W}(\mathfrak{g})^G &\xrightarrow{\Theta^*} \Omega(P)^G \\ \text{CS}_\omega &\longmapsto \text{CS}_\omega(\Theta). \end{aligned}$$

A.4. Equivariant Forms. We now describe an explicit de Rham model $\Omega_G(M)$, often referred to as the Weil model, for $H_G^*(M; \mathbb{R})$. While the definition of $\Omega_G(M)$ does not require G to be compact, the cohomology of $\Omega_G(M)$ is not necessarily isomorphic to $H_G^*(M; \mathbb{R})$ for noncompact G .

Suppose M is a manifold equipped with a smooth left G -action. The tensor product $\mathbb{W}(\mathfrak{g}) \otimes \Omega(M)$ is a G^* -algebra with derivative

$$d_G = d_W \otimes 1 + 1 \otimes d.$$

Definition A.14. The complex of *equivariant differential forms* is the basic sub-complex

$$(\Omega_G^*(M), d_G) := ((\mathbb{W}(\mathfrak{g}) \otimes \Omega(M))_{\text{hor}}^G, d_W \otimes 1 + 1 \otimes d).$$

Note that if $f: M \rightarrow N$ is G -equivariant map, then

$$\mathbb{W}(\mathfrak{g}) \otimes \Omega(N) \xrightarrow{1 \otimes f^*} \mathbb{W}(\mathfrak{g}) \otimes \Omega(M)$$

restricts to the basic complex, giving a naturally induced DGA-morphism we denote

$$\Omega_G(N) \xrightarrow{f_G^*} \Omega_G(M).$$

To relate $\Omega_G(M)$ to the cohomology of $EG \times_G M$, we use the following geometric fact. There exist finite-dimensional smooth n -classifying bundles with connection $(E^{(n)}G, \Theta_{E^{(n)}G}) \rightarrow B^{(n)}G$ ([NR]). This means that if $\dim(X) \leq n$, any bundle with connection $(P, \Theta) \rightarrow X$ is isomorphic to the pullback $f^*(E^{(n)}G, \Theta_{E^{(n)}G})$ for some smooth map $X \xrightarrow{f} B^{(n)}G$. Furthermore, any two maps classifying (P, Θ) are homotopic. The universal bundle $EG \rightarrow BG$ can then be constructed as a direct limit of finite-dimensional manifolds.

$$\begin{array}{ccc} (EG, \Theta_{EG}) & & \\ \downarrow \pi & := & \varinjlim \left(\begin{array}{c} (E^{(n)}G, \Theta_{E^{(n)}G}) \\ \downarrow \pi \\ B^{(n)}G \end{array} \right) \\ BG & & \end{array}$$

When discussing differential forms and cochains, we use the notations

$$\begin{aligned} \Omega^*(EG) &= \varprojlim \Omega^*(E^{(n)}G), & C^*(EG; A) &= \varprojlim C^*(E^{(n)}G; A), \\ \Omega^*(BG) &= \varprojlim \Omega^*(B^{(n)}G), & C^*(BG; A) &= \varprojlim C^*(B^{(n)}G; A). \end{aligned}$$

The connection Θ_{EG} and the Weil homomorphism give a G^* -algebra homomorphism

$$W(\mathfrak{g}) \xrightarrow{\Theta_{EG}^*} \Omega(EG),$$

which in turn gives

(A.15)

$$\Omega_G(M) = (W(\mathfrak{g}) \otimes \Omega(M))_{\text{basic}} \xrightarrow{\Theta_{EG}^* \otimes 1} \Omega(EG \times M)_{\text{basic}} \cong \Omega(EG \times_G M).$$

Theorem A.16 (Equivariant de Rham Theorem). *For G compact, (A.15) induces an isomorphism in cohomology*

$$H^*(\Omega_G(M), d_G) \xrightarrow{\cong} H^*(\Omega(EG \times_G M), d) \cong H_G^*(M; \mathbb{R}).$$

A proof of the above theorem can be found in Theorems 2.5.1 (and Theorems 4.3.1 and 6.7.1) of [GS]. The following lemma is used in the first construction of $\check{H}_G^*(M)$ in Section 2, along with the proof of Proposition 4.18. Though it is certainly well-known, we are unaware of a specific reference, and we prove it directly so that the first construction of $\check{H}_G^*(M)$ does not rely on results from [FreH].

Lemma A.17. *The homomorphism $W(\mathfrak{g}) \xrightarrow{\Theta_{EG}^*} \Omega(EG)$ is injective, as are the induced homomorphisms $\Omega_G(M) \rightarrow \Omega(EG \times_G M)$.*

Proof. The second homomorphism is given by restricting

$$W(\mathfrak{g}) \otimes \Omega(M) \longrightarrow \Omega(EG) \otimes \Omega(M) \cong \Omega(EG \times M)$$

to the basic subcomplex. Showing this map is injective is equivalent to showing $W(\mathfrak{g}) \rightarrow \Omega(EG)$ is injective.

Any bundle with connection is isomorphic to the pullback of (EG, Θ_{EG}) , so it suffices to show that for any element $\alpha \in W(\mathfrak{g})$, there exists some $(P, \Theta) \xrightarrow{\pi} X$ such that $\alpha(\Theta) \neq 0 \in \Omega(P)$. At any point p , $\Lambda T_p^* P$ is an algebra freely generated by the horizontal and vertical cotangent spaces. On the vertical part, $W^{0,*}(\mathfrak{g}) \rightarrow \Lambda T_p^* P^* \cong \Lambda \mathfrak{g}^*$ is an isomorphism, so it suffices to show the Weil homomorphism is injective on $W^{*,0}(\mathfrak{g})$.

Let $\omega \in S^n \mathfrak{g}^*$ be any nonzero element. Then $\omega(\xi_{i_1} \cdots \xi_{i_n}) \neq 0$ for some (i_1, \dots, i_n) , where $\{\xi_i\}$ be a basis of \mathfrak{g} . Let $\mathbb{R}^{2n} \times G \rightarrow \mathbb{R}^{2n}$ be the trivial bundle. In a neighborhood of 0, use the canonical frame p to define a connection Θ by

$$p^* \Theta = x^1 dx^2 \xi_{i_1} + \cdots + x^{2n-1} dx^{2n} \xi_{i_n} \in \Omega^1(\mathbb{R}^{2n}; \mathfrak{g}).$$

The local curvature is given by

$$p^* \Omega = dx^1 dx^2 \xi_{i_1} + \cdots + dx^{2n-1} dx^{2n} \xi_{i_n} + \frac{1}{2} [p^* \Theta \wedge p^* \Theta],$$

and the terms involving $p^* \Theta$ vanish when all $x^i = 0$. Evaluating $p^* \omega(\Theta^{\wedge n})$ at the origin gives

$$p^* \omega(\Theta^{\wedge n})_{(0)}(\partial_1, \dots, \partial_{2n}) = n! \omega(\xi_{i_1} \cdots \xi_{i_n}) \neq 0. \quad \square$$

A.5. Cartan model and Mathai–Quillen. There is another G^* -algebra structure on $W(\mathfrak{g}) \otimes \Omega(M)$ that leads to the Cartan model for $H_G^*(M; \mathbb{R})$. It is obtained by modifying the G^* -structure so that the interior derivative ι only acts on the factor $W(\mathfrak{g})$.

The G -equivariant homomorphism

$$\Omega^1(M) \xrightarrow{\iota_\theta} \Lambda^1 \mathfrak{g}^*,$$

defined by $(\iota_\theta \psi)(X) = \iota_{\theta(X)} \psi = \iota_X \psi$, induces a derivation on associative algebra $W(\mathfrak{g}) \otimes \Omega(M)$. It exponentiates to a G -equivariant automorphism of associative algebras

$$W(\mathfrak{g}) \otimes \Omega(M) \xrightarrow{\exp(\iota_\theta)} W(\mathfrak{g}) \otimes \Omega(M)$$

known as the Mathai–Quillen isomorphism [MaQ]. Conjugating by e^{ι_θ} gives the new operators

$$d_C := (\exp \iota_\theta) d_G (\exp -\iota_\theta), \quad \iota^C := (\exp \iota_\theta) \iota (\exp -\iota_\theta)$$

where $\iota_X^C(\alpha \otimes \psi) = (\iota_X \alpha) \otimes \psi$. The isomorphism between G^* -algebras induces an isomorphism of the basic subcomplexes, one of which is the previously discussed Weil model. The other, known as the Cartan model, is

$$(W(\mathfrak{g}) \otimes \Omega(M))_{\text{Ker } \iota^C}^G = (S\mathfrak{g}^* \otimes \Omega(M))^G,$$

and the derivative takes the form $d_C = d - \iota_{\Omega_{\mathfrak{g}}}$, where $\iota_{\Omega_{\mathfrak{g}}}$ is the degree 1 derivation induced by

$$\Omega^k(M) \xrightarrow{\Omega_{\mathfrak{g}} \otimes 1} S^1 \mathfrak{g}^* \otimes \mathfrak{g} \otimes \Omega^k(M) \xrightarrow{1 \otimes \iota} S^1 \mathfrak{g}^* \otimes \Omega^{k-1}(M).$$

The Mathai–Quillen isomorphism may be used to interpret $\Omega_G(M)$ as the Cartan model throughout the entire paper.

References

- [BeV] BERLINE, NICOLE; VERGNE, MICHÈLE. Zéros d'un champ de vecteurs et classes caractéristiques équivariantes. *Duke Math. J.* **50** (1983), no. 2, 539–549. MR705039 (84i:58114), Zbl 0515.58007, doi:10.1215/S0012-7094-83-05024-X.
- [BoT] BOTT, RAOUL; TU, LORING W. Equivariant characteristic classes in the Cartan model. *Geometry, analysis and applications* (Varanasi, 2000), 3–20, *World Sci. Publ., River Edge, NJ*, 2001. MR1850463 (2002i:57046), Zbl 1016.53024, arXiv:math/0102001.
- [Bry] BRYLINSKI, JEAN-LUC. Loop spaces, characteristic classes and geometric quantization. *Progress in Mathematics*, 107. *Birkhäuser Boston Inc., Boston, MA*, 1993. xvi+300 pp. ISBN: 0-8176-3644-7. MR1197353 (94b:57030), Zbl 0823.55002, doi:10.1007/978-0-8176-4731-5.
- [Bun] BUNKE, ULRICH. Differential cohomology. Preprint, 2012. arXiv:1208.3961.
- [BunNV] BUNKE, ULRICH; NIKOLAUS, THOMAS; VÖLKL, MICHAEL. Differential cohomology theories as sheaves of spectra. *J. Homotopy Relat. Struct.* **11** (2016), no. 1, 1–66. MR3462099, Zbl 1341.57020, arXiv:1311.3188, doi:10.1007/s40062-014-0092-5.

- [CheeS] CHEEGER, JEFF; SIMONS, JAMES. Differential characters and geometric invariants. *Geometry and topology* (College Park, Md., 1983/84), 50–80, Lecture Notes in Math., 1167. Springer, Berlin, 1985. MR0827262 (87g:53059), Zbl 0621.57010, doi: 10.1007/BFb0075216.
- [ChevE] CHEVALLEY, CLAUDE; EILENBERG, SAMUEL. Cohomology theory of Lie groups and Lie algebras. *Trans. Amer. Math. Soc.* **63** (1948), 85–124. MR0024908 (9,567a), Zbl 0031.24803, doi: 10.2307/1990637.
- [FOS] FIGUEROA-O’FARRILL, JOSÉ M.; STANCIU, SONIA. Gauged Wess–Zumino terms and equivariant cohomology. *Phys. Lett. B* **341** (1994), no. 2, 153–159. MR1311654 (96h:81069), arXiv:hep-th/9407196, doi: 10.1016/0370-2693(94)90304-2.
- [FSS] FIORENZA, DOMENICO; SCHREIBER, URS; STASHEFF, JIM. Čech cocycles for differential characteristic classes: an ∞ -Lie theoretic construction. *Adv. Theor. Math. Phys.* **16** (2012), no. 1, 149–250. MR3019405, Zbl 06171205, arXiv:1011.4735, doi: 10.4310/ATMP.2012.v16.n1.a5.
- [Fre] FREED, DANIEL S. Classical Chern–Simons theory. I. *Adv. Math.* **113** (1995), no. 2, 237–303. MR1337109 (96h:58019), Zbl 0844.58039, doi: 10.1006/aima.1995.1039.
- [FreH] FREED, DANIEL S.; HOPKINS, MICHAEL J. Chern–Weil forms and abstract homotopy theory. *Bull. Amer. Math. Soc. (N.S.)* **50** (2013), no. 3, 431–468. MR3049871, Zbl 1281.58001, arXiv:1301.5959, doi: 10.1090/S0273-0979-2013-01415-0.
- [Gom] GOMI, KIYONORI. Equivariant smooth Deligne cohomology. *Osaka J. Math.* **42** (2005), no. 2, 309–337. MR2147734 (2006j:14026), Zbl 1081.14030, arXiv:math/0307373.
- [GS] GUILLEMIN, VICTOR W.; STERNBERG, SHLOMO. Supersymmetry and equivariant de Rham theory. Mathematics Past and Present. Springer-Verlag, Berlin, 1999. xxiv+228 pp. ISBN: 3-540-64797-X. MR1689252 (2001i:53140), Zbl 0934.55007, doi: 10.1007/978-3-662-03992-2.
- [HaL] HARVEY, REESE; LAWSON, BLAINE. From sparks to grundles—differential characters. *Comm. Anal. Geom.* **14** (2006), no. 1, 25–58. MR2230569 (2007d:53125), Zbl 1116.53048, arXiv:math/0306193, doi: 10.4310/CAG.2006.v14.n1.a2.
- [Hei] HEINLOTH, JOCHEN. Notes on differentiable stacks. *Mathematisches Institut, Georg-August-Universität Göttingen: Seminars Winter Term 2004/2005*, 1–32. Universitätsdrucke Göttingen, Göttingen, 2005. MR2206877 (2007c:19009), Zbl 1098.14501.
- [HQ] HOPKINS, MICHAEL J.; QUICK, GEREON. Hodge filtered complex bordism. *J. Topol.* **8** (2015), no. 1, 147–183. MR3335251, Zbl 1349.32009, arXiv:1212.2173, doi: 10.1112/jtopol/jtu021.
- [HS] HOPKINS, MICHAEL J.; SINGER, ISADORE M. Quadratic functions in geometry, topology, and M-theory. *J. Differential Geom.* **70** (2005), no. 3, 329–452. MR2192936 (2007b:53052), Zbl 1116.58018, arXiv:math/0211216, doi: 10.4310/jdg/1143642908.
- [KN] KOBAYASHI, SHOSHICHI; NOMIZU, KATSUMI. Foundations of differential geometry. I. Interscience Publishers, a division of John Wiley & Sons, New York-London, 1963. xi+329 pp. MR0152974 (27 #2945), Zbl 0119.37502, doi: 10.1126/science.143.3603.235-a.
- [KT] KÜBEL, ANDREAS; THOM, ANDREAS. Equivariant differential cohomology. Preprint, 2015. arXiv:1510.06392.
- [LMS] LASHOF, RICHARD K.; MAY, JOHN P.; SEGAL, GRAEME B. Equivariant bundles with abelian structural group. *Proceedings of the Northwestern Homotopy Theory Conference* (Evanston, Ill., 1982), 167–176, Contemp. Math., 19. Amer.

- Math. Soc., Providence, RI*, 1983. MR711050 (85b:55023), Zbl 0526.55020, doi: 10.1090/conm/019/711050.
- [Lur] LURIE, JACOB. Higher topos theory. *Annals of Mathematics Studies*, 170. *Princeton University Press, Princeton, NJ*, 2009. xviii+925 pp. ISBN: 978-0-691-14049-0; 0-691-14049-9. MR2522659 (2010j:18001), Zbl 1175.18001, arXiv:math/0608040, doi: 10.1515/9781400830558.
- [MaQ] MATHAI, VARGHESE; QUILLEN, DANIEL. Superconnections, Thom classes, and equivariant differential forms. *Topology* **25** (1986), no. 1, 85–110. MR836726 (87k:58006), Zbl 0592.55015, doi: 10.1016/0040-9383(86)90007-8.
- [Mei] MEINRENKEN, ECKHARD. Equivariant cohomology and the Cartan model. *Encyclopedia of Mathematical Physics*, 242–250. *Elsevier*, 2006. doi: 10.1016/B0-12-512666-2/00344-8.
- [NR] NARASIMHAN, MUDUMBAI S.; RAMANAN, SUNDARARAMAN. Existence of universal connections. *Amer. J. Math.* **83** (1961), 563–572. MR0133772 (24 #A3597), Zbl 0114.38203, doi: 10.2307/2372896.
- [Red] REDDEN, CORBETT. Harmonic forms on principal bundles. *Asian J. Math.* **16** (2012), no. 4, 637–660. MR3004281, Zbl 1271.58003, arXiv:0810.4578, doi: 10.4310/AJM.2012.v16.n4.a4.
- [Sch] SCHREIBER, URS. Differential cohomology in a cohesive infinity-topos. Preprint, 2013. arXiv:1310.7930.
- [SS] SIMONS, JAMES; SULLIVAN, DENNIS. Axiomatic characterization of ordinary differential cohomology. *J. Topol.* **1** (2008), no. 1, 45–56. MR2365651 (2009e:58035), Zbl 1163.57020, arXiv:math/0701077, doi: 10.1112/jtopol/jtm006.

(Corbett Redden) DEPARTMENT OF MATHEMATICS, LIU POST, LONG ISLAND UNIVERSITY, 720 NORTHERN BLVD, BROOKVILLE, NY 11548, USA
corbett.redden@liu.edu

This paper is available via <http://nyjm.albany.edu/j/2017/23-20.html>.