

Corrigendum to “Topology and arithmetic of resultants, I”, New York J. Math. 22 (2016), 801–821

Benson Farb and Jesse Wolfson

ABSTRACT. This note is meant to correct a mistake in [1]. A corrected version of [1] can be found on the archive: arXiv:1506.02713.

In Step 2 of Theorem 1.2 on page 808 of [1], we claimed that the map of Equation (3.3) (the map Ψ in Equation (1) below) is an isomorphism. This is not true, as pointed out to us by H. Spink and D. Tseng. However, we will see below that it is a bijective morphism. This has the effect that one needs to add the assumption that $\text{char}(K) = 0$ in Theorem 1.2, Corollary 1.3, and Theorem 1.7 of [1]. The corresponding point counts over \mathbb{F}_q still hold.

Step 2 of Theorem 1.2. As to the proof of Theorem 1.2 on page 808 of [1], the entirety of Step 2 should be deleted and replaced by the following.

Let $k \geq 0$. Define a morphism

$$\bar{\Psi} : \mathbb{A}^{m(d-nk)} \times \mathbb{A}^k \rightarrow \mathbb{A}^{md}$$

by

$$\bar{\Psi}(f_1, \dots, f_m, g) := (f_1 g^n, \dots, f_m g^n).$$

The restriction of $\bar{\Psi}$ to $\text{Poly}_n^{d-kn,m} \times \mathbb{A}^k$ gives a morphism

$$\Psi : \text{Poly}_n^{d-kn,m} \times \mathbb{A}^k \rightarrow R_{n,k}^{d,m} - R_{n,k+1}^{d,m} \tag{1}$$

where the target is the space of m -tuples of degree d polynomials with a common n -fold factor of degree equal to k , with no other common n -fold factors. We think of the map Ψ^{-1} as the (non-algebraic) map that extracts a common n -fold factor from a tuple of polynomials. We claim that:

- (i) For any field k the morphism Ψ is bijective.
- (ii) For $k = \mathbb{C}$, the map Ψ is a homeomorphism in the classical topology.

Received February 1, 2019.

2010 *Mathematics Subject Classification.* Primary: 55R80; secondary: 14N20.

Key words and phrases. 0-cycles, rational maps.

B.F. is supported in part by NSF Grant Nos. DMS-1105643 and DMS-1406209. J.W. is supported in part by NSF Grant No. DMS-1400349.

These facts will allow us to analyze $Polyn_n^{d,m}$ recursively. Note that the case $k = 0$ follows by definition:

$$Polyn_n^{d,m} := R_{n,0}^{d,m} - R_{n,1}^{d,m}.$$

To see (i): It is clear from the definitions that Ψ is surjective. The map Ψ is injective because there is a unique n -fold degree k factor in each $f_i g^n$, so if $f_i g^n = u_i v^n$ then this implies $g = v$ and so $f_i = u_i$.

To see (ii): First note that the spaces of polynomials in the range and domain of $\bar{\Psi}$ have Galois covers given by the corresponding spaces of (all possible orderings of) roots, with deck group the appropriate product of symmetric groups. The map $\bar{\Psi}$ lifts to a map between these spaces of roots:

$$\Phi : \mathbb{A}^{m(d-nk)} \times \mathbb{A}^k \rightarrow \mathbb{A}^{md}$$

given by

$$\Phi((\vec{r}_1, \dots, \vec{r}_m), \vec{s}) := ((\vec{r}_1, (\vec{s})^n), \dots, (\vec{r}_m, (\vec{s})^n))$$

where \vec{r}_i is the vector of d roots of f_i ; the vector of roots of g is denoted \vec{s} ; and where $(\vec{s})^n$ denotes the vector $(\vec{s}, \dots, \vec{s})$, where \vec{s} is repeated n times. It follows that the map Φ is closed, and hence the map $\bar{\Psi}$ is closed, and hence the map Ψ is closed. Since Ψ is bijective, it follows that Ψ is a homeomorphism.

Step 3 of Theorem 1.2. In Step 3 on page 808, one should insert the following after Equation (3.6).

We now claim that, when $\text{char}(K) = 0$ then

$$[Polyn_n^{d-kn,m}] \cdot \mathbb{L}^k = [R_{n,k}^{d,m}] - [R_{n,k+1}^{d,m}] \quad (2)$$

To see this, first note that we proved in Step 2 that the map Ψ in (1) is a bijective morphism on K -points for all fields K . It is known (see, e.g., Remark 4.1 of [2]) that if $\text{char}(K) = 0$ then a bijective morphism of K -varieties induces an equality $[X] = [Y]$ in the Grothendieck ring of K -varieties.

The line “Plugging in the expression from Equation (3.3) into Equation (3.6)” should now read: “Plugging in the expression from Equation (2) into Equation 3.6 ”

References

- [1] FARB, BENSON; WOLFSON, JESSE. Topology and arithmetic of resultants, I, *New York Jour. of Math.* **22** (2016), 801–821. arXiv:1506.02713, MR3548124, Zbl 1379.55016. 195
- [2] GÖTTSCHE, LOTHAR. On the motive of the Hilbert scheme of points on a surface, *Math. Res. Lett.* **8** (2001), no. 5-6, 613–627. arXiv:math/0007043v3, doi:10.4310/MRL.2001.v8.n5.a3, MR1879805, Zbl 1079.14500. 196

CORRIGENDUM TO “TOPOLOGY AND ARITHMETIC OF RESULTANTS, I” 197

(Benson Farb) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVENUE, CHICAGO, IL 60637, USA.
farb@math.uchicago.edu

(Jesse Wolfson) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT IRVINE, IRVINE, CA 92697, USA.
wolfson@uci.edu

This paper is available via <http://nyjm.albany.edu/j/2019/25-10.html>.