

Composition operators on the Dirichlet space of the upper half-plane

Ajay K. Sharma, Mehak Sharma and Kuldeep Raj

ABSTRACT. It is well known that Hardy and weighted Bergman spaces of the upper half-plane do not support compact composition operators (see [M99] and [SS03]). In this paper, we prove that unlike Hardy and Bergman spaces, the Dirichlet space of the upper half-plane does support compact composition operators. Furthermore, bounded analytic symbols, which in the case of Hardy and weighted Bergman spaces of the upper half-plane do not even induce bounded composition operators, can induce compact composition operators on the Dirichlet space of the upper half-plane.

CONTENTS

1. Introduction and preliminaries	198
2. Boundedness and compactness of C_φ on \mathcal{D}_{Π^+} .	199
References	205

1. Introduction and preliminaries

Let Ω be a domain in the complex plane \mathbb{C} and let φ be a holomorphic self-map of Ω . Then the equation $C_\varphi f = f \circ \varphi$, for f analytic in Ω , defines a composition operator C_φ with inducing map φ . During the past few decades, composition operators have been studied extensively on spaces of functions analytic on the open unit disk \mathbb{D} . As a consequence of the Littlewood subordination principle it is known that every analytic self-map φ of \mathbb{D} induces a bounded composition operator on Hardy and weighted Bergman spaces of the open unit disk. However, a self-map φ of \mathbb{D} does not necessarily induce a bounded composition operator on the Dirichlet space of the open unit disk \mathbb{D} . An obvious necessary condition for it is that φ be in the Dirichlet space of the open unit disk. But this condition is not sufficient. A necessary and sufficient condition for φ to induce a bounded composition operator on

Received July 16, 2018.

2010 *Mathematics Subject Classification.* 47B33; 30H15; 30H30.

Key words and phrases. composition operator; upper-half plane; Hardy space; Bergman space; Dirichlet space; counting function.

The first author was supported by the research grant 02011/30/2017/R&D II/12565 of NBHM(DAE) (India).

the Dirichlet space of the open unit disk is given in terms of the counting function and Carleson measures (see [JM97] and [Z98] and the references therein). For more about composition operators, we refer to [CoM95] and [S93].

If we move to Hardy and weighted Bergman spaces of the upper half-plane

$$\Pi^+ = \{z \in \mathbb{C} : \Im z > 0\},$$

the situation is entirely different. There do exist analytic self-maps of the upper half-plane which do not induce bounded composition operators. Moreover, Hardy and Bergman spaces of the upper half-plane do not support compact composition operators (see [M99] and [SS03]). Recent work on composition operators on Hardy and weighted Bergman spaces of the upper half-plane can be found in [BT12],[CKS17],[EW11],[EJ12],[M89],[M99],[SS03], and [SiSh80].

Composition operators on the Dirichlet space \mathcal{D}_{Π^+} of the upper half-plane remain untouched so far. In this paper, we characterize compact composition operators on the Dirichlet space of the upper half-plane. Recall that a function f that is analytic in the upper half-plane Π^+ belongs to the Dirichlet space \mathcal{D}_{Π^+} if and only if

$$\int_{\Pi^+} |f'(z)|^2 dA(z) < \infty,$$

where $dA(z) = dx dy$ is ordinary area measure. The norm on \mathcal{D}_{Π^+} is defined as

$$\|f\|_{\mathcal{D}_{\Pi^+}}^2 = |f(i)|^2 + \int_{\Pi^+} |f'(z)|^2 dA(z),$$

For $z \in \Pi^+$ and $0 < r < 1$ we define

$$S(z, r\Im z) = \{w \in \Pi^+ : |w - z| < r\Im z\}.$$

Then for $0 < r < 1/3$, there exists a positive integer M and a sequence $\{z_n\}$ in Π^+ such that

$$\cup_{n=1}^{\infty} S(z_n, r\Im z_n) = \Pi^+$$

and every point in Π^+ belongs to at most M sets in $\{S(z_n, 3r\Im z_n)\}_{n \in \mathbb{N}}$; see [KK01].

2. Boundedness and compactness of C_φ on \mathcal{D}_{Π^+} .

In this section we characterize bounded and compact composition operators on the Dirichlet space of the upper half plane in terms of the counting function and Carleson measures.

Let $\varphi : \Pi^+ \rightarrow \Pi^+$ be an analytic map and w be a point in Π^+ . Let $\{z_k\}$ be the points in the upper half plane for which $\varphi(z_k) = w$ counting multiplicities. Define the counting function

$$n_\varphi : \Pi^+ \rightarrow \mathbb{N} \cup \{\infty\}$$

by

$$n_\varphi(w) = \text{Card}\{z \in \Pi^+ : \varphi(z_k) = w\}$$

when the set $\{z \in \Pi^+ : \varphi(z_k) = w\}$ is finite, and $n_\varphi(w) = \infty$ otherwise. Also we set $n_\varphi(w) = 0$ if $w \notin \varphi(\Pi^+)$. Define

$$\mathfrak{N}_\varphi(w) = \begin{cases} n_\varphi(w) & (w \in \varphi(\Pi^+)) \\ 0 & \text{otherwise.} \end{cases}$$

The counting function \mathfrak{N}_φ produces the following non-univalent change of variables formula. The proof is standard, but we include it for completeness.

Proposition 2.1. *Let $0 < r < 1/3$ be fixed. Let $f \in \mathcal{D}_{\Pi^+}$ and φ be a non-constant analytic self-map of Π^+ . Then*

$$\int_{\Pi^+} |(f' \circ \varphi)(w)|^2 |\varphi'(w)|^2 dA(z) = \int_{\Pi^+} |f'(w)|^2 \mathfrak{N}_\varphi(w) dA(w). \quad (2.1)$$

Proof. Let

$$\Pi_0^+ = \{z \in \Pi^+ : \varphi'(z) \neq 0\}.$$

Then the set $\Pi^+ \setminus \Pi_0^+$ is at most countable. If $z \in \Pi_0^+$, then φ is one-one on $S(z, r\mathfrak{S}z)$ for some $r \in (0, 1/3)$. Thus we have

$$\int_{S(z, r\mathfrak{S}z)} |(f' \circ \varphi)(w)|^2 |\varphi'(w)|^2 dA(w) = \int_{\varphi(S(z, r\mathfrak{S}z))} |f'(w)|^2 dA(w).$$

The disks $S(z, r\mathfrak{S}z)$ form a cover for Π_0^+ and we can pick a countable sub-cover $\{S(z_n, r\mathfrak{S}z_n) : n \in \mathbb{N}\}$. Let $B_1 = S(z_1, ry_1)$ and

$$B_n = S(z_n, r\mathfrak{S}z_n,) \setminus \cup_{k=1}^{n-1} B_k$$

for all $n \geq 2$. Then $\{B_n : n \in \mathbb{N}\}$ is a pairwise disjoint cover of Π_0^+ . Using (2.1), we have

$$\begin{aligned} \int_{\Pi^+} |(f' \circ \varphi)(w)|^2 |\varphi'(w)|^2 dA(w) &= \sum_{n=1}^{\infty} \int_{B_n} |(f' \circ \varphi)(w)|^2 |\varphi'(w)|^2 dA(w) \\ &= \sum_{n=1}^{\infty} \int_{\varphi(B_n)} |f'(w)|^2 dA(w) \\ &= \int_{\Pi^+} |f'(w)|^2 \sum_{n=1}^{\infty} \chi_{\varphi(B_n)}(w) dA(w) \\ &= \int_{\Pi^+} |f'(w)|^2 \mathfrak{N}_\varphi(w) dA(w). \end{aligned}$$

This completes the proof of the non-univalent change of variables formula. \square

From now onwards, constants are denoted by C . They are positive and not necessarily the same at each occurrence.

Theorem 2.2. *Let $0 < r < 1/3$ be fixed and φ be a non-constant analytic self-map of Π^+ . Then C_φ is bounded on \mathcal{D}_{Π^+} if and only if*

$$\sup_{z \in \Pi^+} \frac{1}{(\Im z)^2} \int_{S(z, r\Im z)} \mathfrak{N}_\varphi(w) dA(w) < \infty. \quad (2.2)$$

Proof. Suppose that (2.2) holds. By the closed graph theorem, we need to show that $C_\varphi f \in \mathcal{D}_{\Pi^+}$ whenever $f \in \mathcal{D}_{\Pi^+}$. By Proposition 2.1, we have

$$\int_{\Pi^+} |(C_\varphi f)'(w)|^2 dA(w) = \int_{\Pi^+} |f'(w)|^2 \mathfrak{N}_\varphi(w) dA(w). \quad (2.3)$$

The right side of (2.3) is dominated by

$$\begin{aligned} & \sum_{n=1}^{\infty} \int_{S(z_n, r\Im z_n)} |f'(w)|^2 \mathfrak{N}_\varphi(w) dA(w) \\ & \leq \sum_{n=1}^{\infty} \sup\{|f'(w)|^2 : w \in S(z_n, r\Im z_n)\} \int_{S(z_n, r\Im z_n)} \mathfrak{N}_\varphi(w) dA(w). \end{aligned} \quad (2.4)$$

By the subharmonicity of $|f'(w)|^2$, (2.4) is further dominated by a constant multiple of

$$\begin{aligned} & \sum_{n=1}^{\infty} \left(\frac{1}{(\Im z_n)^2} \int_{S(z_n, 2r\Im z_n)} |f'(w)|^2 dA(w) \right) \int_{S(z_n, r\Im z_n)} \mathfrak{N}_\varphi(w) dA(w) \\ & \leq C \sum_{n=1}^{\infty} \int_{S(z_n, 2r\Im z_n)} |f'(w)|^2 dA(w) \leq CM \|f\|_{\mathcal{D}_{\Pi^+}}^2. \end{aligned}$$

Conversely, suppose that C_φ is bounded, then

$$\|C_\varphi f\|_{\mathcal{D}_{\Pi^+}}^2 \leq C \|f\|_{\mathcal{D}_{\Pi^+}}^2.$$

Let $f_z(w) = \Im z / (w - \bar{z})$. Then $f'_z(w) = \Im z / (w - \bar{z})^2$. If $z = x + iy$ and $w = t + iu$ are in Π^+ , then

$$\begin{aligned} \int_{\Pi^+} \left| \frac{1}{(w - \bar{z})^2} \right|^2 dA(w) &= \int_0^\infty \int_{-\infty}^\infty \frac{1}{((x-t)^2 + (y+u)^2)^2} dt du \\ &\leq \int_0^\infty \int_{-\infty}^\infty \frac{1}{(y+u)^3} \cdot \frac{y+u}{(x-t)^2 + (y+u)^2} dt du \\ &\leq \frac{C}{y^2}, \end{aligned}$$

where we have used the fact that

$$P_y(x, t) = \frac{1}{\pi} \frac{y}{(x-t)^2 + y^2} \quad (x, t \in \mathbb{R}, y > 0)$$

is the Poisson kernel for Π^+ and so

$$\frac{1}{\pi} \int_{-\infty}^\infty \frac{y+u}{(x-t)^2 + (y+u)^2} dt = 1.$$

Thus $f_z \in \mathcal{D}_{\Pi^+}$. Moreover,

$$\sup_{z \in \Pi^+} \|f_z\|_{\mathcal{D}_{\Pi^+}} \leq C.$$

So, from (2.3), we have

$$\begin{aligned} \frac{1}{(\Im z)^2} \int_{S(z, r\Im z)} \mathfrak{N}_\varphi(w) dA(w) &\leq C \int_{S(z, r\Im z)} |f'_z(w)|^2 \mathfrak{N}_\varphi(w) dA(w) \\ &\leq \|C_\varphi f\|_{\mathcal{D}_{\Pi^+}}^2 \leq C. \end{aligned}$$

Since $z \in \Pi^+$ is arbitrary, the desired result follows. \square

It is well known (see [M89], [M99] and [SS03]) that a linear fractional map

$$\tau(z) = \frac{az + b}{cz + d}, \quad a, b, c, d \in \mathbb{R} \text{ and } ad - bc > 0, \quad (2.5)$$

induces a bounded composition operator on the Hardy or weighted Bergman spaces of the upper half plane if and only if $c = 0$. However, in view of Theorem 2.2, every linear fractional map τ defined in (2.5) induces a bounded composition operator on \mathcal{D}_{Π^+} .

As in [CKS17], let $\widehat{\Pi^+}$ denote the set $\overline{\Pi^+} \cup \{\infty\}$. For any function $F(z)$,

$$\lim_{z \rightarrow \partial \widehat{\Pi^+}} F(z) = 0$$

means that

$$\sup_{z \in \Pi^+ \setminus K} |F(z)| \rightarrow 0$$

as the compact set $K \subset \Pi^+$ expands to the whole of Π^+ , or equivalently, that $F(z) \rightarrow 0$ as $\Im z \rightarrow 0^+$ and $F(z) \rightarrow 0$ as $|z| \rightarrow \infty$.

Theorem 2.3. *Let φ be a non-constant holomorphic self-map of Π^+ such that C_φ is bounded on \mathcal{D}_{Π^+} . Then C_φ is compact on \mathcal{D}_{Π^+} if and only if there exists $r \in (0, 1)$ such that*

$$\lim_{z \rightarrow \partial \widehat{\Pi^+}} \frac{1}{(\Im z)^2} \int_{S(z, r\Im z)} \mathfrak{N}_\varphi(w) dA(w) = 0. \quad (2.6)$$

Proof. Arguing by contradiction, first assume that C_φ is compact on \mathcal{D}_{Π^+} but (2.6) does not hold. Then there is a $\delta > 0$ and a sequence $\{z_n\}_{n \in \mathbb{N}}$ in Π^+ such that $\Im z_n \rightarrow 0$ or $|z_n| \rightarrow \infty$ and

$$\frac{1}{(\Im z)^2} \int_{S(z_n, r\Im z_n)} \mathfrak{N}_\varphi(w) dA(w) > \delta$$

for all $n \in \mathbb{N}$. For each $n \in \mathbb{N}$ consider the function

$$f_n(w) = \frac{\Im z_n}{w - \bar{z}_n}.$$

It is clear that f_n is norm bounded and $f_n \rightarrow 0$ uniformly on compact subsets of Π^+ as $\Im z_n \rightarrow 0$ or $|z_n| \rightarrow \infty$. Thus $\|C_\varphi f_n\|_{\mathcal{D}_{\Pi^+}} \rightarrow 0$ as $\Im z_n \rightarrow 0$ or $|z_n| \rightarrow \infty$. On the other hand, by (2.3), we have

$$\begin{aligned} \|C_\varphi f_n\|_{\mathcal{D}_{\Pi^+}} &\geq \int_{S(z_n, r\Im z_n)} |f'_n(w)|^2 \mathfrak{N}_\varphi(w) dA(w) \\ &\geq \frac{C}{(\Im z_n)^2} \int_{S(z_n, r\Im z_n)} \mathfrak{N}_\varphi(w) dA(w) > C\delta, \end{aligned}$$

which is a contradiction.

Next, assume that (2.6) holds. Then for each $\epsilon > 0$ there is a compact subset K of Π^+ such that

$$\int_{S(z, r\Im z)} \mathfrak{N}_\varphi(w) dA(w) < \epsilon (\Im z)^2 \quad (2.7)$$

whenever $z \in \Pi^+ \setminus K$. Let $\{f_m\}$ be a sequence in \mathcal{D}_{Π^+} such that

$$\sup_m \|f_m\|_{\mathcal{D}_{\Pi^+}} \leq M_1$$

and $f_m \rightarrow 0$ uniformly on compact subsets of Π^+ as $m \rightarrow \infty$. Then

$$\begin{aligned} \|C_\varphi f_m\|_{\mathcal{D}_{\Pi^+}}^2 &= |f_m(\varphi(i))|^2 + \int_{\Pi^+} |f'_m(w)|^2 \mathfrak{N}_\varphi(w) dA(w) \\ &= |f_m(\varphi(i))|^2 + \int_K |f'_m(w)|^2 \mathfrak{N}_\varphi(w) dA(w) \\ &\quad + \int_{\Pi^+ \setminus K} |f'_m(w)|^2 \mathfrak{N}_\varphi(w) dA(w). \end{aligned}$$

Note that $|f_m(\varphi(i))|^2 \rightarrow 0$ as $m \rightarrow \infty$ and

$$\int_K |f'_m(w)|^2 \mathfrak{N}_\varphi(w) dA(w) \rightarrow 0 \quad \text{as } m \rightarrow \infty. \quad (2.8)$$

Thus to prove that C_φ is compact on \mathcal{D}_{Π^+} we just need to show that

$$\int_{\Pi^+ \setminus K} |f'_m(w)|^2 \mathfrak{N}_\varphi(w) dA(w) \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

As in the proof of Theorem 2.2, the above term is dominated by

$$\sum_{n=1}^{\infty} \left(\frac{1}{(\Im z_n)^2} \int_{S(z_n, 2r\Im z_n)} |f'_m(w)|^2 dA(w) \right) \int_{S(z_n, r\Im z_n) \cap (\Pi^+ \setminus K)} \mathfrak{N}_\varphi(w) dA(w).$$

By (2.7), we have

$$\int_{\Pi^+ \setminus K} |f'_m(w)|^2 \mathfrak{N}_\varphi(w) dA(w) < \epsilon \sum_{n=1}^{\infty} \int_{S(z_n, 2r\Im z_n)} |f'_m(w)|^2 dA(w) \leq \epsilon M M_1 \quad (2.9)$$

Combining (2.8) and (2.9), we obtain

$$\int_{\Pi^+} |f'_m(w)|^2 \mathfrak{N}_\varphi(z) dA(w) \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Hence C_φ is compact on \mathcal{D}_{Π^+} . \square

In [M89], Matache proved that if φ is a bounded analytic self mapping of Π^+ , then C_φ cannot be bounded on the Hardy space $H^p(\Pi^+)$. Also, bounded analytic self-maps of Π^+ cannot induce bounded composition operators on weighted Bergman spaces of the upper half-plane; see [SS03]. As an application of Theorem 2.3, we prove that there are non-trivial analytic self-maps of the upper half-plane that induce compact composition operators on \mathcal{D}_{Π^+} .

Corollary 2.4. *Let φ be a conformal mapping from Π^+ to a relatively compact subset of Π^+ . Then φ induces a compact composition operator on \mathcal{D}_{Π^+} .*

Proof. Suppose that $K = \varphi(\Pi^+)$ is a relatively compact subset of Π^+ . Then

$$\mathfrak{N}_\varphi(w) = \begin{cases} 1 & \text{if } w \in K \\ 0 & \text{if } w \notin K. \end{cases}$$

Therefore,

$$\begin{aligned} & \sup_{z \in \Pi^+} \frac{1}{(\Im z)^2} \int_{S(z, r\Im z)} \mathfrak{N}_\varphi(w) dA(w) \\ &= \sup_{z \in \Pi^+} \left(\frac{1}{(\Im z)^2} \int_{S(z, r\Im z) \cap K} \mathfrak{N}_\varphi(w) dA(w) \right. \\ & \quad \left. + \frac{1}{(\Im z)^2} \int_{S(z, r\Im z) \cap (\Pi^+ \setminus K)} \mathfrak{N}_\varphi(w) dA(w) \right) \\ & \leq C \sup_{z \in \Pi^+} \frac{1}{(\Im z)^2} A(S(z, r\Im z) \cap \Pi^+) \leq C. \end{aligned}$$

Thus by (2.2), φ induces a bounded composition operator on \mathcal{D}_{Π^+} .

Next, we prove that φ induces a compact composition operator on \mathcal{D}_{Π^+} . In view of Theorem 2.3, we need to show that

$$\lim_{z \rightarrow \partial \Pi^+} \frac{1}{(\Im z)^2} \int_{S(z, r\Im z)} \mathfrak{N}_\varphi(w) dA(w) = 0. \quad (2.10)$$

Since $K = \varphi(\Pi^+)$ is a relatively compact subset of Π^+ and

$$\sup_{z \in \Pi^+} \frac{1}{(\Im z)^2} \int_{S(z, r\Im z)} \mathfrak{N}_\varphi(w) dA(w)$$

is finite, so (2.10) is vacuously true. Hence φ induces a compact composition operator on \mathcal{D}_{Π^+} . \square

Recall that the *valence* of an analytic self-mapping φ of Π^+ is

$$N = \sup_{w \in \Pi^+} n_\varphi(w).$$

The function φ is said to have *bounded valence* if $N < \infty$, that is, if there is a positive integer N such that φ takes every value at most N times in Π^+ .

Corollary 2.5. *Let φ be of bounded valence and φ maps Π^+ to a relatively compact subset of Π^+ . Then φ induces a compact composition operator on \mathcal{D}_{Π^+} .*

The proof follows on the same lines as the proof of Corollary 2.4. We omit the details.

Example 2.6. Let

$$\varphi(z) = 2i + \frac{1}{(z+i)\log(z+ei)} \quad (z \in \Pi^+).$$

Then $\varphi(\Pi^+)$ is a relatively compact subset of Π^+ (see Example 4.4 in [CKS17]). Thus φ induces a compact composition operator on \mathcal{D}_{Π^+} .

Acknowledgements. The authors thank the anonymous referee for providing suggestions that improved the paper. The first author thanks NBHM(DAE) (India) for the project grant 02011/30/2017/R&D II/12565.

References

- [BT12] BERCOVICI, HARI; TIMOTIN, DAN. A note on composition operators in a half-plane. *Arch. Math. (Basel)* **99** (2012), no. 6, 567–576. [MR3001560](#), [Zbl 1262.47035](#), [arXiv:1205.1489](#), doi: [10.1007/s00013-012-0450-7](#). 199
- [CKS17] CHOE, BOO RIM; KOO, HYUNGWOON; SMITH, WAYNE. Difference of composition operators over the half-plane. *Trans. Amer. Math. Soc.* **369** (2017), no. 5, 3173–3205. [MR3605968](#), [Zbl 06682369](#), doi: [10.1090/tran/6742](#). 199, 202, 205
- [CoM95] COWEN, CARL C.; MACCLUER, BARBARA D. Composition operators on spaces of analytic functions. Studies in Advanced Mathematics. *CRC Press Boca Raton, FL*, 1995. xii+388 pp. ISBN: 0-8493-8492-3. [MR1397026](#), [Zbl 0873.47017](#). 199
- [EJ12] ELLIOTT, SAM; JURY, MICHAEL T. Composition operators on Hardy spaces of a half-plane. *Bull. Lond. Math. Soc.* **44** (2012), no. 3, 489–495. [MR2966995](#), [Zbl 1248.47025](#), [arXiv:0907.0350](#), doi: [10.1112/blms/bdr110](#). 199
- [EW11] ELLIOTT, SAM J.; WYNN, ANDREW. Composition operators on weighted Bergman spaces of a half-plane. *Proc. Edinb. Math. Soc.* (2) **54** (2011), no. 2, 373–379. [MR2794660](#), [Zbl 1230.47047](#), [arXiv:0910.0408](#), doi: [10.1017/S0013091509001412](#). 199
- [JM97] JOVOVIĆ, MIRJANA; MACCLUER, BARBARA. Composition operators on Dirichlet spaces. *Acta Sci. Math. (Szeged)* **63** (1997), no. 1–2, 229–247. [MR1459789](#), [Zbl 0880.47019](#). 199
- [KK01] KANG, SI HO; KIM, JA YOUNG. Harmonic Bergman spaces of the half-space and their some operators. *Bull. Korean Math. Soc.* **38** (2001), no. 4, 773–786. [MR1865836](#), [Zbl 1003.47019](#). 199
- [M89] MATACHE, VALENTIN. Composition operators on H^p of the upper half-plane. *An. Univ. Timișoara Ser. Științ. Mat.* **27** (1989), no. 1, 63–66. [MR1140524](#), [Zbl 0791.47031](#). 199, 202, 204
- [M99] MATACHE, VALENTIN. Composition operators on Hardy spaces of a half-plane. *Proc. Amer. Math. Soc.* **127** (1999) no. 5, 1483–1491. [MR1625773](#), [Zbl 0916.47022](#), doi: [10.1090/S0002-9939-99-05060-1](#). 198, 199, 202

- [S93] SHAPIRO, JOEL H. Composition operators and classical function theory. Universitext: Tracts in Mathematics. *Springer-Verlag, New York*, 1993. xvi+223 pp. ISBN: 0-387-94067-7. MR1237406, Zbl 0791.30033, doi: [10.1007/978-1-4612-0887-7](https://doi.org/10.1007/978-1-4612-0887-7). 199
- [SS03] SHAPIRO, JOEL H.; SMITH, WAYNE. Hardy spaces that support no compact composition operators. *J. Funct. Anal.* **205** (2003), no. 1, 62–89. MR2020208, Zbl 1041.46019, doi: [10.1016/S0022-1236\(03\)00215-5](https://doi.org/10.1016/S0022-1236(03)00215-5). 198, 199, 202, 204
- [SiSh80] SINGH, R.K.; SHARMA, S.D. Non-compact composition operators. *Bull. Austral. Math. Soc.* **21** (1980), no. 1, 125–130. MR0569092, Zbl 0411.47024, doi: [10.1017/S0004972700011345](https://doi.org/10.1017/S0004972700011345). 199
- [Z98] ZORBOSKA, NINA. Composition operators on weighted Dirichlet spaces. *Proc. Amer. Math. Soc.* **126** (1998), no. 7, 2013–2023. MR1443862, Zbl 0894.47023, doi: [10.1090/S0002-9939-98-04266-X](https://doi.org/10.1090/S0002-9939-98-04266-X). 199

(A. Sharma) DEPARTMENT OF MATHEMATICS, CENTRAL UNIVERSITY OF JAMMU, (BAGLA) RAYA-SUCHANI, SAMBA-181143, J&K, INDIA.
aksju.76@yahoo.com

(M. Sharma) DEPARTMENT OF MATHEMATICS, CENTRAL UNIVERSITY OF JAMMU, (BAGLA) RAYA-SUCHANI, SAMBA-181143, J&K, INDIA.
smehak14@gmail.com

(Kuldip Raj) DEPARTMENT OF MATHEMATICS, SHRI MATA VAISHNO DEVI UNIVERSITY, KAKRYAL, KATRA-182320, J&K, INDIA.
kuldipraj68@gmail.com

This paper is available via <http://nyjm.albany.edu/j/2019/25-11.html>.