

von Neumann’s inequality for the Hartogs triangle

**Sameer Chavan, Shubham Jain
and Paramita Pramanick**

ABSTRACT. For a commuting pair T of bounded linear operators T_1 and T_2 on a Hilbert space \mathcal{H} , let $D_T = T_2^*T_2 - T_1^*T_1$. If $T_2^*D_T T_2 \leq D_T$ and the Taylor spectrum of T is contained in the Hartogs triangle Δ_H , then for any bounded holomorphic function ϕ on Δ_H , $\|\phi(T)\| \leq \|\phi\|_\infty$. We deduce this fact from an analogue of von Neumann’s inequality for bounded domains in \mathbb{C}^d . The proof of the latter closely follows the model theory approach as developed in [1].

CONTENTS

1. Introduction	791
2. von Neumann’s inequality for a bounded domain	794
References	798

1. Introduction

1.1. Hardy space of the Hartogs triangle. The *Hartogs triangle* Δ_H is given by

$$\Delta_H = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1| < |z_2| < 1\}.$$

Clearly,

$$\Delta_H = \{(z_1, z_2) \in \mathbb{D}^2 : (z_2\bar{z}_2 - z_1\bar{z}_1)(1 - z_2\bar{z}_2) > 0\}, \tag{1}$$

where \mathbb{D} denotes the open unit disc. Moreover, Δ_H is bi-holomorphic to $\mathbb{D} \times \mathbb{D}^*$ via the map $(z_1, z_2) \rightarrow (\frac{z_1}{z_2}, z_2)$, where $\mathbb{D}^* = \mathbb{D} \setminus \{0\}$. Thus, Δ_H is a holomorphically convex domain. However, it is easy to see that Δ_H is not polynomially convex (refer to [19] for the basics of several complex variables and also to [20] for an exposition surveying classical and recent results on the Hartogs triangle).

Let $I = (0, 1)$. For $(s, t) \in I \times I$, let

$$\Delta_{H,s,t} = \{(z_1, z_2) \in \mathbb{C}^2 : \frac{|z_1|}{s} < |z_2| < t\}.$$

Received January 3, 2022.

2020 *Mathematics Subject Classification.* Primary 47A13; Secondary 32A10.

Key words and phrases. Hartogs triangle, commuting tuple, von Neumann inequality.

The distinguished boundary $\partial_d(\Delta_{H_{s,t}})$ of $\Delta_{H_{s,t}}$ is given by

$$\partial_d(\Delta_{H_{s,t}}) = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1| = st, |z_2| = t\}.$$

Following [12, Section 3] (see also [9, Section 6]), we define the *Hardy space on the Hartogs triangle* Δ_H as

$$H^2(\Delta_H) = \left\{ f \in \text{Hol}(\Delta_H) : \sup_{(s,t) \in I \times I} \frac{1}{4\pi^2} \int_{\partial_d(\Delta_{H_{s,t}})} |f|^2 d\sigma_{s,t} < \infty \right\},$$

where $\sigma_{s,t}$ is the surface area measure on $\partial_d(\Delta_{H_{s,t}})$ induced by the Lebesgue measure on the unit torus $\mathbb{T} \times \mathbb{T}$. Then $H^2(\Delta_H)$ is a reproducing kernel Hilbert space endowed with the norm

$$\|f\|_{H^2(\Delta_H)}^2 = \sup_{(s,t) \in I \times I} \frac{1}{4\pi^2} \int_0^{2\pi} \int_0^{2\pi} |f(ste^{i\theta}, te^{i\gamma})|^2 st^2 d\theta d\gamma.$$

A domain Ω in \mathbb{C}^d is a nonempty open subset of \mathbb{C}^d . For a bounded domain Ω in \mathbb{C}^d , let $H^\infty(\Omega)$ denote the Banach algebra of bounded holomorphic functions on Ω endowed with the sup norm $\|\cdot\|_{\infty, \Omega}$. Note that $H^\infty(\mathbb{D}^2) \subsetneq H^\infty(\Delta_H)$ (consider $f(z_1, z_2) = z_1/z_2$). However, any function holomorphic in a neighborhood of Δ_H extends analytically to \mathbb{D}^2 (see [20, Proposition 1.1]). The multiplier algebra of a Hilbert space \mathcal{H} of complex-valued holomorphic functions is denoted by $\text{Mult}(\mathcal{H})$. If $\phi \in \text{Mult}(\mathcal{H})$, then the operator M_ϕ of multiplication by ϕ defines a bounded linear operator. The multiplier norm of ϕ is the operator norm of M_ϕ .

Lemma 1.1. *The following statements are valid:*

(i) *The reproducing kernel for $H^2(\Delta_H)$ is given by*

$$\kappa_H(z, w) = \frac{1}{(z_2 \bar{w}_2 - z_1 \bar{w}_1)(1 - z_2 \bar{w}_2)}.$$

(ii) *The multiplier algebra of $H^2(\Delta_H)$ is equal to $H^\infty(\Delta_H)$ with equality of norms.*

Proof. Note that (i) is precisely [12, Proposition 3.2]. To see (ii), note that the multiplier algebra of $H^2(\Delta_H)$ is contained in $H^\infty(\Delta_H)$ and $\|\phi\|_{\infty, \Delta_H} \leq \|M_\phi\|$ (see [14]). Also, for each $\phi \in H^\infty(\Delta_H)$,

$$\|\phi f\|_{H^2(\Delta_H)} \leq \|\phi\|_{\infty, \Delta_H} \|f\|_{H^2(\Delta_H)}, \quad f \in H^2(\Delta_H).$$

Thus, the multiplier norm of ϕ is equal to $\|\phi\|_{\infty, \Delta_H}$. \square

1.2. Main theorem. Let \mathcal{H} be a complex Hilbert space and let d be a positive integer. By a *commuting d -tuple* $T = (T_1, \dots, T_d)$ on \mathcal{H} , we understand the d -tuple of bounded linear operators T_1, \dots, T_d on \mathcal{H} such that $T_j T_k = T_k T_j$ for all integers $1 \leq j \neq k \leq d$. The Taylor spectrum and approximate point-spectrum of a commuting d -tuple T on \mathcal{H} are denoted by $\sigma(T)$ and $\sigma_{\text{ap}}(T)$, respectively

(the reader is referred to [5] for definition and basic information on the various joint spectra of commuting tuples).

The purpose of this note is to prove von Neumann's inequality for the Hartogs triangle (cf. [22, Theorem 1.1]).

Theorem 1.2. *If $T = (T_1, T_2)$ is a commuting 2-tuple on \mathcal{H} such that $\sigma(T) \subset \Delta_H$ and $T_2^*(T_2^*T_2 - T_1^*T_1)T_2 \leq T_2^*T_2 - T_1^*T_1$, then*

$$\|\phi(T)\| \leq \|\phi\|_{\infty, \Delta_H}, \quad \phi \in H^\infty(\Delta_H). \tag{2}$$

Remark 1.3. Set $D_T := T_2^*T_2 - T_1^*T_1$ and $E_T := D_T - T_2^*D_T T_2$.

(1) By the hereditary functional calculus (see [1]),

$$T_2^*D_T T_2 \leq D_T \iff \frac{1}{\kappa_H}(T, T^*) \geq 0,$$

where $\frac{1}{\kappa_H}(z, w) = (z_2\bar{w}_2 - z_1\bar{w}_1)(1 - z_2\bar{w}_2)$ (cf. (1)).

(2) The condition $E_T \geq 0$ need not imply $\sigma(T) \subset \Delta_H$ (cf. [22, Lemma 4.1]). Indeed, any commuting subnormal 2-tuple T with minimal normal extension N such that

$$\sigma(N) \subset \{(z_1, z_2) \in \mathbb{C}^2 : |z_1| > |z_2| > 1\}$$

satisfies $E_T \geq 0$ (consult the spectral theorem for commuting normal operators). However, since $\sigma(N) \subseteq \sigma(T)$ (see [17]) is a nonempty set (see [5]), $\sigma(T)$ intersects with the complement of Δ_H . Also, by the spectral theorem for commuting normal operators,

$$T_1^*T_1 \geq T_2^*T_2 \geq I.$$

This example also shows that $E_T \geq 0$ need not imply

$$T_1^*T_1 \leq T_2^*T_2 \leq I.$$

(3) Any commuting 2-tuple T such that $E_T \geq 0$ satisfies

$$\sigma_{\text{ap}}(T) \cap \overline{\mathbb{D}}^2 \subseteq \overline{\Delta_H}. \tag{3}$$

Indeed, if $(\lambda_1, \lambda_2) \in \sigma_{\text{ap}}(T)$, then there exists a sequence $\{x_n\}_{n=1}^\infty$ of unit vectors such that

$$\lim_{n \rightarrow \infty} \langle E_T x_n, x_n \rangle = 0 \implies (1 - |\lambda_2|^2)(|\lambda_2|^2 - |\lambda_1|^2) \geq 0,$$

and hence (1) yields the inclusion (3).

Finally, note that (2) implies that Δ_H is a spectral domain for T (see [1, Definition 1.46]).

Since Δ_H is not a P -ball in the sense [16] (indeed, Δ_H is a Reinhardt domain, which is not complete), [16, Corollary 5.4] is not applicable to the case of the Hartogs triangle. It turns out that an application of Lemma 1.1 together with an analogue of von Neumann's inequality (see Theorem 2.1) yields Theorem 1.2.

2. von Neumann's inequality for a bounded domain

Theorem 2.1. *Let Ω be a bounded domain in \mathbb{C}^d . Assume that there exists a reproducing kernel Hilbert space \mathcal{H}_κ with reproducing kernel $\kappa : \Omega \times \Omega \rightarrow \mathbb{C} \setminus \{0\}$ such that $\frac{1}{\kappa}$ is a hereditary function on Ω . If T is a commuting d -tuple on \mathcal{H} such that $\sigma(T) \subset \Omega$ and $\frac{1}{\kappa}(T, T^*) \geq 0$, then*

$$\|\phi(T)\| \leq \|M_\phi\|, \quad \phi \in \text{Mult}(\mathcal{H}_\kappa). \quad (4)$$

If, in addition, the multiplier algebra $\text{Mult}(\mathcal{H}_\kappa)$ of \mathcal{H}_κ is equal to $H^\infty(\Omega)$ (with equality of norms), then

$$\|\phi(T)\| \leq \|\phi\|_{\infty, \Omega}, \quad \phi \in H^\infty(\Omega). \quad (5)$$

Remark 2.2. The above result is applicable to a number of domains Ω in \mathbb{C}^d allowing us to recover variants of several known results ([3, 1, 8, 13, 16, 22]):

- Letting $\Omega = \mathbb{D}^d$ and $\kappa(z, w) = \frac{1}{\prod_{j=1}^d (1 - \bar{w}_j z_j)}$, $z, w \in \mathbb{D}^d$, we get (5) for any commuting d -tuple T satisfying $\sigma(T) \subset \mathbb{D}^d$ and

$$\sum_{\substack{\alpha_j \in \{0,1\} \\ j=1,\dots,d}} (-1)^{|\alpha|} T^{*\alpha} T^\alpha \geq 0,$$

where $T^\alpha = \prod_{j=1}^d T_j^{\alpha_j}$, $|\alpha| = \alpha_1 + \dots + \alpha_d$, $\alpha = (\alpha_1, \dots, \alpha_d)$. This recovers a variant of von Neumann inequality for the polydisc (cf. [6, Theorem 4.7] and [21, Theorem 1.9.1]).

- Letting $\Omega = \mathbb{B}^d$ and $\kappa(z, w) = \frac{1}{1 - \langle z, w \rangle}$, $z, w \in \mathbb{B}^d$, we get (4) for any commuting d -tuple T satisfying $\sigma(T) \subset \mathbb{B}^d$ and $\sum_{j=1}^d T_j^* T_j \leq I$. This recovers Drury's theorem (see [8, pp 300-301]).
- Letting $\Omega = \mathbb{B}^d$ and $\kappa(z, w) = \frac{1}{(1 - \langle z, w \rangle)^d}$, $z, w \in \mathbb{B}^d$, we get (5) for any commuting d -tuple T satisfying $\sigma(T) \subset \mathbb{B}^d$ and

$$\sum_{0 \leq k \leq d} (-1)^k \binom{d}{k} \sum_{|\alpha|=k} \frac{k!}{\alpha!} T^{*\alpha} T^\alpha \geq 0.$$

This is a consequence of [13, Theorem 11].

- For $0 < r < 1$, letting $\Omega = \mathbb{A}_r = \{z \in \mathbb{C} : r < |z| < 1\}$ and $\kappa(z, w) = \frac{1-r^2}{(1-z\bar{w})(1-r^2/z\bar{w})}$, $z, w \in \mathbb{A}_r$, we get (4) for any bounded linear operator T satisfying $\sigma(T) \subset \mathbb{A}_r$ and

$$r^2 I - (1 + r^2) T^* T + T^{*2} T^2 \leq 0.$$

This recovers a part of [22, Theorem 1.1].

Recall from [2] (see also [1, Chapter 4]) that h is a hereditary function on Ω if h is a mapping from $\Omega \times \Omega$ to \mathbb{C} such that $(z, w) \rightarrow h(z, \bar{w})$ is holomorphic function on $\Omega \times \Omega$. A dyad in $\text{Her}(\Omega)$ is a function of the form $(z, w) \rightarrow \overline{\psi(w)} \phi(z)$,

$z, w \in \Omega$, for some functions ϕ and ψ in $\text{Hol}(\Omega)$. Here $\text{Her}(\Omega)$ denotes the set of hereditary functions on Ω .

It has been recorded in [1, Remark 4.15] that for any domain Ω in \mathbb{C}^d , the hereditary functional calculus is well-defined and satisfies the properties as listed in [1, Lemma 4.13]. Alternatively, following [4, Section 1], we may use the iterated Cauchy-Weil integral (see [5, Corollary 5.4]) to define

$$h(T) = \frac{1}{(2\pi i)^{2d}} \int_{\partial\Omega'} \int_{\partial\Omega'} h(z, \bar{w})M(\bar{w} - T^*)M(z - T) \wedge dz \wedge d\bar{w}$$

for every $h \in \text{Her}(\Omega)$, where M is the Martinelli kernel (see [5]) and Ω' is an open domain with C^1 -boundary such that $\sigma(T) \subset \Omega' \subset \overline{\Omega'} \subset \Omega$. In the proof of Theorem 2.1, we need the continuity of the hereditary functional calculus:

Lemma 2.3. *Let Ω be a bounded domain in \mathbb{C}^d . If $\{h_n\}_{n=1}^\infty$ is a sequence in $\text{Her}(\Omega)$, $h \in \text{Her}(\Omega)$, and $h_n \rightarrow h$ uniformly on compact subsets of $\Omega \times \Omega$, then $h_n(T) \rightarrow h(T)$ with respect to the operator norm, whenever T is a commuting d -tuple with $\sigma(T) \subset \Omega$.*

Proof. Let T be a commuting d -tuple satisfying $\sigma(T) \subset \Omega$. As pointed out in [1, Remark 4.15], the proof of this fact relies on the nuclearity of $\text{Hol}(\Omega)$ (see [10, Theorem V.4.1], [7, Theorem 3.64]) and [15, Proposition, p 113]. Alternatively, one may adapt the proof of [1, Proposition 2.87] to the present situation. Since $\sigma(T)$ is contained in Ω' and

$$\lambda \mapsto \left(\sum_{j=1}^d (T_j - \lambda_j)^*(T_j - \lambda_j) \right)^{-1}$$

is bounded on $\partial\Omega'$ (since the approximate point-spectrum is a closed subset of Ω' and the inverse map on invertible bounded linear operators is continuous), by [5, Lemma 5.11], the map $(z, w) \mapsto M(\bar{w} - T^*)M(z - T)$ is bounded on $\partial\Omega'$. It follows that

$$h_n(z, w)M(\bar{w} - T^*)M(z - T) \rightarrow h(z, w)M(\bar{w} - T^*)M(z - T)$$

in the operator norm, uniformly for $z, w \in \partial\Omega'$. Hence, $h_n(T) \rightarrow h(T)$. □

If Ω is a Reinhardt domain in \mathbb{C}^d , then by [11, Proposition 1.7.1(b)&(c)], $(z, w) \rightarrow h(z, \bar{w})$ has a Laurent series expansion converging compactly on $\Omega \times \Omega$, and hence for some complex numbers $a_{\alpha, \beta}$,

$$h(T) = \sum_{\alpha, \beta \in \mathbb{Z}^d} a_{\alpha, \beta} T^{*\beta} T^\alpha,$$

where \mathbb{Z} denotes the set of integers. This observation is applicable to the Hartogs triangle (see [20, Eq (1.1)]).

The following is a counter-part of [1, Theorem 4.4] for any bounded domain in \mathbb{C}^d .

Lemma 2.4. *Let Ω be a bounded domain in \mathbb{C}^d . If A is a positive semi-definite hereditary function on Ω , then there exists a sequence $\{f_n\}_{n=1}^\infty$ in $\text{Hol}(\Omega)$ such that*

$$A(z, w) = \sum_{n=1}^{\infty} \overline{f_n(w)} f_n(z), \quad z, w \in \Omega, \quad (6)$$

where the series in (6) converges uniformly on compact subsets of $\Omega \times \Omega$.

Proof. We closely follow the proof of [1, Theorem 4.4]. Since A is positive semi-definite on Ω , by Moore's theorem (see [1, Theorem 2.5]), there exist a Hilbert space \mathcal{M} and a function $u : \Omega \rightarrow \mathcal{M}$ such that

$$\vee\{u(z) : z \in \Omega\} = \mathcal{M}, \quad A(z, w) = \langle u(z), u(w) \rangle_{\mathcal{M}}, \quad z, w \in \Omega.$$

By the reproducing property of \mathcal{M} , u is easily seen to be a weakly holomorphic function, and hence by [1, Lemma 2.90], u is holomorphic. Also, since Ω is separable, so is \mathcal{M} . Let $\{e_n\}_{n=1}^N$ be an orthonormal basis for \mathcal{M} , where N is a positive integer or $N = \infty$. By Parseval's identity and the continuity of the inner-product, we have

$$A(z, w) = \langle u(z), u(w) \rangle_{\mathcal{M}} = \sum_{n=1}^N \overline{\langle u(w), e_n \rangle_{\mathcal{M}}} \langle u(z), e_n \rangle_{\mathcal{M}}, \quad z, w \in \Omega.$$

Therefore, if we define $f_n = \langle u(\cdot), e_n \rangle_{\mathcal{M}}$, $n = 1, \dots, N$, then $f_n \in \text{Hol}(\Omega)$ and (6) holds (if $N < \infty$, then let $f_n = 0$ for $n > N$). Letting $z = w$ in (6), we may conclude from Dini's theorem (see [18, Theorem 7.13]) that $\sum_{n=1}^{\infty} |f_n(\cdot)|^2$ converges uniformly on compact subsets of Ω . We obtain the desired conclusion now from the Cauchy-Schwarz inequality. \square

Proof of Theorem 2.1. Assume that $\sigma(T) \subset \Omega$ and $\frac{1}{\kappa}(T, T^*) \geq 0$. Let $\phi \in \text{Mult}(\mathcal{H}_{\kappa})$. After multiplying by a scalar, if required, we may assume that ϕ belongs to the closed unit ball of $\text{Mult}(\mathcal{H}_{\kappa})$. By [14, Theorem 5.21], there exists a positive semi-definite kernel $A : \Omega \times \Omega \rightarrow \mathbb{C}$ such that

$$(1 - \overline{\phi(w)}\phi(z))\kappa(z, w) = A(z, w), \quad z, w \in \Omega. \quad (7)$$

Thus, we have the model formula

$$1 - \overline{\phi(w)}\phi(z) = \frac{A(z, w)}{\kappa(z, w)}, \quad z, w \in \Omega.$$

Since A is a positive semi-definite element of $\text{Her}(\Omega)$, by Lemma 2.4, there exists a sequence $\{f_n\}_{n=1}^\infty$ of elements in $\text{Hol}(\Omega)$ such that

$$1 - \phi(z)\overline{\phi(w)} = \sum_{n=1}^{\infty} \overline{f_n(w)} \left(\frac{1}{\kappa}(z, w) \right) f_n(z)$$

converges uniformly on compact subsets of $\Omega \times \Omega$. Since $\sigma(T) \subset \Omega$, by the continuity of the hereditary functional calculus on Ω (see Lemma 2.3),

$$I - \phi(T)^* \phi(T) = \sum_{n=1}^{\infty} f_n(T)^* \left(\frac{1}{\kappa}(T, T^*) \right) f_n(T).$$

Hence, by the assumption $\frac{1}{\kappa}(T, T^*) \geq 0$, we conclude that $I - \phi(T)^* \phi(T) \geq 0$ or equivalently, $\|\phi(T)\| \leq 1$ completing the proof of the first part. The remaining part is now immediate. \square

Proof of Theorem 1.2. An application of Lemma 1.1 to the choice $\kappa = \kappa_H$ together with Theorem 2.1 and Remark 1.3(1) yields (2). \square

Let us discuss some consequences of Theorem 1.2.

Corollary 2.5. *If $T = (T_1, T_2)$ is a commuting 2-tuple on \mathcal{H} such that $\sigma(T) \subset \Delta_H$ and $T_2^*(T_2^*T_2 - T_1^*T_1)T_2 \leq T_2^*T_2 - T_1^*T_1$, then*

$$T_1^*T_1 \leq T_2^*T_2 \leq I. \tag{8}$$

Proof. Letting $\phi(z_1, z_2) = z_2$ in (2), we conclude that $T_2^*T_2 \leq I$. To see the remaining inequality, by the spectral mapping property (see [5, Theorem 5.19]), T_2 is invertible. Applying (2) to the bounded holomorphic function $\phi(z_1, z_2) = z_1/z_2$ on Δ_H , we get $\|T_1T_2^{-1}\| \leq 1$, or equivalently, $(T_1T_2^{-1})^*(T_1T_2^{-1}) \leq I$. Thus, $T_1^*T_1 \leq T_2^*T_2$ completing the proof. \square

Remark 2.6. Assume that (8) holds and T_2 on \mathcal{H} is invertible. Note that 2-tuple $(T_1T_2^{-1}, T_2)$ is a commuting pair of contractions. By Ando's dilation theorem (see [3]), there exists a unitary commuting 2-tuple $U = (U_1, U_2)$ on a Hilbert space $\mathcal{K} \supseteq \mathcal{H}$ such that

$$T_1^m T_2^{n-m} = (T_1 T_2^{-1})^m T_2^n = P_{\mathcal{H}} U_1^m U_2^n, \quad m, n \geq 0,$$

where $P_{\mathcal{H}}$ denotes the orthogonal projection of \mathcal{K} onto \mathcal{H} . Finally, it has been pointed out by the anonymous referee that (8) can also be deduced from a couple of applications of the Gelfand spectral radius formula.

For certain Taylor invertible 2-tuples, the inclusion $\sigma(T) \subset \Delta_H$ appearing in Theorem 1.2 may be replaced by the weaker condition $\sigma(T) \subseteq \Delta_H$.

Corollary 2.7. *Let $T = (T_1, T_2)$ be a commuting 2-tuple on \mathcal{H} such that $\sigma(T) \subseteq \Delta_H$ and*

$$E_T := T_2^*T_2 - T_1^*T_1 - T_2^*(T_2^*T_2 - T_1^*T_1)T_2 \geq 0.$$

*If T is Taylor-invertible and $T_1^*T_1 \leq T_2^*T_2 \leq I$, then for every bounded rational function ϕ with poles off Δ_H , $\|\phi(T)\| \leq \|\phi\|_{\infty, \Delta_H}$.*

Proof. Assume that T is Taylor-invertible and $T_1^*T_1 \leq T_2^*T_2 \leq I$. For $0 < s, t < 1$, let $T_{s,t}$ denote the commuting pair (sT_1, tT_2) . Note that

$$E_{T_{s,t}} = t^2(E_T + (1 - t^2)T_2^*T_2 + (1 - s^2)T_1^*T_1 + (s^2t^2 - 1)T_2^*T_1^*T_1T_2).$$

Thus, if $T_1^*T_1 \leq T_2^*T_2 \leq I$, then

$$\begin{aligned} E_{T_{s,t}} &\geq t^2(E_T) + t^2(1 - t^2 + 1 - s^2 + s^2t^2 - 1)T_2^*T_1^*T_1T_2 \\ &= t^2(E_T) + t^2(1 - t^2)(1 - s^2)T_2^*T_1^*T_1T_2, \end{aligned}$$

which is a positive operator by assumption. Since $(0, 0) \notin \sigma(T)$, by the spectral mapping property (see [5, Theorem 5.19]), $\sigma(T_{s,t}) \subset \Delta_H$. One may now apply Theorem 1.2 to the commuting 2-tuple $T_{s,t}$ and any bounded rational function ϕ with poles off Δ_H , and let s, t tend to 1. \square

We do not know whether the assumption of the Taylor invertibility of T can be relaxed from Corollary 2.7. This constraint may be related to the fact that the origin is a singularity of the boundary of the Hartogs triangle.

References

- [1] AGLER, JIM; MCCARTHY, JOHN E.; YOUNG, NICHOLAS J. Operator analysis. Hilbert space methods in complex analysis. Cambridge Tracts in Mathematics, 219. *Cambridge University Press, Cambridge*, 2020. xv+375 pp. ISBN: 978-1-108-48544-9. [MR4411370](#), [Zbl 07162756](#), doi: [10.1017/9781108751292](#). [791](#), [793](#), [794](#), [795](#), [796](#)
- [2] AGLER, JIM; YOUNG, NICHOLAS J. The magic functions and automorphisms of a domain. *Complex Anal. Oper. Theory* **2** (2008), no. 3, 383–404. [MR2434458](#), [Zbl 1195.32013](#), [arXiv:0709.0096](#), doi: [10.1007/s11785-007-0039-5](#). [794](#)
- [3] ANDÔ, TSUYOSHI. On a pair of commutative contractions. *Acta Sci. Math.(Szeged)* **24** (1963), 88–90. [MR0155193](#), [Zbl 0116.32403](#). [794](#), [797](#)
- [4] ATHAVALE, AMEER. Model theory on the unit ball in \mathbf{C}^m . *J. Operator Theory* **27** (1992), no. 2, 347–358. [MR1249651](#), [Zbl 0804.47005](#). [795](#)
- [5] CURTO, RAÚL E. Applications of several complex variables to multiparameter spectral theory. Surveys of some recent results in operator theory, Vol. II, 25-90. Pitman Res. Notes Math. Ser., 192. *Longman Sci. Tech., Harlow*, 1988. [MR0976843](#), [Zbl 0827.47005](#). [793](#), [795](#), [797](#), [798](#)
- [6] CURTO, RAÚL E.; VASILESCU, FLORIAN-HORIA. Standard operator models in the polydisc. *Indiana Univ. Math. J.* **42** (1993), no. 3, 791-810. [MR1254118](#), [Zbl 0805.47007](#), doi: [10.1512/iumj.1993.42.42035](#). [794](#)
- [7] DINEEN, SEÁN. Complex analysis in locally convex spaces. Notas de Matemática [Mathematical Notes], 83. North-Holland Mathematics Studies, 57. *North-Holland Publishing Co., Amsterdam-New York*, 1981. xiii+492 pp. ISBN: 0-444-86319-2. [MR640093](#), [Zbl 0484.46044](#). [795](#)
- [8] DRURY, STEPHEN W. A generalization of von Neumann's inequality to the complex ball. *Proc. Amer. Math. Soc.* **68** (1978), no. 3, 300–304. [MR480362](#), [Zbl 0377.47016](#), doi: [10.2307/2043109](#). [794](#)
- [9] GALLAGHER, ANNE-KATRIN; GUPTA, PURVI; LANZANI, LOREDANA; VIVAS, LIZ R. Hardy spaces for a class of singular domains. *Math. Z.* **299** (2021), no. 3-4, 2171–2197. [MR4329283](#), [Zbl 07428892](#), [arXiv:2009.02466](#), doi: [10.1007/s00209-021-02755-1](#). [792](#)
- [10] HOGBE-NLEND, HENRI; MOSCATELLI, VINCENZO B. Nuclear and conuclear spaces. Introductory course on nuclear and conuclear spaces in the light of the duality "topology-bornology". Notas de Matemática [Mathematical Notes], 79. North-Holland Mathematics Studies, 52. *North-Holland Publishing Co., Amsterdam-New York*, 1981. x+275 pp. ISBN: 0-444-86207-2. [MR639899](#), [Zbl 0467.46001](#). [795](#)
- [11] JARNICKI, MAREK; PFLUG, PETER. First steps in several complex variables: Reinhardt domains. EMS Textbooks in Mathematics. *European Mathematical Society (EMS), Zürich*,

2008. viii+359 pp. ISBN: 978-3-03719-049-4. [MR2396710](#), [Zbl 1148.32001](#), doi: [10.4171/049.795](#)
- [12] MONGUZZI, ALESSANDRO. Holomorphic function spaces on the Hartogs triangle. *Math. Nachr.* **294** (2021), no. 11, 2209–2231. [MR4371295](#), [arXiv:1910.13741](#), doi: [10.1002/mana.201900477](#). [792](#)
- [13] MÜLLER, VLADIMÏR; VASILESCU, FLORIAN-HORA. Standard models for some commuting multioperators. *Proc. Amer. Math. Soc.* **117** (1993), no. 4, 979–989. [MR1112498](#), [Zbl 0777.47009](#), doi: [10.2307/2159525](#). [794](#)
- [14] PAULSEN, VERN I.; RAGHUPATHI, MRINAL. An introduction to the theory of reproducing kernel Hilbert spaces. Cambridge Studies in Advanced Mathematics, 152. *Cambridge University Press, Cambridge*, 2016. x+182 pp. ISBN: 978-1-107-10409-9. [MR3526117](#), [Zbl 1364.46004](#), doi: [10.1017/CBO9781316219232](#). [792](#), [796](#)
- [15] PIETSCH, ALBRECHT. Nuclear locally convex spaces. Translated from the second German edition by William H. Ruckle. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 66. Springer-Verlag, New York-Heidelberg*, 1972. ix+193 pp. [MR0350360](#), [Zbl 0236.46002](#), [795](#)
- [16] POTT, SANDRA. Standard models under polynomial positivity conditions. *J. Operator Theory* **41** (1999), no. 2, 365–389. [MR1681579](#), [Zbl 0995.47003](#). [793](#), [794](#)
- [17] PUTINAR, MIHAI. Spectral inclusion for subnormal n -tuples. *Proc. Amer. Math. Soc.* **90** (1984), no. 3, 405–406. [MR728357](#), doi: [10.2307/2044482](#). [793](#)
- [18] RUDIN, WALTER. Principles of mathematical analysis. Third edition. International Series in Pure and Applied Mathematics. *McGraw-Hill Book Co., New York-Auckland-Düsseldorf*, 1976. x+342 pp. [MR0385023](#), [Zbl 0346.26002](#). [796](#)
- [19] SCHEIDEMANN, VOLKER. Introduction to complex analysis in several variables. *Birkhäuser Verlag, Basel*, 2005. viii+171 pp. ISBN: 978-3-7643-7490-7; 3-7643-7490-X 32-01. [MR2176976](#), [Zbl 1085.32001](#). doi: [10.1007/3-7643-7491-8](#). [791](#)
- [20] SHAW, MEI-CHI. The Hartogs triangle in complex analysis. *Geometry and topology of submanifolds and currents*, 105–115. *Contemp. Math.*, 646. *Amer. Math. Soc., Providence, RI*, 2015. [MR3384976](#), [Zbl 1348.32001](#), doi: [10.1090/conm/646/12944](#). [791](#), [792](#), [795](#)
- [21] SZ.-NAGY, BÉLA; FOIAS, CIPRIAN; BERCOVICI, HARI; KÉRCHY, LÁSZLÓ. Harmonic analysis of operators on Hilbert space. Second edition. *Universitext. Springer, New York*, 2010. xiv+474 pp. ISBN: 978-1-4419-6093-1. [MR2760647](#), [Zbl 1234.47001](#), doi: [10.1002/zamm.19720520821](#). [794](#)
- [22] TSIKALAS, GEORGIOS. A von Neumann type inequality for an annulus. *J. Math. Anal. Appl.* **506** (2022), no. 2, Paper No. 125714, 12 pp. [MR4321662](#), [Zbl 07412879](#), [arXiv:2106.06013](#), doi: [10.1016/j.jmaa.2021.125714](#). [793](#), [794](#)

(Sameer Chavan) DEPARTMENT OF MATHEMATICS AND STATISTICS, INDIAN INSTITUTE OF TECHNOLOGY KANPUR, INDIA
chavan@iitk.ac.in

(Shubham Jain) DEPARTMENT OF MATHEMATICS AND STATISTICS, INDIAN INSTITUTE OF TECHNOLOGY KANPUR, INDIA
shubjain@iitk.ac.in

(Paramita Pramanick) SCHOOL OF MATHEMATICS, HARISH-CHANDRA RESEARCH INSTITUTE, CHHATNAG ROAD, JHUNSI, ALLAHABAD 211019, INDIA
paramitapramanick@hri.res.in

This paper is available via <http://nyjm.albany.edu/j/2022/28-31.html>.