

Slice filtration of certain C_{pq} -spectra

Surojit Ghosh

ABSTRACT. For p, q distinct odd primes and a virtual C_{pq} -representation α , we compute the slices of the C_{pq} -spectrum $S^\alpha \wedge H\underline{\mathbb{Z}}$, and prove the existence of *spherical slice* for this spectrum.

CONTENTS

| | |
|--|------|
| 1. Introduction | 1399 |
| Organization. | 1400 |
| Acknowledgement | 1400 |
| 2. Preliminaries | 1400 |
| 3. Some important Mackey functors | 1403 |
| 4. Spherical C_p -slices for the spectrum $S^W \wedge H\underline{\mathbb{Z}}$ | 1405 |
| 5. C_{pq} -slices for $S^V \wedge H\underline{\mathbb{Z}}$ | 1410 |
| References | 1417 |

1. Introduction

For a fixed finite group G , a central computational tool in G -equivariant stable homotopy theory is the $RO(G)$ -graded homotopy groups of a G -spectrum, but there are difficult to compute.

It was first observed by Voevodsky ([17]) that the stable motivic homotopy theory can be understood by their slice filtration. This slice filtration yields a spectral sequence, a tool computing various cohomology theories in algebraic geometry context. In equivariant homotopy theory, for any G -spectrum X , we have an analogous equivariant *slice tower*

$$\dots P^{n+1}X \rightarrow P^nX \rightarrow P^{n-1}X \rightarrow \dots \quad (1)$$

In this tower, P^nX is the localization with respect to the localizing category of G -spectra generated by $\Sigma_G^\infty G_+ \wedge_H S^{k\rho_H}$ where ρ_H is the regular representation of $H \leq G$ and $k|H| \geq n$. The homotopy fiber P_n^nX of $P^nX \rightarrow P^{n-1}X$ is called the n -*slice* of X . An n -slice is called *spherical* if it is of the form $S^V \wedge H\underline{\mathbb{Z}}$, where V is a (non-virtual) representation of G of dimension n .

Received August 2, 2021.

2010 *Mathematics Subject Classification*. Primary: 55N91, 55P91; Secondary: 57S17, 14M15.

Key words and phrases. Bredon cohomology, Mackey functor, Slice tower.

The homotopy spectral sequence of (1) is an essential tool in modern homotopy theory due to Hill-Hopkins-Ravenel in [10]. This is called the *slice spectral sequence* and it eventually computes the $RO(G)$ -graded homotopy groups of spectra.

The homotopy groups of the C_{2^n} -spectrum $N_2^{2^n}MU_{\mathbb{R}}$ played a crucial role in solving the Kervaire invariant one problem. In [10, Theorem 6.5], Hill, Hopkins, and Ravenel proved that the 0-th slice of this spectrum is the Eilenberg-Mac Lane spectrum $H\mathbb{Z}$. This yields that in the slice spectral sequence of any $N_2^{2^n}MU_{\mathbb{R}}$ -module, all the differentials and extensions live in $H\mathbb{Z}$ -modules. This motivates us to compute the slice filtration of the certain $H\mathbb{Z}$ -modules. The computation of the slice tower for $S^n \wedge H\mathbb{Z}$ was made in [20] and for $S^{n\xi} \wedge H\mathbb{Z}$ in [11] for the group C_{p^n} . Recently, in [8], Guillou and Yarnall computed the $C_2 \times C_2$ -slice tower for the spectrum $S^n \wedge H\mathbb{F}_2$.

In this article, for a C_{pq} -representation V , we investigate the spectrum $S^V \wedge H\mathbb{Z}$ and provide a description of the slices of it as follows:

Theorem A. *For any d -dimensional C_{pq} -representation V , the d -slice of the C_{pq} -spectrum $S^V \wedge H\mathbb{Z}$ is spherical.*

See Theorem 5.7.

As a direct consequence, we derive:

Corollary B. *For $\alpha \in RO(C_{pq})$ there exists $\beta \in RO(C_{pq})$ such that $S^\alpha \wedge H\mathbb{Z}$ has a $\dim(\alpha)$ -slice $S^\beta \wedge H\mathbb{Z}$. The other slices are suspensions of $H\mathcal{K}_p\langle\mathbb{Z}/p\rangle$ or $H\mathcal{K}_q\langle\mathbb{Z}/q\rangle$ or wedges of them.*

See Corollary 5.8.

Organization. Section 2 provides a short tour of equivariant homotopy theory. A few useful Mackey functors both in the context of C_p and C_{pq} have been described, and as a consequence, we prove certain equivalences of spectra in Section 3. Section 4 is dedicated to explaining the slice filtration of the C_p -spectrum $S^W \wedge H\mathbb{Z}$. Finally, in Section 5, we provide a careful analysis of the C_{pq} -spectrum $S^V \wedge H\mathbb{Z}$ with a result concerning the existence of a *spherical slice*.

Notation 1.1. *For an orthogonal G -representation V , $S(V)$ denotes the unit sphere, $D(V)$ the unit disk, and S^V the one-point compactification $\cong D(V)/S(V)$.*

Acknowledgement. The author wants to thank Samik Basu for suggesting the problem and David Blanc for a careful reading of the first draft of this work, which was partially supported by Israel Science Foundation grant 770/16.

2. Preliminaries

In this section, we recall specific fundamental ideas and techniques in Bredon cohomology. Throughout the paper, G will be a finite group.

Definition 2.1. The *Burnside category* of G is the category Burn_G of finite G -sets, with $\text{Hom}(S, T)$ the group completion of the set of correspondences $S \leftarrow U \rightarrow T$.

A functor $\underline{M} : \text{Burn}_G^{\text{op}} \rightarrow \text{Ab}$ from the Burnside category into abelian groups is called *Mackey functor*.

Explicitly, a G -Mackey functor \underline{M} is a collection of commutative $W_G(H)$ -groups $\underline{M}(G/H)$ one for each subgroup $H \leq G$, each accompanied by *transfer* $\text{tr}_K^H : \underline{M}(G/K) \rightarrow \underline{M}(G/H)$ and *restriction* $\text{res}_K^H : \underline{M}(G/H) \rightarrow \underline{M}(G/K)$ for $K \leq H \leq G$ such that

- (1) $\text{tr}_J^H = \text{tr}_K^H \text{tr}_J^K$ and $\text{res}_J^H = \text{res}_J^K \text{res}_K^H$ for all $J \leq K \leq H$.
- (2) $\text{tr}_K^H(\gamma \cdot x) = \text{tr}_K^H(x)$ for all $x \in \underline{M}(G/K)$ and $\gamma \in W_H(K)$.
- (3) $\gamma \cdot \text{res}_K^H(x) = \text{res}_K^H(x)$ for all $x \in \underline{M}(G/H)$ and $\gamma \in W_H(K)$.
- (4) $\text{res}_K^H \text{tr}_K^J(x) = \sum_{\gamma \in W_H(K)} \gamma \cdot \text{tr}_{J \cap K}^K(x)$ for all subgroups $J, H \leq K$.

Usually, we describe a Mackey functor by a *Lewis diagram* (see [13]). For a C_p -Mackey functor \underline{N} , we shall use the diagram:

$$\underline{N} : \begin{array}{ccc} & \underline{N}(C_p/C_p) & \\ & \text{res}_e^{C_p} \downarrow \uparrow \text{tr}_e^{C_p} & \\ & \underline{N}(C_p/e) & \end{array}$$

and a C_{pq} -Mackey functor \underline{M} , we shall associate the following diagram.

$$\underline{M} : \begin{array}{ccccc} & & \underline{M}(C_{pq}/C_{pq}) & & \\ & \text{res}_{C_p}^{C_{pq}} \swarrow & & \searrow \text{res}_{C_q}^{C_{pq}} & \\ & \text{tr}_{C_p}^{C_{pq}} \swarrow & & \searrow \text{tr}_{C_q}^{C_{pq}} & \\ \underline{M}(C_{pq}/C_p) & & & & \underline{M}(C_{pq}/C_q) \\ & \text{tr}_e^{C_p} \swarrow & & \searrow \text{tr}_e^{C_q} & \\ & \text{res}_e^{C_p} \swarrow & \underline{M}(C_{pq}/e) & \searrow \text{res}_e^{C_q} & \end{array}$$

We denote the category of G -Mackey functors by Mack_G .

Example 2.2. For an abelian group C , the *constant Mackey functor* \underline{C} is given by the assignment $\underline{C}(G/H) = C$ with $\text{res}_K^H = \text{Id}_C$ and $\text{tr}_K^H =$ multiplication by the index $[H : K]$. One can define the *dual constant Mackey functor* \underline{C}^* by interchanging the role of tr_K^H and res_K^H for $K \leq H \leq G$. In particular, for $G = C_p$, cyclic group of order p , the constant and dual constant C_p -Mackey functors are listed as follows.

$$\underline{\mathbb{Z}} : \quad \begin{array}{c} \mathbb{Z} \\ \uparrow \\ 1 \quad \downarrow \\ \mathbb{Z} \end{array} \quad \underline{\mathbb{Z}}^* : \quad \begin{array}{c} \mathbb{Z} \\ \uparrow \\ p \quad \downarrow \\ \mathbb{Z} \end{array}$$

Let Sp^G be the category of orthogonal G -spectra ([14]). It has a symmetric model category structure $(\text{Sp}^G, \wedge, \mathbb{S}^0)$ as in [10, Appendix A, B], and we use Sp^G as our model for the G -equivariant stable homotopy theory.

It seems that the equivariant homotopy groups are more naturally graded on $RO(G)$, the free abelian group generated by the irreducible representations of G . For any $\alpha = V - W \in RO(G)$, we define

$$\pi_\alpha(X) := [S^V, S^W \wedge X]^G,$$

where the right hand side denotes the set of homotopy classes of maps in Sp^G . Thus, any orthogonal G -spectrum X has $RO(G)$ -graded homotopy groups, denoted by $\pi_\star(X)$. It induces a Mackey functor $\underline{\pi}_\alpha(X)$ such that

$$\underline{\pi}_\alpha(X)(G/H) := \pi_\alpha(\Sigma_G^\infty G/H_+ \wedge X).$$

For any G -Mackey functor \underline{M} there is an equivariant Eilenberg-Mac Lane spectrum \underline{HM} [7, Theorem 5.3] with

$$\underline{\pi}_k(\underline{HM}) = \begin{cases} \underline{M} & \text{for } k = 0, \\ 0 & \text{otherwise.} \end{cases}$$

The equivariant Eilenberg-Mac Lane spectrum \underline{HM} gives rise to a reduced cohomology theory given by

$$\hat{H}_G^\alpha(X; \underline{M}) \cong [X, \Sigma^\alpha \underline{HM}]^G.$$

This cohomology is called the $RO(G)$ -graded Bredon cohomology with coefficients in the Mackey functor \underline{M} .

It is interesting to note that the suspension isomorphism for $RO(G)$ -graded cohomology theory $\hat{H}_G^\alpha(-; \underline{M})$ takes the form $\hat{H}_G^\alpha(X; \underline{M}) \cong \hat{H}_G^{\alpha+V}(S^V \wedge X; \underline{M})$ for every based G -space X and representation V .

Notation 2.3. For each odd cyclic group C_n , the representation ring $RO(C_n)$ is generated by the trivial representation 1, and the 2-dimension representation ξ^j given by the rotation by the angle $\frac{2\pi j}{n}$ for $j = 1, \dots, \frac{n-1}{2}$.

Proposition 2.4 ([22], Proposition 4.25). Let p be an odd prime. In Sp^{C_p} , there is an equivalence $H\underline{\mathbb{Z}} \wedge S^\xi \simeq H\underline{\mathbb{Z}} \wedge S^{\xi^j}$ whenever $p \nmid j$.

We recall that there are change of groups functors on equivariant spectra. The restriction functor $i_H^* : \text{Sp}^G \rightarrow \text{Sp}^H$ from G -spectra to H -spectra has a left

adjoint given by smashing with $G_+ \wedge_H (-)$. For a G -CW complex X , this also induces an isomorphism for cohomology with Mackey functor coefficients

$$\tilde{H}_G^\alpha(G_+ \wedge_H X; \underline{M}) \cong \tilde{H}_H^\alpha(i_H^* X; \downarrow_H^G(\underline{M})).$$

Here $\downarrow_H^G(\underline{M})$ is the H -Mackey functor defined by $\downarrow_H^G(\underline{M})(H/L) := \underline{M}(G/H \times H/L)$.

The $RO(G)$ -graded cohomology theories may also be assumed to be Mackey functor-valued as in the definition below.

Definition 2.5. Let X be a pointed G -space, \underline{M} be any G -Mackey functor, and $\alpha \in RO(G)$. Then the Mackey functor valued cohomology $\underline{H}_G^\alpha(X; \underline{M})$ ([16, §2.3]) is defined:

$$\underline{H}_G^\alpha(X; \underline{M})(G/K) = \tilde{H}_G^\alpha((G/K)_+ \wedge X; \underline{M}).$$

The restriction and transfer maps are induced by the appropriate maps of G -spectra.

3. Some important Mackey functors

We start by recalling the C_p -Mackey functors from Example 2.2 along with the following.

$$\langle \mathbb{Z}/p \rangle : \begin{array}{c} \mathbb{Z}/p \\ \left(\begin{array}{c} \uparrow \\ \downarrow \end{array} \right) \\ 0 \end{array}$$

Now observe that the Burnside category $\text{Burn}_{C_{pq}}$ is isomorphic to $\text{Burn}_{C_p} \otimes \text{Burn}_{C_q}$ formed as the product set of objects and tensor product set of morphisms. Thus, we may define a C_{pq} -Mackey functor by tensoring Mackey functors on C_p and C_q . This is defined by a functor $\boxtimes : \text{Mack}_{C_p} \times \text{Mack}_{C_q} \rightarrow \text{Mack}_{C_{pq}}$ denoted by $(\underline{M}, \underline{N}) \mapsto \underline{M} \boxtimes \underline{N}$ such that on objects $G/H = C_p/H_1 \times C_q/H_2$,

$$\underline{M} \boxtimes \underline{N}(G/H) = \underline{M}(C_p/H_1) \otimes \underline{N}(C_q/H_2).$$

The restrictions and transfers are given by tensoring the restrictions and transfers for \underline{M} and \underline{N} (§3, [1]). The following Mackey functors have particular importance in our case.

Definition 3.1. For a C_p -Mackey functor \underline{M} , define C_{pq} -Mackey functors

$$\mathcal{C}_p \underline{M} := \underline{M} \boxtimes \underline{\mathbb{Z}}^*, \quad \mathcal{K}_p \underline{M} := \underline{M} \boxtimes \underline{\mathbb{Z}}.$$

The Lewis diagrams for the Mackey functors $\mathcal{K}_p \langle \mathbb{Z}/p \rangle$ and $\mathcal{K}_q \langle \mathbb{Z}/q \rangle$ are

$$\begin{array}{ccc} \begin{array}{c} \mathbb{Z}/p \\ \uparrow \quad \downarrow \\ \mathbb{Z}/p \quad 0 \\ \downarrow \quad \uparrow \\ 0 \end{array} & \begin{array}{c} \mathbb{Z}/q \\ \uparrow \quad \downarrow \\ 0 \quad \mathbb{Z}/q \\ \downarrow \quad \uparrow \\ 0 \end{array} & (2) \end{array}$$

$$\mathcal{K}_p\langle \mathbb{Z}/p \rangle \qquad \mathcal{K}_q\langle \mathbb{Z}/q \rangle$$

Also we denote the tensor products

$$\underline{\mathbb{Z}} \boxtimes \underline{\mathbb{Z}} \text{ and } \underline{\mathbb{Z}}^* \boxtimes \underline{\mathbb{Z}}^*$$

again by $\underline{\mathbb{Z}}$ and $\underline{\mathbb{Z}}^*$ respectively, and these are the constant and dual constant Mackey functors for the group C_{pq} . Recall [18, §7] that

Definition 3.2. A G -Mackey functor \underline{M} is called *cohomological Mackey functor* if, for $K \leq H \leq G$, the composition $\text{tr}_K^H \text{res}_K^H : \underline{M}(G/H) \rightarrow \underline{M}(G/H)$ of the restriction $\text{res}_K^H : \underline{M}(G/H) \rightarrow \underline{M}(G/K)$ with the transfer $\text{tr}_K^H : \underline{M}(G/K) \rightarrow \underline{M}(G/H)$ is equal to the multiplication by index $[H : K]$.

Any module over the constant Mackey functor $\underline{\mathbb{Z}}$ is a cohomological Mackey functor (see [21]). For the group $G = C_{pq}$, any cohomological Mackey functor satisfies the following:

Proposition 3.3. *Let \underline{M} be a cohomological C_{pq} -Mackey functor such that both the abelian groups $\underline{M}(C_{pq}/C_p)$ and $\underline{M}(C_{pq}/C_q)$ are trivial. Then, $\underline{M}(C_{pq}/C_{pq})$ is the trivial abelian group.*

Proof. Since \underline{M} is cohomological, for $K \leq H \leq G$, the composition

$$\text{tr}_K^H \text{res}_K^H : \underline{M}(G/H) \rightarrow \underline{M}(G/H)$$

is given by the multiplication by index $[H : K]$. Let $x \in \underline{M}(C_{pq}/C_{pq})$; applying above map we get

$$px = 0 \text{ and } qx = 0.$$

Since, p and q are relatively prime, $x = 0$. The result follows. □

With the above notations, we recall from [1] the additive structure of the Mackey functor $H_{C_{pq}}^\alpha(S^0; \underline{\mathbb{Z}})$ as follows:

Theorem 3.4 (Theorem 7.3, [1]). *Let $\alpha \in RO(C_{pq})$. Then the Mackey functor*

$$H_{C_{pq}}^\alpha(S^0; \underline{\mathbb{Z}}) \cong \begin{cases} \mathcal{K}_p\langle \mathbb{Z}/p \rangle & \text{if } |\alpha| < 0, |\alpha^{C_p}| > 1, |\alpha^{C_q}| \leq 1 \text{ odd} \\ \mathcal{K}_q\langle \mathbb{Z}/q \rangle & \text{if } |\alpha| < 0, |\alpha^{C_p}| \leq 1, |\alpha^{C_q}| > 1 \text{ odd} \\ \mathcal{K}_p\langle \mathbb{Z}/p \rangle \oplus \mathcal{K}_q\langle \mathbb{Z}/q \rangle & \text{if } |\alpha| < 0, |\alpha^{C_p}| > 1, |\alpha^{C_q}| > 1 \text{ odd} \\ \mathcal{K}_p\langle \mathbb{Z}/p \rangle \oplus \mathcal{K}_q\langle \mathbb{Z}/q \rangle & \text{if } |\alpha| > 0, |\alpha^{C_p}| \leq 0, |\alpha^{C_q}| \leq 0 \text{ even} \\ \mathcal{K}_p\langle \mathbb{Z}/p \rangle & \text{if } |\alpha| > 0, |\alpha^{C_p}| \leq 0, |\alpha^{C_q}| > 0 \text{ even} \\ \mathcal{K}_q\langle \mathbb{Z}/q \rangle & \text{if } |\alpha| > 0, |\alpha^{C_p}| > 0, |\alpha^{C_q}| \leq 0 \text{ even} \\ \underline{\mathbb{Z}} & \text{if } |\alpha| = 0, |\alpha^{C_p}| \leq 0, |\alpha^{C_q}| \leq 0 \\ \underline{\mathbb{Z}}^* & \text{if } |\alpha| = 0, |\alpha^{C_p}| > 0, |\alpha^{C_q}| > 0 \\ \mathcal{K}_p\underline{\mathbb{Z}}^* & \text{if } |\alpha| = 0, |\alpha^{C_p}| > 0, |\alpha^{C_q}| \leq 0 \\ \mathcal{K}_q\underline{\mathbb{Z}}^* & \text{if } |\alpha| = 0, |\alpha^{C_p}| \leq 0, |\alpha^{C_q}| > 0 \\ 0 & \text{otherwise.} \end{cases}$$

Using Theorem 3.4, we have the following equivalences of spectra which simplify computations:

Lemma 3.5. (a) *There are the equivalences*

- (1) $S^\xi \wedge H\underline{\mathbb{Z}} \simeq S^{\xi^j} \wedge H\underline{\mathbb{Z}}$ if $(j, pq) = 1$.
- (2) $S^{\xi^p} \wedge H\underline{\mathbb{Z}} \simeq S^{\xi^{jp}} \wedge H\underline{\mathbb{Z}}$ if $(j, q) = 1$.
- (3) $S^{\xi^q} \wedge H\underline{\mathbb{Z}} \simeq S^{\xi^{jq}} \wedge H\underline{\mathbb{Z}}$ if $(j, p) = 1$.
- (4) $\Sigma^{\xi^p} H\underline{\mathbb{Z}} \simeq \Sigma^\xi H(\mathcal{K}_p \underline{\mathbb{Z}}^*)$.
- (5) $H\underline{\mathbb{Z}}^* \simeq \Sigma^{2-\xi} H\underline{\mathbb{Z}}$.

(b) *The equivalence (4) induces a map $u_{\xi-\xi^p} : H\underline{\mathbb{Z}} \rightarrow \Sigma^{\xi-\xi^p} H\underline{\mathbb{Z}}$ and equivalence (5) induces $u_\xi : H\underline{\mathbb{Z}} \rightarrow \Sigma^{\xi-2} H\underline{\mathbb{Z}}$.*

Proof. (a) These follow by computing homotopy groups of $S^{\xi^j-\xi^k} \wedge H\underline{\mathbb{Z}}$ and using Theorem 3.4.

(b) Consider the C_p -Mackey functor map $\underline{\mathbb{Z}}^* \rightarrow \underline{\mathbb{Z}}$ which is the identity on the orbit C_p/e . By tensoring with the constant C_q -Mackey functor $\underline{\mathbb{Z}}$ and using equivalence (4), we obtain

$$H\underline{\mathbb{Z}} \xrightarrow{\cong} \Sigma^{\xi-\xi^p} H\mathcal{K}_p \underline{\mathbb{Z}}^* \rightarrow \Sigma^{\xi-\xi^p} H\underline{\mathbb{Z}}.$$

Next, consider the canonical C_{pq} -Mackey functor map $\underline{\mathbb{Z}}^* \rightarrow \underline{\mathbb{Z}}$ which is the identity at level C_{pq}/e . Applying (5) yields u_ξ . \square

Remark 3.6. As a consequence of Lemma 3.5, only the C_{pq} -representations of the form

$$V = a + b\xi + c\xi^p + d\xi^q \text{ for } a, b, c, d \in \mathbb{Z}$$

are useful for our computations.

4. Spherical C_p -slices for the spectrum $S^W \wedge H\underline{\mathbb{Z}}$

In this section, we recall certain general facts about the slice tower of any genuine G -spectra. We also prove a few results for C_p -spectra of the form $S^W \wedge H\underline{\mathbb{Z}}$.

Recall the *localizing subcategories* of Sp^G are those closed under weak equivalences, cofibrations, extensions, coproducts, and well-ordered homotopy colimits ([15, §2]). For each integer n , let $\tau_{\geq n}$ denote the localizing subcategory of Sp^G generated by the G -spectra of the form $\Sigma_G^\infty G_+ \wedge_H S^{k\rho_H}$, where H ranges over all subgroups of G , ρ_H is the regular representation of H and $k|H| \geq n$.

Associated to the category $\tau_{\geq n}$ there is a natural localization functor P^{n-1} . As $\tau_{\geq n+1} \subseteq \tau_{\geq n}$, there is a natural transformation $P^n \rightarrow P^{n-1}$, which yields the slice tower (1) for each spectrum X . The homotopy fibre of $P^n X \rightarrow P^{n-1} X$, denoted by $P_n^n X$, is called the *n-slice* for the spectrum X (see [15, 4]).

A spectrum $X \in \tau_{\geq n}$ is called *n-slice connective* written $X \geq n$. A criterion for this to hold is produced by

Theorem 4.1 (Theorem A, [12]). *A G-spectrum $X \geq n$ if and only if the non equivariant homotopy groups*

$$\pi_k \Phi^H(X) = 0$$

for all $H \leq G$ and $k < \frac{n}{|H|}$.

Here $\Phi^H(X)$ is the H -geometric fixed points of the spectrum X , that is, the geometric fixed point of $i_H^*(X)$, the restriction of the G -spectrum X to $H \leq G$.

Remark 4.2. For a fixed point free G -representation V , $a_V \in \pi_{-V}(S^0)$ is represented by the map $S^0 = \{0, \infty\} \subseteq S^V$. For any $X \in \text{Sp}^G$, $\Phi^G(X)$ is the fixed point spectrum of the localization $X[a_{\bar{\rho}_G}^{-1}]$ by the element $a_{\bar{\rho}_G}$. For $G = C_{pq}$, it is evident that $\xi^p \oplus \xi^q \subset \bar{\rho}_{C_{pq}}$. Therefore, $a_{\bar{\rho}_{C_{pq}}}$ is trivial by the ring structure described in [2]. Hence, the spectrum $\Phi^{C_{pq}}(X)$ has trivial homotopy groups.

The local objects in Sp^G with respect to $\tau_{\geq n}$ are those X such that $[Y, X]^G = 0$ for all $Y \in \tau_{\geq n}$. These spectra are called *n-slice coconnective* spectra, and we write $X \leq n$ in this case. Equivalently,

Lemma 4.3. *A spectrum is n-slice coconnective if the restriction $i_H^*(X) \leq n$ and $[S^{k\rho_G}, X]^G = 0$ for all $k \geq 0$ such that $k|G| > n$. We denote such an X by $X \leq n$.*

By [9, Proposition 2.6], if a spectrum X satisfies $k \leq X \leq n$, the same holds for all its restrictions. Moreover, $n \leq X \leq n$ if and only if $X = P_n^n X$ in which case X is called an *n-slice*.

In general, $\Sigma^V P_n^n X$ need not equal $P_{n+\dim(V)}^{n+\dim(V)}(\Sigma^V X)$, but it does commute with suspensions of the following forms.

Proposition 4.4 (Corollary 4.25, [10]). *If X an m-slice, $\Sigma^{k\rho_G} X$ is $(m+k|G|)$ -slice for all $k \in \mathbb{Z}$, that is,*

$$P_{m+k|G|}^{m+k|G|}(\Sigma^{k\rho_G} X) \simeq \Sigma^{k\rho_G} P_m^m(X).$$

The homotopy spectral sequence of the slice tower (1) of a G -spectrum X is called the *slice spectral sequence* for X , with

$$E_2^{s,t} = \underline{\pi}_{t-s}(P_t^t X) \Rightarrow \underline{\pi}_{t-s}(X)$$

(more generally, it computes the $RO(G)$ -graded homotopy groups $\underline{\pi}_\star(X)$). The associated filtration on the homotopy groups of the spectrum X is

$$F^s \underline{\pi}_t(X) = \text{Ker}(\underline{\pi}_t(X) \rightarrow \underline{\pi}_t(P^{t+s-1} X)).$$

In [15], Ullman proved following

Proposition 4.5 (Corollary 8.6, [15]). *If $n > 0$, then for a $(n + 1)$ -coconnective spectrum X ,*

$$F^s \underline{\pi}_{-n}(X) \cong \mathcal{F}^{(s+n-1)/n} \underline{\pi}_{-n} X.$$

The filtration $\mathcal{F}^k \underline{M}$ for a Mackey functor \underline{M} is given by:

$$\mathcal{F}^k \underline{M}(G/H) = \{x \in \underline{M}(G/H) : i_{|J|}^* x = 0, \text{ for all } J \subset H, |J| \leq k\}.$$

Where $i_a^* : \text{Mack}_G \rightarrow \text{Mack}_G$ such that

$$i_a^* \underline{M}(G/H) = \begin{cases} 0 & \text{if } |H| > a, \\ \underline{M}(G/H) & \text{otherwise,} \end{cases}$$

with restrictions and transfers induced from \underline{M} .

For the rest of this section, $G = C_p$, for p odd prime order. Let W be a (non-virtual) representation C_p of the form $m + n\xi$ with $m, n \geq 0$. To describe the $\dim(W)$ -slice of $S^W \wedge H\underline{\mathbb{Z}}$, we begin with the n -coconnective case:

Lemma 4.6. *For a C_p -representation $W = m + n\xi$, the C_p -spectrum $S^W \wedge H\underline{\mathbb{Z}}$ is slice $\dim(W)$ -coconnective if and only if*

$$m \leq \frac{2n + 3p}{p - 1}.$$

Proof. By Lemma 4.3, we need to find conditions on m, n such that

$$[S^{k\rho}, S^W \wedge H\underline{\mathbb{Z}}]^{C_p} = \tilde{H}_{C_p}^{W-k\rho}(S^0; \underline{\mathbb{Z}}) = 0$$

for all $k \geq 0$ with $k\rho > m + 2n$. These cohomology groups are computed in [5, Corollary B.10]. For completeness, recall that

$$\underline{H}_{C_p}^\alpha(S^0; \underline{\mathbb{Z}}) \cong \begin{cases} \underline{\mathbb{Z}} & \text{if } |\alpha| = 0, |\alpha^{C_p}| \leq 0 \\ \underline{\mathbb{Z}}^* & \text{if } |\alpha| = 0, |\alpha^{C_p}| > 0 \\ \langle \mathbb{Z}/p \rangle & \text{if } |\alpha| > 0, |\alpha^{C_p}| \leq 0 \text{ even} \\ \langle \mathbb{Z}/p \rangle & \text{if } |\alpha| < 0, |\alpha^{C_p}| \geq 3 \text{ odd} \\ 0 & \text{otherwise.} \end{cases} \quad (3)$$

Let $\alpha = W - k\rho$. Then $(|\alpha|, |\alpha^{C_p}|) = (m + 2n - k\rho, m - k)$. Therefore, since $m + 2n - k\rho < 0$, then by (3), the group $\tilde{H}_{C_p}^{W-k\rho}(S^0; \underline{\mathbb{Z}})$ is non-zero if and only if $m - k \geq 3$ odd, equivalently (by transforming the inequalities),

$$\frac{m + 2n}{p} < k \leq m - 3.$$

So there exists k with $0 \leq k$ and $k\rho > m + 2n$ such that $\tilde{H}_{C_p}^{W-k\rho}(S^0; \underline{\mathbb{Z}})$ is non-trivial if and only if $m - 3 > \frac{m+2n}{p}$. Therefore, $S^W \wedge H\underline{\mathbb{Z}} \leq \dim(W)$ if and only if $m \leq \frac{2n+3p}{p-1}$. The result follows. \square

Lemma 4.7. *Let $W = m + n\xi$ be a representation of C_p . Then the spectrum $S^W \wedge H\underline{\mathbb{Z}} \geq \dim(W)$ if and only if*

$$\frac{2n}{p-1} \leq m.$$

Proof. Write $\dim(W) = d$, the C_p -spectrum $S^W \wedge H\underline{\mathbb{Z}} \geq d$ if and only if $\pi_k \Phi^{C_p}(S^W \wedge H\underline{\mathbb{Z}}) = 0$ for all $k \leq d/p$, as the statement of Theorem 4.1 for $H = e$ is always true for $X = S^W \wedge H\underline{\mathbb{Z}}$. Since $\Phi^{C_p}(\Sigma^W \wedge H\underline{\mathbb{Z}}) \cong S^m \wedge H\underline{\mathbb{Z}}/p$, $\pi_k \Phi^{C_p}(\Sigma^W \wedge H\underline{\mathbb{Z}}) \neq 0$ if and only if $k = m$. Therefore, the statement is true if and only if $m \geq d/p = (m + 2n)/p$. \square

Lemma 4.6 and Lemma 4.7 together directly imply the following

Proposition 4.8. *For a C_p -representation W , the spectrum $S^W \wedge H\underline{\mathbb{Z}}$ is a $\dim(W)$ -slice if and only if*

$$\frac{\dim(W)}{p} \leq \dim(W^{C_p}) \leq \frac{\dim(W)}{p} + 3.$$

Remark 4.9. For $p = 3$, we have an alternative proof: Recall that the regular representation is $\rho_{C_3} = 1 + \xi$. Therefore, we have $S^{m+n\xi} \wedge H\underline{\mathbb{Z}} \cong S^{n\rho_{C_3}} \wedge \Sigma^{m-n} H\underline{\mathbb{Z}}$. So, the problem reduces to showing that $\Sigma^{m-n} H\underline{\mathbb{Z}}$ is an $(m - n)$ -slice. Clearly, $\Sigma^{m-n} H\underline{\mathbb{Z}} \geq m - n$. By [5, Corollary B.10], it follows that if $0 \leq m - n \leq 4$, then the spectrum $\Sigma^{m-n} H\underline{\mathbb{Z}} \leq m - n$.

For $n > 0$, recall [20, §3.1] d_n to be the number of the integers of the same parity as n that lie between $\frac{n}{p}$ and $n - 2$. It can be expressed using the following formula:

$$d_n := \frac{1}{2} \left(n - \frac{(n - n_0)}{p} - \delta \right) \tag{4}$$

where n_0 is the residue of n modulo p and

$$\delta = \begin{cases} 2 & \text{if } n_0 \text{ is even} \\ 1 & \text{if } n_0 \text{ is odd} \\ 0 & \text{if } n_0 = 0. \end{cases} \tag{5}$$

Corollary 4.10. *For a C_p -representation W , if the C_p -spectrum $S^W \wedge H\underline{\mathbb{Z}}$ is slice $\dim(W)$ -coconnective then the integer*

$$\frac{1}{2}(\dim(W) - \dim(W^{C_p})) - d_{\dim(W)} \geq -1.$$

The equality holds, if and only if $S^W \wedge H\underline{\mathbb{Z}}$ is a $\dim(W)$ -slice.

Proof. Using (4),

$$\begin{aligned} \frac{1}{2}(\dim(W) - \dim(W^{C_p})) - d_{\dim(W)} &= \frac{1}{2} \left(\frac{\dim(W)}{p} - \dim(W^{C_p}) - \frac{\dim(W)_0}{p} + \delta \right) \\ &\geq \frac{1}{2} \left(-3 - \frac{\dim(W)_0}{p} + \delta \right) \text{ (by Proposition 4.8)} \\ &\geq -\frac{3}{2} \text{ (by (5)).} \end{aligned}$$

If the equality holds, then

$$\frac{\dim(W)}{p} - \dim(W^{C_p}) = \frac{\dim(W)_0}{p} - 2 - \delta \leq 0 \text{ (by (5))}.$$

Then by Proposition 4.8, the spectrum $S^W \wedge H\underline{\mathbb{Z}}$ is a $\dim(W)$ -slice. The result follows. \square

For the cyclic group $G = C_{p^k}$, Yarnall ([20, Main Theorem, §4]) computed an explicit formula of the n -slice associated to the C_{p^k} -spectrum $S^n \wedge H\underline{\mathbb{Z}}$ when $n \geq 3$. To summarise her formula for $k = 1$, define two C_p -representations

$$W(n) = (n - 2d_n) + d_n\xi \text{ and } W'(n) = W(n) + (2 - \xi),$$

and then

$$P_n^n(S^n \wedge H\underline{\mathbb{Z}}) = \begin{cases} S^{W(n)} \wedge H\underline{\mathbb{Z}} & \text{if } p \nmid n, n_0 = \text{even} \\ S^{W'(n)} \wedge H\underline{\mathbb{Z}} & \text{if } p \nmid n, n_0 = \text{odd} \\ S^{W'(n)} \wedge H\underline{\mathbb{Z}} & \text{if } p \mid n. \end{cases} \quad (6)$$

Let W be a C_p -representation. The slice formula for the non-trivial $RO(C_p)$ -suspensions $S^W \wedge H\underline{\mathbb{Z}}$ of $H\underline{\mathbb{Z}}$ can be derived from the following result.

Theorem 4.11 (Theorem C, [12]). *Let p be odd. For the C_p -spectrum $X = S^W \wedge H\underline{\mathbb{Z}}$, we have equivalences for any $a \in \mathbb{Z}$*

$$P_{ap}^{ap}(X) \simeq S^{a\rho} \wedge H(\underline{H}_{C_p}^{W-a\rho}(S^0; \underline{\mathbb{Z}})).$$

$$P_{ap+2k+1}^{ap+2k+1}(X) \simeq S^{a\rho+k\xi+1} \wedge H(\mathcal{P}^0 \underline{H}_{C_p}^{W-a\rho-k\xi-1}(S^0; \underline{\mathbb{Z}})), \quad 0 \leq k \leq \frac{p-3}{2}.$$

$$P_{ap+2k+2}^{ap+2k+2}(X) \simeq S^{a\rho+(k+1)\xi} \wedge H(EC_p \otimes \underline{H}_{C_p}^{W-a\rho-(k+1)\xi}(S^0; \underline{\mathbb{Z}})), \quad 0 \leq k \leq \frac{p-3}{2}.$$

Here, for a G -Mackey functor \underline{M} , $EG \otimes \underline{M}$ denote the subMackey functor generated by $\underline{M}(G/e)$. \mathcal{P}^0 is the functor that takes a Mackey functor to the largest quotient in which the restriction maps are injections.

As a consequences of the above discussion one may easily derive the formula for $\dim(W)$ -slice for the spectrum $S^W \wedge H\underline{\mathbb{Z}}$ as follows.

Proposition 4.12. *Let W be a C_p -representation with dimension ω . Then*

(1) *If the spectrum $S^W \wedge H\underline{\mathbb{Z}}$ is slice ω -coconnective, then either $S^W \wedge H\underline{\mathbb{Z}}$ itself a ω -slice or $S^{W(\omega)} \wedge H\underline{\mathbb{Z}}$ is the ω -slice of $S^W \wedge H\underline{\mathbb{Z}}$.*

(2) *If the spectrum $S^W \wedge H\underline{\mathbb{Z}}$ is slice ω -connective, then either $S^W \wedge H\underline{\mathbb{Z}}$ itself a ω -slice or*

$$P_\omega^\omega(S^W \wedge H\underline{\mathbb{Z}}) = \begin{cases} S^{W(\omega)} \wedge H\underline{\mathbb{Z}} & \text{if } p \mid \omega \\ S^{W(\omega)} \wedge H\underline{\mathbb{Z}} & \text{if } p \nmid \omega \text{ and } \omega_0 = \text{odd} \\ S^{W(\omega)} \wedge H\underline{\mathbb{Z}} & \text{if } p \nmid \omega \text{ and } \omega_0 = \text{even}. \end{cases}$$

Proof. Here we prove part (1); the proof of part (2) is similar to (1). Set $X := S^W \wedge H\underline{\mathbb{Z}}$. For (1), first consider $p \mid \omega$, that is, $\omega = ap$ for some integer a . One readily computes

$$W(\omega) = a + \left(\frac{a(p-1)}{2}\right)\xi.$$

Since we consider X as slice ω -coconnective, thus by Lemma 4.10, one derives either $\dim(W^{C_p}) \leq a$ or X itself is a ω -slice. Then by Theorem 4.11, the ω -slice of X ,

$$P_\omega^\omega(X) \simeq S^{a\rho} \wedge H(\underline{H}_{C_p}^{W-a\rho}(S^0; \underline{\mathbb{Z}})) \cong S^{a\rho} \wedge H\underline{\mathbb{Z}} \cong S^{W(\omega)} \wedge H\underline{\mathbb{Z}} \text{ (by Proposition 2.4).}$$

If ω of the form $ap+2k+1$ where $0 \leq k \leq \frac{p-3}{2}$, then $W(\omega) = a+1+(\frac{a(p-1)}{2}+k)\xi$. If X is not a ω -slice, then Lemma 4.10 computes $\dim W^{C_p} \leq a + 1$. Thus by Theorem 4.11,

$$\begin{aligned} P_\omega^\omega(X) &\simeq S^{a\rho+k\xi+1} \wedge H\mathcal{P}^0(\underline{H}_{C_p}^{W-a\rho-k\xi-1}(S^0; \underline{\mathbb{Z}})) \\ &\cong S^{a\rho+k\xi+1} \wedge H\mathcal{P}^0(\underline{\mathbb{Z}}) \\ &= S^{a\rho+k\xi+1} \wedge H\underline{\mathbb{Z}} \\ &\cong S^{W(\omega)} \wedge H\underline{\mathbb{Z}} \text{ (by Proposition 2.4)} \end{aligned}$$

Finally, if ω is of the form $ap+2k+2$ for some $0 \leq k \leq \frac{p-3}{2}$, then we compute $W(\omega) = a + 2 + (\frac{a(p-1)}{2} + k)\xi$. The result follows by analogous computations above along with the observation $EC_p \otimes \underline{\mathbb{Z}}^* \cong \underline{\mathbb{Z}}^* \cong \sum^{\xi-2} H\underline{\mathbb{Z}}$. □

Example 4.13. (1) Let us consider the spectrum $X = S^{2+5\xi} \wedge H\underline{\mathbb{Z}}$. Using Proposition 4.8, one observes X is 12-coconnective but not slice 12-connective. Also note that $S^{W(12)} \wedge H\underline{\mathbb{Z}} \cong S^{4+4\xi} \wedge H\underline{\mathbb{Z}}$ is a 12-slice (by Theorem 4.12). (2) Next, consider $X = S^{8+2\xi} \wedge H\underline{\mathbb{Z}}$, which is slice 12-connective but not slice 12-coconnective. Here we compute $S^{W'(12)} \wedge H\underline{\mathbb{Z}} \cong S^{6+3\xi} \wedge H\underline{\mathbb{Z}}$, and it is the 12-slice of X by Theorem 4.12.

5. C_{pq} -slices for $S^V \wedge H\underline{\mathbb{Z}}$

In this section, we compute the C_{pq} -slices for the spectrum $S^\alpha \wedge H\underline{\mathbb{Z}}$ for each $\alpha \in RO(C_{pq})$. Slices of any C_{pq} -spectrum have a special feature: under a mild condition, the information of the C_p - and C_q -slices of the corresponding restrictions give the slices of C_{pq} -spectrum as follows.

Proposition 5.1. *Let X be a C_{pq} -spectrum such that $\underline{\pi}_\star(X)$ is cohomological. Then X is a k -slice if and only if both $i_{C_p}^*(X)$ and $i_{C_q}^*(X)$ are k -slices.*

Proof. Assume both $i_{C_p}^*(X)$ and $i_{C_q}^*(X)$ are k -slices. Then, by Proposition 3.3 X is a k -slice for the group C_{pq} . The other direction follows from [10, Proposition 4.13]. □

For each $k \geq 0$, the Mackey functor $\underline{\pi}_k S^V \wedge H\underline{\mathbb{Z}}$ is cohomological (see Definition 3.2). Therefore, as a direct consequence of Proposition 5.1, one may extend the detection result (Proposition 4.8) for C_p to C_{pq} case:

Corollary 5.2. *Let $p < q$ be odd primes and V be a C_{pq} -representation of the form $V = a + b\xi + c\xi^p + d\xi^q$. Then the spectrum $S^V \wedge H\underline{\mathbb{Z}}$ is a C_{pq} - $\dim(V)$ -slice if and only if*

$$\begin{aligned} \text{i)} \quad & \frac{2(b+d)}{p-1} \leq a + 2c \leq \frac{2(b+d)+3p}{p-1}. \\ \text{ii)} \quad & \frac{2(b+c)}{q-1} \leq a + 2d \leq \frac{2(b+c)+3q}{q-1}. \end{aligned}$$

Proof. If $S^V \wedge H\underline{\mathbb{Z}}$ is a slice, it must be a $\dim(V)$ -slice. Using Proposition 5.1, it is enough to show that both the spectra $i_{C_p}^*(S^V \wedge H\underline{\mathbb{Z}})$ and $i_{C_q}^*(S^V \wedge H\underline{\mathbb{Z}})$ are $\dim(V)$ -slices. Hence, the result follows from Proposition 4.8. \square

Remark 5.3. If the spectrum $S^V \wedge H\underline{\mathbb{Z}}$ is not an n -slice, then we will construct certain (co)fiber sequences to study the slices of this spectrum. For $\ell \geq 1$, repeated applications of the map $u_{(\xi-\xi^p)}$ in Lemma 3.5 yields a map $H\underline{\mathbb{Z}} \rightarrow S^{\ell(\xi-\xi^p)} \wedge H\underline{\mathbb{Z}}$. Then smashing with S^V yields a map $S^V \wedge H\underline{\mathbb{Z}} \rightarrow S^{V+\ell(\xi-\xi^p)} \wedge H\underline{\mathbb{Z}}$. We denote it by $u_{\ell(\xi-\xi^p)}^V$ and its cofiber by $C(V, \ell, p)$. For the prime q , one has the similar construction. This cofiber spectrum plays an important role in analyzing slices of the spectrum $S^V \wedge H\underline{\mathbb{Z}}$. To understand the slices of $C(V, \ell, p)$, we begin with the following result.

Lemma 5.4. *The spectrum $\Sigma^n H\mathcal{K}_p\langle\mathbb{Z}/p\rangle$ is a pn -slice and $\Sigma^n H(\mathcal{K}_p\langle\mathbb{Z}/p\rangle \oplus \mathcal{K}_q\langle\mathbb{Z}/q\rangle)$ is a npq -slice.*

Proof. As $\Sigma^n H\mathcal{K}_p\langle\mathbb{Z}/p\rangle$ has homotopy groups concentrated only in degree n , it is $(n+1)$ -coconnective. So, by Proposition 4.5 $F^{s+1}\underline{\pi}_n \Sigma^n H\mathcal{K}_p\langle\mathbb{Z}/p\rangle \neq F^s \underline{\pi}_n \Sigma^n H\mathcal{K}_p\langle\mathbb{Z}/p\rangle$ if and only if $\mathcal{F}^{(s+n)/n} \mathcal{K}_p\langle\mathbb{Z}/p\rangle \neq \mathcal{F}^{(s+n-1)/n} \mathcal{K}_p\langle\mathbb{Z}/p\rangle$. This can only happen when

$$(s+n-1)/n < p \text{ and } (s+n)/n \geq p.$$

This gives $s = n(p-1)$ and so,

$$F^{s+1}\underline{\pi}_n \Sigma^n H\mathcal{K}_p\langle\mathbb{Z}/p\rangle = 0 \text{ and } F^s \underline{\pi}_n \Sigma^n H\mathcal{K}_p\langle\mathbb{Z}/p\rangle \cong \mathcal{K}_p\langle\mathbb{Z}/p\rangle.$$

The quotient $F^s \underline{\pi}_{t-s}(\Sigma^n H\mathcal{K}_p\langle\mathbb{Z}/p\rangle) / F^{s+1} \underline{\pi}_{t-s}(\Sigma^n H\mathcal{K}_p\langle\mathbb{Z}/p\rangle)$ can only be non-zero when $t-s = n$ and hence, $t = np$. Therefore, $\Sigma^n H\mathcal{K}_p\langle\mathbb{Z}/p\rangle$ is an np -slice.

The result for $\mathcal{K}_p\langle\mathbb{Z}/p\rangle \oplus \mathcal{K}_q\langle\mathbb{Z}/q\rangle$ can be proved analogously. \square

Proposition 5.5. *For $\ell \geq 1$, the cofiber spectrum $C(V, \ell, p)$ has only kp -slices for each $k \in \{\dim(V^{C_p}) - 2\ell, \dots, \dim(V^{C_p}) - 2\}$.*

Proof. First, we compute the Mackey functor valued homotopy groups of the cofiber $C(V, \ell, p)$. Note that the restriction of the cofiber to the subgroup C_q is

trivial and the homotopy groups of $C(V, \ell, p)$ are cohomological. Using Proposition 3.3, we conclude that $\underline{\pi}_k C(V, \ell, p)$ is non-zero if and only if $\underline{\pi}_k i_{C_p}^* C(V, \ell, p)$ is non-zero. Set $i_{C_p}^*(V) = m + n\xi$ for $m, n \geq 0$. The cofiber sequence

$$S^{m+n\xi} \wedge H\underline{\mathbb{Z}} \rightarrow S^{m+n\xi+\ell(\xi-2)} \wedge H\underline{\mathbb{Z}} \rightarrow i_{C_p}^* C(V, \ell, p)$$

yields the long exact sequence:

$$\dots \underline{H}_{C_p}^{m-k+n\xi}(S^0) \rightarrow \underline{H}_{C_p}^{m-k-2\ell+(n+\ell)\xi}(S^0) \rightarrow \underline{\pi}_k i_{C_p}^* C(V, \ell, p) \rightarrow \underline{H}_{C_p}^{m+1-k+n\xi}(S^0) \dots$$

Incorporating the computation (3) in the above long exact sequence, one yields

$$\underline{\pi}_k(C(V, \ell, p)) \cong \begin{cases} \mathcal{K}_p\langle \mathbb{Z}/p \rangle & \text{if } k \in \{m - 2\ell, \dots, m - 2\} \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, the map $u_{\xi-\xi p}^V$ induces a cofiber sequence as follows

$$C(V, \ell - 1, p) \xrightarrow{u_{\xi-\xi p}^V} C(V, \ell, p) \rightarrow \Sigma^{m-2\ell} H\mathcal{K}_p\langle \mathbb{Z}/p \rangle \tag{7}$$

with $C(V, 1, p) \cong \Sigma^{m-2} H\mathcal{K}_p\langle \mathbb{Z}/p \rangle$.

To compute the slices of $C(V, \ell, p)$, we use induction on ℓ . For $\ell = 1$, it is clear from Lemma 5.4 that $C(V, 1, p)$ has only one $(m - 2)p$ -slice. Using (7) it is clear that $C(V, \ell, p)$ has an $(m - 2)p$ -slice as $\Sigma^{m-2} H\mathcal{K}_p\langle \mathbb{Z}/p \rangle$. The other slices of $C(V, \ell, p)$ are obtained from the induction hypothesis and [19, Proposition 2.32]. Hence, the result follows. \square

Remark 5.6. Using Lemma 3.5 (b), construct a map

$$u_{\ell\xi}^V : S^V \wedge H\underline{\mathbb{Z}} \rightarrow S^{V+\ell(\xi-2)} \wedge H\underline{\mathbb{Z}}$$

analogous to $u_{\ell(\xi-\xi p)}^V$. We let $\mathcal{S}(p) = \{\dim(V^{C_p}) - 2\ell, \dots, \dim(V^{C_p}) - 2\}$. Then direct computation yields

$$\underline{\pi}_k(\text{cofiber}(u_{l\xi})) = \begin{cases} \mathcal{K}_p\langle \mathbb{Z}/p \rangle \oplus \mathcal{K}_q\langle \mathbb{Z}/q \rangle & \text{if } k \in \mathcal{S}(p) \cap \mathcal{S}(q) \\ \mathcal{K}_p\langle \mathbb{Z}/p \rangle & \text{if } k \in \mathcal{S}(p) \cap \mathcal{S}(q)^c \\ \mathcal{K}_q\langle \mathbb{Z}/q \rangle & \text{if } k \in \mathcal{S}(p)^c \cap \mathcal{S}(q) \\ 0, & \text{otherwise.} \end{cases}$$

This allows us to identifies the slices for the cofiber of $u_{l\xi}$ using Lemma 5.4.

Theorem 5.7. *For a real C_{pq} -representation V with dimension ν , the ν -slice of the C_{pq} -spectrum $S^V \wedge H\underline{\mathbb{Z}}$ is spherical. In particular, if $i_{C_p}^*(S^V \wedge H\underline{\mathbb{Z}}) \leq \nu$ and $i_{C_q}^*(S^V \wedge H\underline{\mathbb{Z}}) \leq \nu$, then for $0 \leq k < \nu$, the k -slice of $S^V \wedge H\underline{\mathbb{Z}}$ is either $\Sigma^{k/p} H\mathcal{K}_p\langle \mathbb{Z}/p \rangle$ or $\Sigma^{k/q} H\mathcal{K}_q\langle \mathbb{Z}/q \rangle$ or the wedge of the two.*

Proof. Using Remark 3.6, assume $V = a + b\xi + c\xi^p + d\xi^q$ for some a, b, c and d are non-negative integers. Note that for any C_p -representation W , by Lemma 4.6 and Lemma 4.7, the C_p -spectrum $S^W \wedge H\underline{\mathbb{Z}}$ is either slice d -connective or slice d -coconnective, for $d = \dim(W)$. Therefore, to identify the slices of the C_{pq} -spectrum $S^V \wedge H\underline{\mathbb{Z}}$ it is enough to prove the following four cases:

Case (i). $i_{C_p}^*(S^V \wedge H\underline{\mathbb{Z}}) \leq \nu$ and $i_{C_q}^*(S^V \wedge H\underline{\mathbb{Z}}) \leq \nu$.

We want to construct a spherical ν -slice mapping to $S^V \wedge H\underline{\mathbb{Z}}$ with cofiber $\leq \nu - 1$. By Lemma 4.6, the hypothesis of (i) yields

$$a + 2c \leq \frac{2(b+d) + 3p}{p-1} \text{ and } a + 2d \leq \frac{2(b+c) + 3q}{q-1}.$$

By Proposition 4.12, there exist C_p - and C_q -representations $m + n\xi$ and $m' + n'\xi$, respectively, such that $S^{m+n\xi} \wedge H\underline{\mathbb{Z}}$ is the ν -slice of $i_{C_p}^*(S^V \wedge H\underline{\mathbb{Z}})$ and $S^{m'+n'\xi} \wedge H\underline{\mathbb{Z}}$ is the ν -slice of $i_{C_q}^*(S^V \wedge H\underline{\mathbb{Z}})$. By Lemma 4.10, it follows that $b + d > n$ and $b + c > n'$.

For $V' = V - (b + d - n)(\xi - \xi^p)$, we have

$$i_{C_p}^*(V') = m + n\xi \text{ and } i_{C_q}^*(V') = i_{C_q}^*(V).$$

By Proposition 5.5, there exists a map

$$u_{(b+d-n)(\xi-\xi^p)}^{V'} : S^{V'} \wedge H\underline{\mathbb{Z}} \rightarrow S^V \wedge H\underline{\mathbb{Z}},$$

and the cofiber $C(V', b + d - n, p)$ has slices only in filtrations $p(m - 2(b + d - n)), \dots, p(m - 2)$.

By Proposition 4.12,

$$\begin{aligned} m = \dim(V'^{C_p}) &= \dim(V) - 2d_{\dim(V)} = \frac{\dim(V) - \dim(V)_0}{p} + \delta \\ &\leq \frac{\dim(V) - 1}{p} + 1 \text{ (by (5)).} \end{aligned}$$

It readily follows $p(m - 2) < \nu$. Therefore, the cofiber $C(V', b + d - n, p)$ is slice $(\nu - 1)$ -connective.

Since $i_{C_q}^*(S^V \wedge H\underline{\mathbb{Z}}) \leq \nu$, we may define a representation

$$V'' := V - (b + c - n')(\xi - \xi^q),$$

and deduce $i_{C_q}^*(V'') = m' + n'\xi$ and $i_{C_p}^*(V'') = i_{C_p}^*(V)$. By Proposition 5.5, there is a map $u_{(b+c-n')(\xi-\xi^q)}^{V''} : S^{V''} \wedge H\underline{\mathbb{Z}} \rightarrow S^V \wedge H\underline{\mathbb{Z}}$ and as above the associated cofiber $C(V'', b + c - n', q)$ has slices in filtration $< \nu$, so $C(V'', b + c - n', q) \leq \nu - 1$.

Finally, set

$$\widehat{V} := V - (b + d - n)(\xi - \xi^p) - (b + c - n')(\xi - \xi^q).$$

Then, $i_{C_p}^*(S^{\widehat{V}} \wedge H\underline{\mathbb{Z}})$ (respectively, $i_{C_q}^*(S^{\widehat{V}} \wedge H\underline{\mathbb{Z}})$) is the ν -slice for $i_{C_p}^*(S^V \wedge H\underline{\mathbb{Z}})$ (respectively, $i_{C_q}^*(S^V \wedge H\underline{\mathbb{Z}})$). By Proposition 5.1, the C_{pq} -spectrum $S^{\widehat{V}} \wedge H\underline{\mathbb{Z}}$ is thus the ν -slice of $S^V \wedge H\underline{\mathbb{Z}}$.

Case (ii). $i_{C_p}^*(S^V \wedge H\underline{\mathbb{Z}}) \geq \nu$ and $i_{C_q}^*(S^V \wedge H\underline{\mathbb{Z}}) \leq \nu$.

By Lemma 4.6,

$$\ell_p = \lceil \frac{p \dim(V^{C_p}) - \dim(V) - 3p}{2p} \rceil \tag{8}(p)$$

is the least positive integer such that the C_{pq} -spectrum $S^{V+\ell_p(\xi-\xi^p)} \wedge H\underline{\mathbb{Z}} \leq \nu$. Then the spectrum $i_{C_p}^*(S^{V+\ell_p(\xi-\xi^p)} \wedge H\underline{\mathbb{Z}}) \leq \nu$. For such l , by Proposition 5.5, there exists a map

$$u_{\ell_p(\xi-\xi^p)}^V : S^V \wedge H\underline{\mathbb{Z}} \rightarrow S^{V+\ell_p(\xi-\xi^p)} \wedge H\underline{\mathbb{Z}}$$

and the fiber of this map (equivalently, $\Sigma^{-1}C(V, \ell_p, p)$) has slice filtration $\geq \nu + 1$. Now, as $i_{C_p}^*(S^{V+\ell_p(\xi-\xi^p)} \wedge H\underline{\mathbb{Z}}) \leq \nu$ and $i_{C_q}^*(S^{V+\ell_p(\xi-\xi^p)} \wedge H\underline{\mathbb{Z}}) \leq \nu$, by case (i), we have the spherical ν -slice and also all the lower slices can be determined. Hence, the result follows.

Case (iii). $i_{C_p}^*(S^V \wedge H\underline{\mathbb{Z}}) \leq \nu$ and $i_{C_q}^*(S^V \wedge H\underline{\mathbb{Z}}) \geq \nu$.

It is analogous to case (ii).

Case (iv). $i_{C_p}^*(S^V \wedge H\underline{\mathbb{Z}}) \geq \nu$ and $i_{C_q}^*(S^V \wedge H\underline{\mathbb{Z}}) \geq \nu$.

Then $\ell_0 := \max\{\ell_p, \ell_q\}$ is the positive integer such that $i_{C_p}^*(S^{V+\ell_0(\xi-2)} \wedge H\underline{\mathbb{Z}}) \leq \nu$ and $i_{C_q}^*(S^{V+\ell_0(\xi-2)} \wedge H\underline{\mathbb{Z}}) \leq \nu$. Now we are in case (i). Therefore, by Proposition 4.12, there exists a C_p -representation $m+n\xi$ and a C_q -representation $m'+n'\xi$ such that $S^{m+n\xi} \wedge H\underline{\mathbb{Z}}$ (resp., $S^{m'+n'\xi} \wedge H\underline{\mathbb{Z}}$) is the spherical ν -slice of $i_{C_p}^*(S^{V+\ell_0(\xi-2)} \wedge H\underline{\mathbb{Z}})$ (resp., $i_{C_q}^*(S^{V+\ell_0(\xi-2)} \wedge H\underline{\mathbb{Z}})$).

By hypothesis (iv), Proposition 4.1 yields

$$a + 2c \geq \frac{\nu}{p} \text{ and } a + 2d \geq \frac{\nu}{q},$$

which implies that $b + d + \ell_0 \leq n$ and $b + c + l_0 \leq n'$. So, we set

$$\widehat{V} = V + \ell_0(\xi - 2) + (n - b - d - \ell_0)(\xi - \xi^p) + (n' - b - c - \ell_0)(\xi - \xi^q),$$

as in case (i) we see that the C_{pq} -spectrum $S^{\widehat{V}} \wedge H\underline{\mathbb{Z}}$ is the ν -slice for $S^V \wedge H\underline{\mathbb{Z}}$, which is spherical. \square

Corollary 5.8. Any $\alpha \in RO(C_{pq})$ has $\beta \in RO(C_{pq})$ such that $S^\alpha \wedge H\underline{\mathbb{Z}}$ has a $\dim(\alpha)$ -slice $S^\beta \wedge H\underline{\mathbb{Z}}$. The other slices of $S^\alpha \wedge H\underline{\mathbb{Z}}$ are suspensions of $H\mathcal{K}_p\langle \mathbb{Z}/p \rangle$ or $H\mathcal{K}_q\langle \mathbb{Z}/q \rangle$ or wedges of the two.

Proof. We can always find some $k \in \mathbb{Z}$ such that $\alpha + k\rho_{C_{pq}}$ is a non-virtual representation of C_{pq} . Therefore, using Proposition 4.4 it is enough to consider $\alpha = V = a + b\xi + c\xi^p + d\xi^q$ for a, b, c , and d non-negative integers. Proposition 5.5 and Theorem 5.7 together then imply the result. \square

Example 5.9. For the cyclic group C_{15} with $p = 3$ and $q = 5$ we shall write a slice tower of the C_{15} -spectrum $S^6 \wedge H\underline{\mathbb{Z}}$. Note that this type of spectrum was studied by Yarnall in [20] for $G = C_{p^k}$, in particular for C_p . In our case, the restrictions satisfy

$$i_{C_3}^*(S^6 \wedge H\underline{\mathbb{Z}}) \geq 6 \text{ and } i_{C_5}^*(S^6 \wedge H\underline{\mathbb{Z}}) \geq 6,$$

so, we are in case (iv) of the Theorem 5.7. Here $\ell_0 = \max\{\ell_3, \ell_5\} = 2$ and $n = 2 = n'$. Therefore, by construction of \widehat{V} in case (iv) of the theorem:

$$\widehat{V} = 6 + 2(\xi - 2) + (2 - 2)(\xi - \xi^3) + (2 - 2)(\xi - \xi^5) = 2 + 2\xi.$$

Hence, using Remark 5.6, the slice tower is:

$$\begin{array}{ccc} 45 - \text{slice} : & \Sigma^3 H(\mathcal{K}_q\langle \mathbb{Z}/q \rangle \oplus \mathcal{K}_p\langle \mathbb{Z}/p \rangle) & \longrightarrow S^6 \wedge H\underline{\mathbb{Z}} \\ & & \downarrow u_\xi \\ 6 - \text{slice} : & & S^{4+\xi} \wedge H\underline{\mathbb{Z}}. \end{array}$$

Example 5.10. Consider the C_{15} -spectrum $X = S^{11\xi^5} \wedge H\underline{\mathbb{Z}}$. Note that

$$i_{C_3}^*(X) \cong S^{11\xi} \wedge H\underline{\mathbb{Z}} \text{ and } i_{C_5}^*(X) \cong S^{22} \wedge H\underline{\mathbb{Z}}.$$

Therefore, we are in case (iv) of Theorem 5.7 as $i_{C_3}^*(X) \leq 22$ and $i_{C_5}^*(X) \geq 22$.

By (8)(q), $l_5 = 8$, so $i_{C_5}^*(S^{8(\xi - \xi^5)} \wedge X) \leq 22$, and $i_{C_5}^*(S^{8\xi + 3\xi^5} \wedge H\underline{\mathbb{Z}}) \leq 22$. Above the 22-slice, all higher dimension slices are obtained by the computations of the fiber of the map $u_{\xi - \xi^q}$. By repeated use of Theorem 5.7 (iv), we obtain the slice tower of $S^{11\xi^5} \wedge H\underline{\mathbb{Z}}$ in filtrations > 22 .

Since, the spectrum $S^{8\xi + 3\xi^5} \wedge H\underline{\mathbb{Z}}$ satisfies both

$$i_{C_3}^*(S^{8\xi + 3\xi^5} \wedge H\underline{\mathbb{Z}}) \leq 22 \text{ and } i_{C_5}^*(S^{8\xi + 3\xi^5} \wedge H\underline{\mathbb{Z}}) \leq 22,$$

so we are in case (i) of Theorem 5.7. Now we compute $d_{22} = 7$ for $p = 3$ and $d_{22} = 8$ for $p = 5$. (see Remark 4.9) By Theorem 5.7 we have

$$\widehat{V} : = 8\xi + 3\xi^5 - (11 - 7)(\xi - \xi^3) = 4\xi + 4\xi^3 + 3\xi^5$$

with $S^{\widehat{V}} \wedge H\underline{\mathbb{Z}}$ the 22-slice of $S^{11\xi^5}$.

Thus, the slice tower for $S^{11\xi^q} \wedge H\underline{\mathbb{Z}}$ is

$$\begin{array}{rcl}
 95\text{-slice:} & \Sigma^{19}H\mathcal{K}_q\langle\mathbb{Z}/q\rangle & \longrightarrow S^{11\xi^q} \wedge H\underline{\mathbb{Z}} \\
 & & \downarrow u_{\xi-\xi^q} \\
 85\text{-slice:} & \Sigma^{17}H\mathcal{K}_q\langle\mathbb{Z}/q\rangle & \longrightarrow S^{\xi+10\xi^q} \wedge H\underline{\mathbb{Z}} \\
 & & \downarrow u_{\xi-\xi^q} \\
 75\text{-slice:} & \Sigma^{15}H\mathcal{K}_q\langle\mathbb{Z}/q\rangle & \longrightarrow S^{2\xi+9\xi^q} \wedge H\underline{\mathbb{Z}} \\
 & & \downarrow u_{\xi-\xi^q} \\
 65\text{-slice:} & \Sigma^{13}H\mathcal{K}_q\langle\mathbb{Z}/q\rangle & \longrightarrow S^{3\xi+8\xi^q} \wedge H\underline{\mathbb{Z}} \\
 & & \downarrow u_{\xi-\xi^q} \\
 55\text{-slice:} & \Sigma^{11}H\mathcal{K}_q\langle\mathbb{Z}/q\rangle & \longrightarrow S^{4\xi+7\xi^q} \wedge H\underline{\mathbb{Z}} \\
 & & \downarrow u_{\xi-\xi^q} \\
 45\text{-slice:} & \Sigma^9H\mathcal{K}_q\langle\mathbb{Z}/q\rangle & \longrightarrow S^{5\xi+6\xi^q} \wedge H\underline{\mathbb{Z}} \\
 & & \downarrow u_{\xi-\xi^q} \\
 35\text{-slice:} & \Sigma^7H\mathcal{K}_q\langle\mathbb{Z}/q\rangle & \longrightarrow S^{6\xi+5\xi^q} \wedge H\underline{\mathbb{Z}} \\
 & & \downarrow u_{\xi-\xi^q} \\
 25\text{-slice:} & \Sigma^5H\mathcal{K}_q\langle\mathbb{Z}/q\rangle & \longrightarrow S^{7\xi+4\xi^q} \wedge H\underline{\mathbb{Z}} \\
 & & \downarrow u_{\xi-\xi^q} \\
 22\text{-slice:} & S^{4\xi+4\xi^3+3\xi^5} \wedge H\underline{\mathbb{Z}} & \longrightarrow S^{8\xi+3\xi^q} \wedge H\underline{\mathbb{Z}} \\
 & & \downarrow u_{\xi-\xi^p} \\
 18\text{-slice:} & \Sigma^6H\mathcal{K}_p\langle\mathbb{Z}/p\rangle & \longrightarrow \bigvee_{i \in \{0,1,2,3\}} \Sigma^{2i} HK_p \langle \mathbb{Z}/p \rangle \\
 & & \downarrow u_{\xi-\xi^p} \\
 12\text{-slice:} & \Sigma^4H\mathcal{K}_p\langle\mathbb{Z}/p\rangle & \longrightarrow \bigvee_{i \in \{0,1,2\}} \Sigma^{2i} HK_p \langle \mathbb{Z}/p \rangle \\
 & & \downarrow u_{\xi-\xi^p} \\
 6\text{-slice:} & \Sigma^2H\mathcal{K}_p\langle\mathbb{Z}/p\rangle & \longrightarrow H\mathcal{K}_p\langle\mathbb{Z}/p\rangle \vee \Sigma^2H\mathcal{K}_p\langle\mathbb{Z}/p\rangle \\
 & & \downarrow u_{\xi-\xi^p} \\
 0\text{-slice:} & & H\mathcal{K}_p\langle\mathbb{Z}/p\rangle
 \end{array}$$

References

- [1] BASU, SAMIK; GHOSH, SUROJIT. Computations in C_{pq} -Bredon cohomology. *Math. Z.* **293** (2019), no. 3-4, pp. 1443-1487. MR4024594, Zbl 1435.55003, arXiv:1612.02159, doi: 10.1007/s00209-019-02248-2. 1403, 1404
- [2] BASU, SAMIK; GHOSH, SUROJIT. Equivariant cohomology for cyclic groups of square-free order. Preprint, 2020. arXiv:2006.09669. 1406
- [3] CARUSO, JEFFREY L. Operations in equivariant \mathbb{Z}/p -cohomology. *Math. Proc. Cambridge Philos. Soc.* **126** (1999), no. 3, pp. 521-541. MR1684248, Zbl 0933.55009, doi: 10.1017/S0305004198003375.
- [4] DUGGER, DANIEL. An Atiyah-Hirzebruch spectral sequence for KR -theory. *K-Theory* **35** (2005), no. 3-4, pp. 213-256. MR2240234, Zbl 1109.14024, arXiv:math/0304099, doi: 10.1007/s10977-005-1552-9. 1405
- [5] FERLAND, KEVIN K. On the $RO(G)$ -graded equivariant ordinary cohomology of generalized G -cell complexes for $G = \mathbb{Z}/p$. Ph.D. Thesis, Syracuse University, 1999. 176 pp. ISBN: 978-0599-42024-3. MR2699528. 1407, 1408
- [6] GREENLEES, JOHN P. C.; MAY, J. PETER. Generalized Tate cohomology. *Mem. Amer. Math. Soc.* **113** (1995), no. 543, viii+178 pp. MR1230773, Zbl 0876.55003, doi: 10.1090/memo/0543.
- [7] GREENLEES, JOHN P. C.; MAY, J. PETER. Equivariant stable homotopy theory. *Handbook of algebraic topology*, 277-323, North-Holland, Amsterdam, 1995. MR1361893, Zbl 0866.55013, doi: 10.1016/B978-044481779-2/50009-2.
- [8] GUILLOU, BERTRAND J.; YARNALL, CAROLYN. The Klein four slices of $\Sigma^n H\mathbb{F}_2$. *Math. Z.* **295** (2020), no. 3-4, pp. 1405-1441. MR4125695, Zbl 1454.55004, doi: 10.1007/s00209-019-02433-3. 1402
- [9] HILL, MICHAEL A. The equivariant slice filtration: a primer. *Homology Homotopy Appl.* **14** (2012), no. 2, pp. 143-166. MR3007090, Zbl 1403.55003, arXiv:1107.3582, doi: 10.4310/HHA.2012.v14.n2.a9. 1400
- [10] HILL, MICHAEL A.; HOPKINS, MICHAEL J.; RAVENEL, DOUGLAS, C. On the nonexistence of elements of Kervaire invariant one. *Ann. of Math. (2)* **184** (2016), no. 1, pp. 1-262. MR3505179, Zbl 1366.55007, arXiv:0908.3724, doi: 10.4007/annals.2016.184.1.1. 1406
- [11] HILL, MICHAEL A.; HOPKINS, MICHAEL J.; RAVENEL, DOUGLAS, C. The slice spectral sequence for the C_4 analog of real K -theory. *Forum Math.* **29** (2017), no. 2, pp. 383-447. MR3619120, Zbl 1362.55009, arXiv:1502.07611, doi: 10.1515/forum-2016-0017. 1400, 1402, 1406, 1410
- [12] HILL, MICHAEL A.; YARNALL, CAROLYN. A new formulation of the equivariant slice filtration with applications to C_p -slices. *Proc. Amer. Math. Soc.* **146** (2018), no. 8, pp. 3605-3614. MR3803684, Zbl 1395.55014, arXiv:1703.10526, doi: 10.1090/proc/13906. 1400
- [13] LEWIS, L. GAUNCE, JR. The $RO(G)$ -graded equivariant ordinary cohomology of complex projective spaces with linear \mathbb{Z}/p actions. *Algebraic topology and transformation groups* (Göttingen, 1987), 53-122. Lecture Notes in Math., 1361, Math. Gottingensis. Springer, Berlin, 1988. MR979507, Zbl 0669.57024, doi: 10.1007/BFb0083034.
- [14] MANDELL, MICHAEL A. & MAY, J. PETER. Equivariant orthogonal spectra and S -modules. *Mem. Amer. Math. Soc.* **159** (2002), no. 755, x+108 pp. MR1922205, Zbl 1025.55002, doi: 10.1090/memo/0755. 1406, 1409
- [15] ULLMAN, JOHN. On the regular slice spectral sequence. Ph.D. Thesis, Massachusetts Institute of Technology, 2013. *ProQuest LLC*. MR3211466. 1401
- [16] SHULMAN, MEGAN. Equivariant local coefficients and the $RO(G)$ -graded cohomology of classifying spaces. Ph.D. Thesis, University of Chicago, 2010. 127 pp. MR2941379, arXiv:1405.1770. 1402

- [17] VOEVODSKY, VLADIMIR. Open problems in the motivic stable homotopy theory. I. *Motives, polylogarithms and Hodge theory, Part I* (Irvine, CA, 1998), 3–34. Int. Press Lect. Ser., 3, I. Int. Press, Somerville, MA, 2002. MR1977582, Zbl 1047.14012. 1405, 1406
1403
- [18] WEBB, PETER. A guide to Mackey functors. *Handbook of algebra*, 2, 805–836. Handb. Algebr., 2, Elsevier/North-Holland, Amsterdam, 2000. MR1759612, Zbl 0972.19001, doi: 10.1016/S1570-7954(00)80044-3. 1399
1404
- [19] WILSON, DYLAN. On categories of slices. Preprint, 2017. arXiv:1711.03472. 1412
- [20] YARNALL, CAROLYN. The slices of $S^n \wedge H\mathbb{Z}$ for cyclic p -groups. *Homology Homotopy Appl.* **19** (2017), no. 1, pp. 1–22. MR3628673, Zbl 1394.55007, doi: 10.4310/HHA.2017.v19.n1.a1.
- [21] YOSHIDA, TOMOYUKI. On G -functors. II. Hecke operators and G -functors. *J. Math. Soc. Japan* **35** (1983), no. 1, 179–190. MR679083, Zbl 0507.20010, doi: 10.2969/jmsj/03510179. 1400, 1408, 1409, 1415
- [22] ZENG, MINGCONG. Equivariant Eilenberg–MacLane spectra in cyclic p -groups. Preprint, 2017. arXiv:1710.01769. 1404
1402

(Surojit Ghosh) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY, ROORKEE, UTTARAKHAND-247667, INDIA
surojitghosh89@gmail.com; surojit.ghosh@ma.iitr.ac.in

This paper is available via <http://nyjm.albany.edu/j/2022/28-59.html>.