

One-sided type-D Ricci-flat metrics

Paul Tod

ABSTRACT. We consider four-dimensional, Riemannian, Ricci-flat metrics for which one or other of the self-dual or anti-self-dual Weyl tensors is type-D in the Petrov-Pirani-Penrose classification. Such metrics always admit a valence-2 Killing spinor, and therefore a Hermitian structure and at least one Killing vector. We rederive the results of Przanowski and collaborators, [14, 15, 16, 17], that these metrics can all be given in terms of a solution of the $SU(\infty)$ -Toda field equation, and show that, when there is a second Killing vector commuting with the first, the method of Ward, [22], can be applied to show that the metrics can also be given in terms of an axisymmetric solution of the flat three-dimensional Laplacian. Thus in particular the field equations linearise.

As a corollary, we show that the same technique linearises the field equations for a four-dimensional Einstein metric with anti-self-dual Weyl tensor and two commuting symmetries.

We reduce the Einstein equations with non-zero scalar curvature and one-sided type-D Weyl tensor, excluding the Kähler-Einstein case, to a modified but not integrable Toda equation.

Some examples of the constructions are given.

CONTENTS

1. Introduction	283
2. One-sided type-D	284
3. A second Killing vector	288
4. Ward's linearisation of the Toda field equation	291
5. One-sided type-D Einstein metrics	295
6. An apparent puzzle	296
7. Examples	297
References	300

Received November 19, 2025.

2020 *Mathematics Subject Classification.* 53B35.

Key words and phrases. gravitational instantons, toric Hermitian metrics.

Thanks to the Institut Mittag-Leffler in Djursholm, Sweden for hospitality as part of the programme ‘General Relativity, Geometry and Analysis’ during September 2019, supported by the Swedish Research Council under grant no. 2016-06596.

1. Introduction

In this note we consider four-dimensional, Riemannian, Ricci-flat metrics for which one or other of the self-dual or anti-self-dual Weyl tensors is type-D in the Petrov-Pirani-Penrose classification (for which see e.g. [12]). We'll call these metrics *one-sided type-D*, a term in use at least since 1984 (see [16]). Our motivation comes indirectly from the discovery of the Chen-Teo metric, [7]. This metric has two commuting Killing vectors and is Riemannian and Ricci-flat and was obtained by inverse-scattering methods, which are available since the Einstein equations in this case are known to be completely integrable, (see e.g. [4]). It was discovered by Aksteiner [2] (see also [3]) that the Chen-Teo metric is one-sided type-D, and therefore, as we shall see, Hermitian. In this note we ask: can one do more with it, in the light of the general solution of one-sided type-D vacuum metrics in [16, 17]?

In [16, 17] the Einstein vacuum equations, subject to these restrictions, are reduced to the $SU(\infty)$ -Toda equation (we'll omit the qualifier " $SU(\infty)$ " henceforth). It's known that the Toda equation linearises if the solution has an extra symmetry, [22], which suggests that the field equations for one-sided type-D Ricci-flat metrics with a second symmetry commuting with the first linearise, and we shall see here that they do.

The main result of this note is the explicit metric form given by (35) subject to (36-38). While this is quite complicated in appearance, it has contributed usefully to the classification of Ricci-flat, toric, Hermitian metrics in [6].

Our method is to start in Section 2 with the assumption of a 4-dimensional Riemannian, Ricci-flat metric which is one-sided type-D, or equivalently (as we show) is Hermitian. Then we use the two-component spinor formalism to rederive the expressions of [17] for the metric in terms of a solution u of the Toda field equation (19). The metric automatically has a Killing vector which arises from a valence-2 Killing spinor, which in turn is a consequence of the type-D-ness (by [21]), and our rederivation serves to explain the occurrence of this Killing vector. Then in Section 3 we add the assumption that there is a second Killing vector and deduce that, after possible redefinitions of coordinates and u preserving the Toda field equation, the second Killing vector must be a symmetry of u corresponding to an ignorable coordinate which can be taken to be y in (19). Then in Section 4 we exploit the observation in [22] that solutions of the Toda field equation independent of y correspond to axisymmetric solutions of the flat 3-dimensional Laplacian. We arrive at our main result: that Riemannian, Ricci-flat, one-sided type-D 4-metrics with two commuting Killing vectors are in one-to-one correspondence with axisymmetric solutions of the flat three-dimensional Laplacian. The field equations, known to be completely integrable in this case, in fact linearise. We also give a corollary: by [14] (see also [18], [20]) the general ASD Einstein metric with a symmetry and non-zero Ricci scalar can be found in terms of a solution of the Toda field equation; consequently if there is a second Killing vector commuting with the first then

again by [22] the solution can be given in terms of a solution of the axisymmetric Laplace equation – in this case too the Einstein equations linearise. In Section 5 we show that the field equations for one-sided type-D Einstein metrics (and by assumption *not* ASD) reduce to a modified Toda field equation which is not integrable ([5]), in Section 6 we deal with an apparently puzzling feature of the constructions, and in Section 7 we give some examples of all the reductions.

Acknowledgements: I am grateful to Dr Steffen Aksteiner, formerly of the Albert Einstein Institute, Golm, for discussions about the Chen-Teo metric, and for telling me that it was one-sided type-D, and to the Institut Mittag-Leffler in Djursholm, Sweden for hospitality as part of the programme ‘General Relativity, Geometry and Analysis’ during September 2019, supported by the Swedish Research Council under grant no. 2016-06596. I am grateful to Dr Maciej Dunajski of DAMTP Cambridge for telling me of references [14, 15, 16, 17] and for useful discussions, in the course of which we realised that the construction given here must be possible, and I am grateful to Profs Jenya Ferapontov and Boris Kruglikov for information about integrability of the modified Toda equation (46).

2. One-sided type-D

In this section, we rederive the results of [17] in the style of [18, 20]. The virtue of this rederivation is that one sees how it follows from the algebraic assumption on the Weyl spinor, and one also sees the origin of the first Killing vector in this assumption. For background on the 2-component spinor formalism see [10], [12] or [11].

Start then with a Riemannian, Ricci-flat and one-sided type-D metric. The unprimed Weyl spinor (as a matter of convenience, call this the SD Weyl spinor – it would more properly be the ASD Weyl spinor but this makes very little difference for us, and reduces the number of primed spinor indices in formulae) can be taken to be

$$\psi_{ABCD} = \psi o_{(A} o_B o_C^\dagger o_{D)}^\dagger \quad (1)$$

with real $\psi = 6\Psi_2$ in NP conventions (so this is essentially the definition of type-D; note in a Riemannian space we have the Hermitian conjugate, denoted by dagger, taking spinors to spinors), and the spinor normalisation

$$o_A o^{\dagger A} = 1.$$

We’ll assume the metric is non-trivially type-D, in the sense that the SD Weyl spinor is not zero, so that ψ is not the zero function. The argument of [21] still goes through to show that the spinor field ω_{AB} defined as¹

$$\omega_{AB} = i\psi^{-1/3} o_{(A} o_{B)}^\dagger \quad (2)$$

is a (real) Killing spinor, or equivalently

$$\nabla_{A'(A} \omega_{B)C} = 0.$$

¹What follows is a local calculation, so we restrict to points where ψ is non-zero.

Therefore

$$\nabla_{A'A}\omega_{BC} = \epsilon_{A(B}K_{C)A'} \quad (3)$$

for some real vector $K_{AA'}$, which is then necessarily a Killing vector. To see this necessity, calculate commutators as follows (conventions following [12]): by Ricci-flat-ness

$$0 = \Delta_{A'B'}\omega^{CD} = \nabla_{(A'}^{(C}K_{B')D)}^D, \quad (4)$$

and with zero Ricci scalar

$$\Delta_{AB}\omega_{CD} = \psi_{EABC}\omega_D^E + \psi_{EABD}\omega_C^E, \quad (5)$$

whence

$$\psi_{E(ABC}\omega_{D)}^E = 0,$$

as we know (this follows from (1), (2)), and the trace of (5) on BC gives

$$\nabla_{AA'}K_D^{A'} = \psi_{ABDE}\omega^{BE} = -\frac{1}{3}\psi\omega_{AD}.$$

From the trace of this we deduce $\nabla_a K^a = 0$, so with (4) we see that K^a is a Killing vector, as claimed, and furthermore we may write

$$\nabla_{AA'}K_{BB'} = -\frac{1}{6}\psi\omega_{AB}\epsilon_{A'B'} + \chi_{A'B'}\epsilon_{AB}, \quad (6)$$

for some symmetric $\chi_{A'B'}$ (which we use in Section 6). The identity

$$\nabla_a\nabla_b K_c = R_{bad}K^d,$$

which holds for any Killing vector, now gives

$$\nabla_{AA'}(-\frac{1}{6}\psi\omega_{BC}) = -\psi_{ABCD}K_{A'}^D,$$

which is equivalent to

$$\omega_{AC}K_{A'}^C = -\frac{1}{6}\psi^{-5/3}\nabla_a\psi, \quad (7)$$

and which also follows by contracting (3) with ω^{BC} , so this is an identity, and

$$\nabla_{AA'}\chi_{B'C'} = -\psi_{A'B'C'D'}K_A^{D'}. \quad (8)$$

We find, introducing ϕ_{AB} as

$$\phi_{AB} = 2io_{(A}o_{B)}^\dagger = 2\psi^{1/3}\omega_{AB}, \quad (9)$$

that there is an almost complex structure determined by

$$J_a^b := \phi_A^B\delta_{A'}^{B'}, \quad (10)$$

which is then easily seen to be integrable – thus the metric is Hermitian by virtue of being type-D and Ricci-flat. Conversely, if there is an integrable complex structure of the form of (10) (i.e. with this duality) then necessarily o^A is geodesic and shear-free and therefore, by the Goldberg-Sachs Theorem (see e.g. [12]), is a repeated PND of the Weyl spinor. Then $o^{\dagger A}$ is another and so the Weyl spinor is type-D. Therefore the properties of being Hermitian and being one-sided type-D are here seen to be equivalent.

With (9) we obtain

$$\phi_{AB} K_{A'}^B = 2\psi^{1/3} \omega_{AB} K_{A'}^B = \nabla_a(\psi^{-1/3}). \quad (11)$$

(By a general argument, this must be a gradient as it defines the Hamiltonian for K).

We can proceed to find a metric ansatz as follows: set $W^{-1} := K^b K_b$ and start to construct an orthonormal basis of one-forms with

$$\theta^0 = W^{1/2} K = W^{-1/2}(dt + \omega),$$

where $K^a \partial_a = \partial_t$ and we've introduced a presently unknown one form ω . Introduce θ^1 as

$$\theta^1 = J\theta^0 = W^{1/2} JK = W^{1/2} \phi_A^B K_{BA'} = -W^{1/2} d(\psi^{-1/3}),$$

using (11), and so $\theta^1 = W^{1/2} dz$ with $z = -\psi^{-1/3}$ (a real coordinate). Next we can choose θ^2, θ^3 orthogonal to θ^0, θ^1 and such that $J\theta^2 = \theta^3$. There will be a complex coordinate ζ such that

$$\theta^2 + i\theta^3 = W^{1/2} e^{u/2} d\zeta,$$

for some real u , and the metric is

$$g = W^{-1}(dt + \omega)^2 + W(dz^2 + e^u d\zeta d\bar{\zeta}), \quad (12)$$

in terms of functions u, W and the one-form ω , all to be determined.

We haven't exhausted the information in integrability of the complex structure. The holomorphic one-forms are

$$e^1 := \theta^0 + i\theta^1 = W^{-1/2}(dt + \omega) + iW^{1/2} dz, \quad e^2 := \theta^2 + i\theta^3 = W^{1/2} e^{u/2} d\zeta,$$

and at once

$$e^1 \wedge e^2 \wedge de^2 = 0,$$

but the vanishing of $e^1 \wedge e^2 \wedge de^1$ leads to

$$d\zeta \wedge (d\omega - idz \wedge dW) = 0, \quad (13)$$

which we leave for the moment (this will be part of the 'monopole' equation, (21) below).

From (6) we have, written as forms

$$dK = -\frac{1}{6}\psi^{2/3}J + \text{ASD terms}, \quad (14)$$

with J the complex structure as a 2-form:

$$J = 2(\theta^0 \wedge \theta^1 + \theta^2 \wedge \theta^3).$$

However, with

$$d\omega = \alpha dy \wedge dz + \beta dz \wedge dx + \gamma dx \wedge dy, \quad (15)$$

for unknowns α, β, γ to be found, we obtain

$$\begin{aligned} dK &= d(W^{-1}(dt + \omega)) = (dt + \omega) \wedge W^{-2} dW + W^{-1} d\omega \\ &= c_1 z^{-2} (\theta^0 \wedge \theta^1 + \theta^2 \wedge \theta^3) + f_1 (\theta^0 \wedge \theta^1 - \theta^2 \wedge \theta^3) + f_2 (\theta^0 \wedge \theta^2 - \theta^3 \wedge \theta^1) + f_3 (\theta^0 \wedge \theta^3 - \theta^1 \wedge \theta^2) \end{aligned}$$

where the f_i are the coefficients of the ASD terms in (14). From this we read off

$$\alpha = -W_x, \beta = -W_y, \gamma = -e^u(W_z - 2c_1 \frac{W^2}{z^2}), \quad (16)$$

and c_1 is a constant fixed by earlier choices (in fact $c_1 = -1/3$ but it is convenient to leave it in the formulae). Note (13) is a consequence of these, but there is a stronger integrability condition to check for (15), namely

$$0 = d^2\omega = (\alpha_x + \beta_y + \gamma_z)dx \wedge dy \wedge dz. \quad (17)$$

We'll come back to this when we have an expression for W .

We next calculate the SD curvature with this choice for $d\omega$ by Cartan calculus based on a normalised triad of SD 2-forms: with an orthonormal basis of one-forms chosen as

$$\theta^0 = W^{-1/2}(dt + \omega), \theta^1 = W^{1/2}dz, \theta^2 = W^{1/2}e^{u/2}dx, \theta^3 = W^{1/2}e^{u/2}dy,$$

define an orthonormal basis of SD 2-forms as

$$\phi^1 = \theta^0 \wedge \theta^1 + \theta^2 \wedge \theta^3, \phi^2 = \theta^0 \wedge \theta^2 + \theta^3 \wedge \theta^1, \phi^3 = \theta^0 \wedge \theta^3 + \theta^1 \wedge \theta^2,$$

solve

$$d\phi^i = -\alpha^i_j \wedge \phi^j \quad (18)$$

for the connection one-forms α^i_j to find

$$\alpha_2^1 = C\theta^2, \alpha_1^3 = -C\theta^3, \alpha_3^2 = E\theta^0 + G\theta^2 + H\theta^3,$$

with

$$E = -c_1 z^{-2} W^{1/2}, G = \frac{1}{2} W^{-1/2} e^{-u/2} u_y, H = -\frac{1}{2} W^{-1/2} e^{-u/2} u_x,$$

and

$$C = -\frac{1}{2} W^{-1/2} u_z - c_1 z^{-2} W^{1/2}.$$

Now obtain the curvature components subject to Ricci flatness from

$$\Omega^i_j = d\alpha^i_j + \alpha^i_k \wedge \alpha^k_j = \Omega^i_{j\cdot k} \phi^k,$$

with $\epsilon^{ijk} \Omega_{ij\cdot k} = 0$ (which encodes vanishing Ricci scalar; these indices are raised and lowered by δ_{ij}, δ^{ij}), to find from $\Omega_2^1 + i\Omega_3^1$ that necessarily

$$W = -\frac{z^2 u_z}{2c_1} + f(z),$$

with

$$f' - 2\frac{f}{z} + c_1 \frac{f^2}{z^2} = 0,$$

so that

$$f = \frac{z^2}{c_1 z + c_2},$$

for a constant c_2 . Then from Ω_3^2 we find that u satisfies the *Toda field equation*:

$$u_{xx} + u_{yy} + (e^u)_{zz} = 0, \quad (19)$$

and $c_1 f = z$ so that $c_2 = 0$ and

$$W = \frac{1}{c_1} z \left(1 - \frac{1}{2} z u_z\right). \quad (20)$$

Now we can check the integrability condition (17). In fact $G = W/z$ satisfies a familiar monopole equation:

$$G_{xx} + G_{yy} + (e^u G)_{zz} = 0. \quad (21)$$

The SD curvature components are now

$$\Omega_2^1 = c_1 z^{-3} \phi^3, \quad \Omega_1^3 = c_1 z^{-3} \phi^2, \quad \Omega_3^2 = -2c_1 z^{-3} \phi^1,$$

which is recognisably type-D, as expected (in this setting, type-D is the condition that the SD Weyl tensor, which is here represented by the real trace-free symmetric matrix $E_{ij} := \epsilon_i^{mn} \Omega_{mn,j}$, should be degenerate, in having a repeated eigen-value).

We have recovered the expressions in [16] as deductions from the assumption of Riemannian, Ricci-flat and either Hermitian or one-sided type-D, and we see why there is always a Killing vector preserving the complex structure.

Note that

- The holomorphic one-forms are now

$$e^1 = W^{-1/2}(dt + \omega) + iW^{1/2}dz, \quad e^2 = W^{1/2}e^{u/2}(dx + idy). \quad (22)$$

- We can always set $c_1 = 1$ by a constant rescaling of the metric and redefinition of some coordinates, and this is the value arrived at in [17]: we've recovered their expressions exactly.
- The 2-form $z^{-2}\phi^1$ is closed: this is the rescaling that makes the metric Kähler (for the fact that there must be one, see e.g. [8]).

3. A second Killing vector

Now suppose there is a second Killing vector, then we may write it as

$$L = A\partial_t + B\partial_x + C\partial_y + D\partial_z,$$

(with C not as in the previous section). We start by showing it must have a restricted form, which then allows the Toda field equation to be linearised.

Write \mathcal{L} for the Lie derivative along L , and assume L commutes with $K = \partial_t$ so that

$$\mathcal{L}(K) = 0,$$

which forces A, B, C, D to be independent of t . Also

$$0 = \mathcal{L}(g(K, K)) = \mathcal{L}(W^{-1}) \text{ so that } \mathcal{L}(W) = 0.$$

The curvature is constant along L so that $\mathcal{L}\psi_{ABCD}$ is zero, therefore $\mathcal{L}\psi = 0$ (with ψ the scalar in (1)) and then by the previous section $\mathcal{L}z = 0$ and so $D = 0$. Also the Lie derivative of ϕ_{AB} must be zero (since $-4\psi_{ABCD} = \psi\phi_{(AB}\phi_{CD)}$), and

so the complex structure is preserved. From the vanishing of $\mathcal{L}(W)$ we now deduce

$$\mathcal{L}(u_z) = 0. \quad (23)$$

Since K, W and J are preserved by L , we deduce

$$\mathcal{L}(e^1) = 0,$$

when, by raising the index, we'll also have

$$0 = \mathcal{L}(W^{1/2}\partial_t + iW^{-1/2}\partial_z) = iW^{-1/2}\mathcal{L}(\partial_z),$$

so that $\mathcal{L}(\partial_z) = 0$ and A, B, C are independent of z .

What can we say about $\mathcal{L}(e^2)$? The complex structure is preserved so

$$\mathcal{L}(e^2) = \alpha e^1 + \beta e^2$$

for some α, β (not to be confused with α, β in (15)). Already

$$\mathcal{L}(e^1) = 0 \text{ so also } \mathcal{L}(\bar{e}^1) = 0$$

and

$$\mathcal{L}(g(\bar{e}^1, e^2)) = 0 \text{ whence } g(\bar{e}^1, \mathcal{L}(e^2)) = 0,$$

so $\alpha = 0$, and by considering $\mathcal{L}(g(\bar{e}^2, e^2))$ we find β pure imaginary: L rotates the o.n. basis in the (θ^2, θ^3) -plane.

From (12) we have the freedom

$$t \rightarrow t + f(x, y), \quad \omega \rightarrow \omega - df,$$

and under this

$$L = A\partial_t + B\partial_x + C\partial_y \rightarrow A\partial_t + B(\partial_x + f_x\partial_t) + C(\partial_y + f_y\partial_t),$$

and in particular

$$A \rightarrow A + Bf_x + Cf_y,$$

which we can exploit to set $A = 0$.

From

$$0 = \mathcal{L}(K) = \mathcal{L}(W^{-1}(dt + \omega)) \text{ we deduce } \mathcal{L}(\omega) = 0.$$

Now all that remains of the Killing equation for L is

$$\mathcal{L}(e^u(dx^2 + dy^2)) = 0, \quad (24)$$

i.e. $L = B(x, y)\partial_x + C(x, y)\partial_y$ is a Killing vector of the 2-metric $h = e^u(dx^2 + dy^2)$. We consider solving this reduced problem in a subsection.

3.1. Solving (24) for L . We are considering the 2-metric

$$h := e^u(dx^2 + dy^2),$$

with $u(x, y, z)$ which for our problem we can assume analytic (as solutions of the Toda equation must be). We perform the following calculation at a fixed but arbitrary z , which we can take to be $z = 0$. First calculate the Ricci scalar of h :

$$R = -e^{-u}(u_{xx} + u_{yy}).$$

We are requiring the existence of a Killing vector

$$L = B(x, y)\partial_x + C(x, y)\partial_y,$$

which must preserve R and so must be orthogonal to dR , so for some $\Omega(x, y)$ we have

$$B = -\Omega R_y, \quad C = \Omega R_x. \quad (25)$$

The Killing equations are

$$0 = \mathcal{L}_L h_{ab} = L^c \partial_c h_{ab} + h_{ac} \partial_b L^c + h_{cb} \partial_a L^c.$$

With $ab = 12$ this gives

$$B_y + C_x = 0, \quad (26)$$

while with $ab = 11$ or 22

$$B_x = C_y = -\frac{1}{2}L(u). \quad (27)$$

Substitute from (25) into (26) and (27) to obtain the following system for $d\Omega$:

$$\begin{aligned} \Omega_x R_x - \Omega_y R_y &= \Omega(R_{yy} - R_{xx}) \\ \Omega_x R_y + \Omega_y R_x &= -2\Omega R_{xy}, \end{aligned}$$

which algebraically solves to give

$$\begin{aligned} \frac{\Omega_x}{\Omega} &= D^{-1}(R_x(R_{yy} - R_{xx}) - 2R_y R_{xy}), \\ \frac{\Omega_y}{\Omega} &= D^{-1}(-R_y(R_{xx} - R_{yy}) - 2R_x R_{xy}), \end{aligned}$$

with $D = (R_x)^2 + (R_y)^2 = |dR|^2$ (note this is *not* $h(dR, dR)$). This can be simplified by moving some terms to the left since

$$D_x = 2R_x R_{xx} + 2R_y R_{xy}, \quad D_y = 2R_x R_{xy} + 2R_y R_{yy}$$

so that

$$\begin{aligned} (\log(\Omega D))_x &= D^{-1}R_x(R_{xx} + R_{yy}), \\ (\log(\Omega D))_y &= D^{-1}R_y(R_{xx} + R_{yy}), \end{aligned}$$

or

$$d(\log(\Omega D)) = D^{-1}\Delta_0 R dR, \quad (28)$$

with $\Delta_0 R := R_{xx} + R_{yy}$ (which again is *not* Δ_h). Integrability for (28) is then

$$d(D^{-1}\Delta_0 R) \wedge dR = 0,$$

which is solved by

$$D^{-1}\Delta_0 R = F(R)$$

for some (analytic) F . This says

$$\Delta_0 R + F(R)|\nabla R|^2 = 0$$

so solve $G''/G' = F$ for G to deduce

$$\nabla \cdot (G' \nabla R) = 0,$$

and

$$\Delta(G(R)) = 0,$$

with Δ_0 or Δ_h .

We have a **necessary condition: some function of R is harmonic**. Next we exploit the coordinate freedom: with $\zeta = x + iy$ we can make the change

$$\zeta \rightarrow \hat{\zeta} = f(\zeta), \hat{x} = \phi(x, y), \hat{y} = \psi(x, y),$$

where ϕ, ψ are conjugate harmonic functions, if we accompany this by

$$u \rightarrow \hat{u} = u - \log f' - \log(\bar{f}'),$$

and then R is unchanged (this is also of course a symmetry of the Toda field equation (19)). Now if $G(R)$ is harmonic then there is a new coordinate system in which it is \hat{x} . Drop the hats then w.l.o.g. $R = R(x)$ and the candidate Killing vector is ∂_y . (If preferred we could choose new coordinates so that $R = R(r)$ i.e. radially symmetric at least in some neighbourhood, possibly not including either the origin or a complete circle.)

We have

$$L = \partial_y \text{ and } u_y = 0 \text{ at } z = 0. \quad (29)$$

By (23) we shall also have

$$0 = \mathcal{L}(u_z) = u_{zy}$$

at $z = 0$. Now by uniqueness of solution for the Toda field equation (19) we shall have $u_y = 0$ for all z , and the second Killing vector is, without loss of generality, $L = \partial_y$ everywhere.

4. Ward's linearisation of the Toda field equation

In this section, we follow [22] to relate the y -independent Toda field equation:

$$u_{xx} + (e^u)_{zz} = 0, \quad (30)$$

to the axisymmetric Laplace equation in cylindrical polars:

$$V_{zz} + R^{-1}(RV_R)_R = 0. \quad (31)$$

This will lead to a solution of the Ricci-flat equations for the general one-sided type-D metric with an extra symmetry in terms of a solution of the axisymmetric Laplace equation in three dimensions.

To see how u and V are related, and following [22], set

$$x = V_Z, \quad z = \frac{1}{2}RV_R, \quad u = \log(R^2/4), \quad (32)$$

(so we need to suppose that V_R, V_Z are not constant). We calculate the Jacobian matrix

$$\frac{\partial(x, z)}{\partial(R, Z)} = \begin{pmatrix} V_{ZR} & -\frac{1}{2}RV_{ZZ} \\ V_{ZZ} & \frac{1}{2}RV_{RZ} \end{pmatrix}$$

with the aid of (31), and the inverse is

$$\frac{\partial(R, Z)}{\partial(x, z)} = \Delta^{-1} \begin{pmatrix} \frac{1}{2}RV_{ZR} & \frac{1}{2}RV_{ZZ} \\ -V_{ZZ} & V_{RZ} \end{pmatrix} \quad (33)$$

with $\Delta = \frac{1}{2}R((V_{RZ})^2 + (V_{ZZ})^2)$ (since V_R, V_Z are not constant, Δ is nonzero). In particular therefore

$$R_x = \frac{1}{2}\Delta^{-1}RV_{RZ}, \quad R_z = -\Delta^{-1}V_{ZZ},$$

so with $u = \log(R^2/4)$ as in (32) we deduce

$$u_x = \frac{2}{R}R_x = \Delta^{-1}V_{RZ} = Z_z, \quad e^u u_z = -\frac{R}{2}\Delta^{-1}V_{ZZ} = -Z_x, \quad (34)$$

and by cross-differentiating, we see that u satisfies (30).

With $u_y = 0$, $d\omega$ as in (15) subject to (16) becomes

$$d\omega = -W_x dy \wedge dz - e^u (W_z - 2c_1 \frac{W^2}{z^2}) dx \wedge dy = d(F dy),$$

with

$$F_z = W_x = -\frac{z^2 u_{xz}}{2c_1},$$

and

$$F_x = -e^u (W_z - 2c_1 \frac{W^2}{z^2}) = \frac{1}{2c_1} e^u (-z^2 (u_{zz} + (u_z)^2) + 2zu_z - 2).$$

To integrate for F we need the function H conjugate to V . From (31) this satisfies

$$H_R = RV_Z, \quad H_Z = -RV_R,$$

and then one verifies that

$$F = \frac{1}{2c_1} (z^2 u_x - \frac{1}{2} x R^2 + H).$$

Now transform the metric. First note that

$$\begin{aligned} dz^2 + e^u dx^2 &= \left(\frac{1}{2}(RV_R)_R dR + \frac{1}{2}RV_{RZ} dZ \right)^2 + \frac{R^2}{4} (V_{RZ} dR + V_{ZZ} dZ)^2 \\ &= \frac{R^2}{4} ((-V_{ZZ} dR + V_{RZ} dZ)^2 + (V_{RZ} dR + V_{ZZ} dZ)^2) \\ &= \frac{1}{2}R\Delta(dR^2 + dZ^2), \end{aligned}$$

with the aid of (31) and the definition of Δ . Now the metric is

$$\begin{aligned} g &= W^{-1}(dt + \omega)^2 + W(dz^2 + e^u(dx^2 + dy^2)) \\ &= W^{-1}(dt + Fdy)^2 + We^u dy^2 + \frac{1}{2}WR\Delta(dR^2 + dZ^2) \\ &= \begin{pmatrix} dt & dy \end{pmatrix} \begin{pmatrix} W^{-1} & FW^{-1} \\ FW^{-1} & F^2W^{-1} + We^u \end{pmatrix} \begin{pmatrix} dt \\ dy \end{pmatrix} + \Omega^2(dR^2 + dZ^2) \quad (35) \end{aligned}$$

which is in the canonical form for a Ricci-flat metric with two commuting Killing vectors. Note that the determinant of the matrix of Killing vector contractions is $e^u = R^2/4$ so that R (up to a constant factor) is the standard radial coordinate, and then Z is its harmonic conjugate. The metric components can be given explicitly in terms of V, R, Z via

$$W = \frac{V_R}{2c_1} \left(\frac{R((V_{RZ})^2 + (V_{ZZ})^2) + V_R V_{ZZ}}{((V_{RZ})^2 + (V_{ZZ})^2)} \right), \quad (36)$$

$$F = \frac{1}{2c_1} \left(H + \frac{R(V_R)^2 V_{RZ} - R^2 V_Z ((V_{RZ})^2 + (V_{ZZ})^2)}{2((V_{RZ})^2 + (V_{ZZ})^2)} \right), \quad (37)$$

$$\Omega^2 = \frac{1}{8c_1} R^2 V_R (R((V_{RZ})^2 + (V_{ZZ})^2) + V_R V_{ZZ}). \quad (38)$$

Note that:

- We have linearised the field equations for a Ricci-flat metric with two commuting symmetries. These field equations are already known to be linear if one of the symmetries is hypersurface orthogonal. We can be sure that we haven't inadvertently reduced to this case by looking at the examples which follow, specifically the Riemannian Kerr solution which does not in general admit a hypersurface orthogonal Killing vector.
- We have an expression for the metric given harmonic $V(R, Z)$. We could think about getting back to V starting from g . One route is to set $Q = RV_R$ when

$$8c_1\Omega^2 = Q((Q_Z)^2 + (Q_R)^2 - R^{-1}QQ_R)$$

which needs to be solved for Q given Ω , with Q also subject to

$$Q_{RR} - R^{-1}Q_R + Q_{ZZ} = 0.$$

4.1. A Corollary: linearising the SD Einstein equations with two commuting symmetries. As a corollary to the previous section, we recall that the general solution to the (four-dimensional) SD (or ASD) Einstein equations with a symmetry was given in [14] (see also [18], [20]) and it also depends on a solution to the Toda field equation. Thus with a second symmetry, the Ward transformation can be applied again to give the general solution in this case in terms of an axisymmetric solution of the Laplace equation.

Recall, from [18], the metric in this case is

$$g = \frac{1}{Pz^2}(dt + \omega)^2 + \frac{P}{z^2}(dz^2 + e^u(dx^2 + dy^2)), \quad (39)$$

with

$$u_{xx} + u_{yy} + (e^u)_{zz} = 0, \quad 2\Lambda P = zu_z - 2,$$

where Λ is proportional to the (constant) Ricci scalar, and

$$d\omega = -P_x dy \wedge dz - P_y dz \wedge dx - (Pe^u)_z dx \wedge dy.$$

If we add a second symmetry then the argument goes through as before, and without loss of generality we can suppose $u_y = 0$ and the second symmetry is $L = \partial_y$. We can solve for ω :

$$\omega = Fdy, \quad F = \frac{1}{2\Lambda}(zu_x - Q),$$

with Q conjugate to u in the sense

$$u_x = Q_z, \quad (e^u)_z = -Q_x.$$

We follow the Ward transformation as before, set:

$$x = V_Z, \quad z = \frac{1}{2}RV_R, \quad u = \log(R^2/4),$$

then as before

$$dz^2 + e^u dx^2 = \frac{1}{2}R\Delta(dR^2 + dZ^2),$$

with

$$\Delta = \frac{1}{2}R((V_{RZ})^2 + (V_{ZZ})^2).$$

The metric becomes

$$\begin{aligned} g = & \frac{8\Lambda\Delta}{R^2V_R^2(2\Delta + V_RV_{ZZ})}(dt + Fdy)^2 + \frac{(2\Delta + V_RV_{ZZ})}{2\Lambda\Delta V_R^2}dy^2 \\ & + \frac{(2\Delta + V_RV_{ZZ})}{RV_R^2}(dR^2 + dZ^2). \end{aligned} \quad (40)$$

For F we note from (34) that $Q = Z$ and from the Jacobian matrix (33)

$$zu_x = \frac{1}{2\Delta}RV_RV_{ZZ},$$

so that

$$F = \frac{1}{2\Lambda}(\frac{1}{2\Delta}RV_RV_{ZZ} - Z).$$

This then is the general SD Einstein metric with $\Lambda \neq 0$ and two commuting symmetries, written in terms of a solution V of the axisymmetric Laplace equation (31). We'll give an example in Section 7.

5. One-sided type-D Einstein metrics

It's natural to ask whether the addition of a non-zero scalar curvature interferes with the integrability of the field equations: for ASD vacuum with a symmetry it does not while for vacuum with two commuting Killing vectors it does. In the case considered in this section it does – we can reduce the field equations to a modified Toda field equation but not an integrable one.²

To see this, suppose we still have (1) but with a nonzero Λ , in NP conventions. Then (2) still defines a Killing spinor (the argument in [21] for its existence uses the contracted Bianchi identity, which is unchanged in this case). We need to assume that ψ is not constant as we shall want to use it as a coordinate. By making this assumption we are eliminating some cases: if ψ is constant then K^a is zero and ω_{AB} defines a Kähler form, thus we are excluding Einstein-Kähler metrics. Equation (3) still defines a Killing vector K^a , but (6) is changed to

$$\nabla_{AA'}K_{BB'} = -\frac{1}{6}(\psi + 12\Lambda)\omega_{AB}\epsilon_{A'B'} + \chi_{A'B'}\epsilon_{AB}, \quad (41)$$

with a corresponding change in (14). With $d\omega$ as in (15) we find α and β as in (16) unchanged but

$$\gamma = -e^u(W_z + \frac{2}{3}(\psi + 12\Lambda)\psi^{-1/3}). \quad (42)$$

We may introduce $z = -\psi^{-1/3}$ as before and then it is convenient to introduce

$$g(z) = \frac{1}{3}\psi^{-1/3}(\psi + 12\Lambda) = \frac{12\Lambda z^3 - 1}{3z^2}. \quad (43)$$

The connection coefficients α_j^i of (18) take the same form with G and H unchanged but with

$$C = W^{1/2}g - \frac{1}{2}W^{-1/2}u_z, \quad (44)$$

and

$$E = gW^{1/2}. \quad (45)$$

We calculate the curvature forms Ω_j^i but this time with $\epsilon_{ijk}\Omega_{ij,k} = 24\Lambda$. From $\Omega_2^1 + i\Omega_3^1$ we find

$$gW = -\frac{1}{2}u_z + h(z)$$

with $h' = h^2$ so that $h = -(z - c)^{-1}$ for some c . Then Ω_3^2 leads to $c = 0$ and the modified Toda equation

$$u_{xx} + u_{yy} + (e^u)_{zz} + e^u(A(z)u_z + B(z)) = 0, \quad (46)$$

²This class of metrics was considered in [15] but it is not straightforward to compare results obtained.

for u , with

$$A = \frac{72\Lambda z^2}{1 - 12\Lambda z^3}, \quad B = -\frac{144\Lambda z}{1 - 12\Lambda z^3}.$$

One can now verify that the expression for $d\omega$ is consistent (i.e. that $d(d\omega) = 0$). One can also check that (46) cannot just be transformed back into the Toda equation by the simple change of variable:

$$z \rightarrow Z = f(z), \quad u \rightarrow U = u + F(z). \quad (47)$$

This in fact shows that (46) is not integrable: it is known ([5]) that the only modifications of Toda of the form of (46) which are integrable are those obtained by the transformation (47). We can check the expressions for $A(z)$ and $B(z)$ by obtaining the known one-sided type D Einstein space which is the Schwarzschild-de Sitter metric. We do this in the final section, with other examples. In the next section we dispose of an apparent puzzle concerning the one-sided type-D vacuum and Einstein metrics.

6. An apparent puzzle

From the assumption of one-sided type-D-ness and either vacuum or Einstein, we have been able to reduce the Einstein equations to one of two PDEs for a single function u . It is striking that at no stage do we consider the ASD Weyl spinor, only ever the SD Weyl spinor, but the ASD Weyl spinor is evidently uniquely defined from the metric. This could be contrasted with linear (gravity) theory in which there are degrees of freedom and therefore free data to be specified for both SD and ASD linearised Weyl spinors. One way to explain this apparent puzzle is to point to the Killing vector arising from the Killing spinor, which necessarily satisfies the identity

$$\nabla_a \nabla_b K_c = R_{bcad} K^d, \quad (48)$$

a relation that one doesn't have in linear theory but which, in the full theory, clearly connects the ASD and SD Weyl spinors. Dealing first with one-sided type-D vacuum solutions, we have the Killing spinor ω_{AB} from (2), defining the Killing vector K^a via (3), which in turn satisfies (6). If we contract (6) with K^b we obtain

$$\frac{1}{2} \nabla_a W^{-1} = K^b \nabla_a K_b = -\frac{1}{6} \psi \omega_{AB} K_{A'}^B - \chi_{A'B'} K_A^{B'}$$

which we may rearrange as

$$\chi_{A'B'} K_A^{B'} = \frac{1}{36} \psi^{-2/3} \nabla_a \psi - \frac{1}{2} \nabla_a W^{-1} = \nabla_a Q$$

where we introduce $Q = -\frac{1}{2} W^{-1} - \frac{1}{12} z^{-1}$. Multiplying by $K_{A'}^A$ gives an expression for $\chi_{A'B'}$ as

$$\chi_{A'B'} = W K_{A'}^B \nabla_{B'} Q$$

which is therefore known explicitly when we know u . This in turn is a potential for the SD Weyl spinor via (8) so that, as must be the case, knowledge of u fixes both the ASD and the SD Weyl spinor – there is no more free data.

7. Examples

- Starting with one-sided type-D vacua, flat space is *not* an example of the construction, as we're assuming the SD Weyl spinor isn't zero. However, we could have chosen the zero solution of the Toda equation, when, taking $c_1 = 1$ for simplicity,

$$u = 0, \quad W = z, \quad \omega = xdy,$$

and the metric is

$$g = \frac{1}{z}(dt + xdy)^2 + z(dz^2 + dx^2 + dy^2),$$

which is recognisably the Gibbons-Hawking metric with potential z (see e.g. [9]). In particular, this metric is hyper-Kähler with the other orientation so that the primed Weyl spinor is zero. With $T = \frac{2}{3}z^{3/2}$, it can be written

$$g = dT^2 + \left(\frac{3T}{2}\right)^{3/2} (dx^2 + dy^2) + \left(\frac{2}{3T}\right)^{3/2} (dt + xdy)^2,$$

which makes the isometry group manifest: this is LRS Bianchi-type II. Because $u = 0$, this doesn't have a Ward form.

- There are separable solutions of the Toda equation (19) in the sense $u = f(x, y) + g(z)$ (see e.g. [19]) some of which can be written

$$u = -2 \log(1 + x^2 + y^2) + \log(4(z^2 + 2mz + a)),$$

for real constants a, m , when

$$W = \frac{z(a + mz)}{z^2 + 2mz + a}, \quad \omega = -a \cos \theta d\phi,$$

and we've introduced polar coordinates by $\zeta = \tan(\theta/2)e^{i\phi}$. The metric can be written

$$\begin{aligned} g = & \frac{(z^2 + 2mz + a)}{z(a + mz)} (dt - a \cos \theta d\phi)^2 + \frac{z(a + mz)}{z^2 + 2mz + a} dz^2 \\ & + z(a + mz)(d\theta^2 + \sin^2 \theta d\phi^2), \end{aligned} \tag{49}$$

which has LRS Bianchi-type-IX form. When $a = 0$ it is the Riemannian Schwarzschild solution; with $a = m^2, n = -m^{3/2}/2$ it is the self-dual Taub-NUT metric as given in equation (3.9) of [9]; with $m = 0$ it is the Eguchi-Hanson metric as in (3.20) of [9] but with a^4 there replaced by $-16a^3$ from here (in particular it *isn't* the Riemannian Kerr solution).

There will be more, probably unfamiliar, one-sided type-D metrics determined by the 'quadric ansatz' [19] for solutions of the Toda equation.

- To illustrate the Ward transformation on a one-sided type-D vacuum with an extra symmetry, we consider the particular separable solution of the Toda equation given by

$$u = 2 \log \operatorname{sech} x + \log(z^2 + 2mz + a),$$

when

$$W = \frac{z(mz + a)}{z^2 + 2mz + a}, \quad \omega = a \operatorname{tanh} x dy,$$

and with $\cos \theta = \tanh x$, $\phi = -y$ we arrive at the metric in (49) again, but this time with u such that $u_y = 0$. For simplicity put $a = m^2$, and then

$$R = 2(z + m) \operatorname{sech} x, \quad Z = -2(z + m) \tanh x,$$

and

$$V = -2m \log R + (R^2 + Z^2)^{1/2} - Z \tanh^{-1} \left(\frac{Z}{(R^2 + Z^2)^{1/2}} \right),$$

which one verifies is harmonic. (In spherical polars the terms independent of m are

$$V = r(1 - \cos \theta \log \cot(\theta/2)).$$

- For the transformation (32) to be nontrivial, we need V_R and V_Z to be nonconstant, so for an example with a simple V consider

$$V = R^2 - 2Z^2.$$

Then

$$x = V_Z = -4Z, \quad z = \frac{1}{2}RV_R = R^2, \quad u = \log(R^2/4),$$

and we have the simple solution $u = \log(z/4)$ of (30). After a change of variable, the metric (12) becomes the LRS Riemannian Kasner solution:

$$g = dT^2 + T^{-2/3}dU^2 + T^{4/3}(dX^2 + dY^2).$$

- **Riemannian Kerr:** We start with the NP tetrad (of vectors) tied to the Principal Null Directions for Lorentzian Kerr as given in [1], (24)–(26), lower to one-forms and transform to Boyer-Lindquist coordinates via (35) of [1] to obtain

$$\begin{aligned} \ell &= dt - \frac{\Sigma^2}{\tilde{\Delta}} dr - a \sin^2 \theta d\phi, \quad n = \frac{\tilde{\Delta}}{2\Sigma^2}(dt - a \sin^2 \theta d\phi) + \frac{1}{2} dr, \\ m &= \frac{1}{\sqrt{2}\Gamma}(ia \sin \theta dt - \Sigma^2 d\theta - i(r^2 + a^2) \sin \theta d\phi), \end{aligned}$$

where

$$\Sigma^2 = r^2 + a^2 \cos^2 \theta, \quad \tilde{\Delta} = r^2 - 2mr + a^2, \quad \Gamma = r + ia \cos \theta.$$

In this form we can analytically continue to Riemannian signature by changing (t, a) to (it, ia) . It's convenient to boost and rotate the basis a little to arrive at

$$L = (-i\tilde{\Theta})\ell = \tilde{\Theta}(dt - a \sin^2 \theta d\phi) + \frac{i}{2\tilde{\Theta}}dr, \quad N = \bar{L},$$

and

$$M = \frac{1}{\sqrt{2}\Sigma}(-\Sigma^2 d\theta + i \sin \theta (adt + (r^2 - a^2)d\phi))$$

where now

$$\tilde{\Theta} = \frac{\sqrt{\tilde{\Delta}}}{\Sigma\sqrt{2}}, \quad \tilde{\Delta} = r^2 - 2mr - a^2, \quad \Sigma^2 = r^2 - a^2 \cos^2 \theta.$$

Since both Weyl spinors are type-D we have a choice of complex structures, both of them integrable: one, say J_1 , has L, M as holomorphic one-forms and the other, J_2 , has L, \bar{M} . We lower the Killing vector $K = \partial_t$ and take its exterior derivative to obtain

$$dK = iX(L \wedge \bar{L} + M \wedge \bar{M}) + iY(L \wedge \bar{L} - M \wedge \bar{M})$$

with

$$X = -m(r - a \cos \theta)^{-2}, \quad Y = -m(r + a \cos \theta)^{-2},$$

so if we stick with J_1 then $i(L \wedge \bar{L} + M \wedge \bar{M})$ is the 2-form corresponding to the Killing spinor under consideration, the scalar ψ is a multiple of $(r - a \cos \theta)^{-3}$ and the coordinate z is a multiple of $r - a \cos \theta$. Comparing the Riemannian Kerr metric, which is now

$$\begin{aligned} g = 2\tilde{\Theta}^2(dt - a \sin^2 \theta d\phi)^2 + \frac{dr^2}{2\tilde{\Theta}^2} + \Sigma^2 d\theta^2 \\ + \frac{\sin^2 \theta}{\Sigma^2}(adt + (r^2 - a^2)d\phi)^2 \end{aligned}$$

with the metric form (12) and using $z = r - a \cos \theta$, we are led to

$$e^u d\zeta d\bar{\zeta} = \tilde{\Delta} \sin^2 \theta ((a\tilde{\Delta}^{-1}dr - \csc \theta d\theta)^2 + d\phi^2),$$

when a choice for ζ is

$$\zeta = x + iy = \log \left(\left(\frac{r - m - b}{r - m + b} \right)^{-a/2b} \tan(\theta/2) e^{i\phi} \right),$$

where $b^2 = a^2 + m^2$, so in particular $y = \phi$, and then u is given by

$$e^u = (r^2 - 2mr - a^2) \sin^2 \theta,$$

in agreement with (32) (as this is the determinant of the (ϕ, t) -part of the metric). It is straightforward to verify that

$$u_x = Z_z, \quad e^u u_z = -Z_x,$$

with $Z = 2(r - m) \cos \theta$, so that u does satisfy (30) but we can't obtain $u(x, z)$ explicitly. From (32) we have

$$R^2 = 4e^u = 4(r^2 - 2mr - a^2) \sin^2 \theta,$$

together with

$$Z = 2(r - m) \cos \theta,$$

so that (r, θ) are ellipsoidal coordinates in the (R, Z) plane. It is straightforward to obtain $V(r, \theta)$: we find

$$\begin{aligned} V = & 2(r - a \cos \theta) + 2((r - m) \cos \theta - a) \log \tan(\theta/2) \\ & + 2m \log \sin \theta + ((m + b) - \frac{a}{b}(r - m) \cos \theta) \log(r - m - b) \\ & + ((m - b) + \frac{a}{b}(r - m) \cos \theta) \log(r - m + b). \end{aligned}$$

This isn't simple for V in terms of R, Z (or x, z).

- We can give a simple example of the construction described in the Corollary (Section 4.1) by again taking $V = R^2 - 2Z^2$. With $\Lambda = 2$ for convenience (which leads to Ricci scalar equal to -24), and $Y = y/4$, the metric turns out to be

$$g = \frac{4}{R^4} (dt - ZdY)^2 + \frac{1}{R^2} (dR^2 + dY^2 + dZ^2). \quad (50)$$

This is easily to be seen to be ASD Einstein-Kähler with constant holomorphic sectional curvature and negative Ricci scalar, so it must be the Bergman metric.

- As an example of one-sided type-D Einstein from Section 5, we note that (46) has solutions of the form

$$u = -2 \log(1 + k(x^2 + y^2)) + F(z),$$

with e^F a 2-parameter family of quartic polynomials in z :

$$e^F = c_2(1 + 24\Lambda z^3) + c_3z(1 + 6\Lambda z^3) + 4kz^2,$$

and these include the Schwarzschild-de Sitter metric if $k = 1, c_2 = 0$, and c_3 is negative and related to the mass parameter. This calculation also confirms the accuracy of the expressions for A and B in (46).

References

- [1] ADAMO, TIM; NEWMAN, EZRA T. The Kerr-Newman metric: a review. *Scholarpedia* **9** (2014), no. 10, Article ID 31791. arXiv:1410.6626, doi: 10.4249/scholarpedia.31791. 298
- [2] AKSTEINER, S. Private communication to the author. (2019). 283
- [3] AKSTEINER, STEFFEN; ANDERSSON, LARS. Gravitational instantons and special geometry. *J. Differential Geom.* **128** (2024), no. 3, 928–958. MR4809333 Zbl 1564.53040 arXiv:2112.11863, doi: 10.4310/jdg/1729092451. 283
- [4] BELINSKI, VLADIMIR A.; VERDAGUER, ENRIC. Gravitational solitons. Cambridge Monographs on Mathematical Physics. *Cambridge University Press, Cambridge*, 2001. xiv+258 pp. ISBN: 0-521-80586-4. MR1839019, Zbl 0984.83001, doi: 10.1017/CBO9780511535253. 283

- [5] BERJAWI, SOBHI; FERAPONTOV, EUGENE V.; KRUGLIKOV, BORIS S.; NOVIKOV, VLADIMIR S. Second-order PDEs in 3D with Einstein-Weyl conformal structure. *Ann. Henri Poincaré* **23** (2022), no. 7, 2579–2609. MR4439924, Zbl 1492.35333, arXiv:2104.02716, doi: 10.1007/s00023-021-01140-2. With additional private communication from Jenya Ferapontov. 284, 296
- [6] BIQUARD, OLIVIER; GAUDUCHON, PAUL. On toric Hermitian ALF gravitational instantons. *Comm. Math. Phys.* **399** (2023), no. 1, 389–422. MR4567377, Zbl 1512.53031, arXiv:2112.12711, doi: 10.1007/s00220-022-04562-z. 283
- [7] CHEN, YU; TEO, EDWARD. A new AF gravitational instanton. *Phys. Lett. B* **703** (2011), no. 3, 359–362. MR2831866, arXiv:1107.0763, doi: 10.1016/j.physletb.2011.07.076. 283
- [8] DUNAJSKI, MACIEJ; TOD, PAUL. Four-dimensional metrics conformal to Kähler. *Math. Proc. Cambridge Philos. Soc.* **148** (2010), no. 3, 485–503. MR2609304, Zbl 1188.53078, arXiv:0901.2261, doi: 10.1017/S030500410999048X. 288
- [9] GIBBONS, GARY W.; HAWKING, STEPHEN W. Classification of gravitational instanton symmetries. *Comm. Math. Phys.* **66** (1979), no. 3, 291–310. MR0535152, doi: 10.1007/BF01197189. 297
- [10] HUGGETT, STEPHEN A.; TOD, KENNETH P. An introduction to twistor theory. Second edition. London Mathematical Society Student Texts, 4. *Cambridge University Press, Cambridge*, 1994. xii + 178 pp. ISBN: 0-521-45157-4; 0-521-45689-4. MR1292461, Zbl 0803.00012, doi: 10.1017/CBO9780511624018. 284
- [11] NEWMAN, EZRA T.; PENROSE, ROGER. Spin-coefficient formalism. *Scholarpedia* **4** (2009), no. 6, Article ID 31791. doi: 10.4249/scholarpedia.7445. 284
- [12] PENROSE, ROGER; RINDLER, WOLFGANG. Spinors and space-time. Vol. 1. Two-spinor calculus and relativistic fields. Cambridge Monographs on Mathematical Physics. *Cambridge University Press, Cambridge*, 1984. x+458 pp. ISBN: 0-521-24527-3. MR0776784, Zbl 0538.53024.
- PENROSE, ROGER; RINDLER, WOLFGANG. Spinors and space-time. Vol. 2. Spinor and twistor methods in space-time geometry. Cambridge Monographs on Mathematical Physics. *Cambridge University Press, Cambridge*, 1986. x+501 pp. ISBN: 0-521-25267-9. MR0838301, Zbl 0591.53002. 283, 284, 285
- [13] PONTECORVO, MASSIMILIANO. On twistor spaces of anti-self-dual Hermitian surfaces. *Trans. Amer. Math. Soc.* **331** (1992), no. 2, 653–661. MR1050087, Zbl 0754.53053, doi: 10.1090/S0002-9947-1992-1050087-0.
- [14] PRZANOWSKI, MACIEJ. Killing vector fields in self-dual, Euclidean Einstein spaces with $\Lambda \neq 0$. *J. Math. Phys.* **32** (1991), no. 4, 1004–1010. MR1097787, Zbl 0737.53073, doi: 10.1063/1.529375. 282, 283, 284, 294
- [15] PRZANOWSKI, MACIEJ. Notes on the Einstein equations for locally Hermite-Einstein spaces. *Acta Phys. Polon. B* **22** (1991), no. 4, 357–365. MR1128156. 282, 284, 295
- [16] PRZANOWSKI, MACIEJ; BAKA, B. One-sided type-D gravitational instantons. *Gen. Relativity Gravitation* **16** (1984), no. 9, 797–803. MR0759976, doi: 10.1007/BF00762933. 282, 283, 284, 288
- [17] PRZANOWSKI, MACIEJ; BIALECKI, S. Lie-Bäcklund transformation and gravitational instantons. *Acta Phys. Polon. B* **18** (1987), no. 10, 879–889. <https://www.actaphys.uj.edu.pl/R/18/10/879>. 282, 283, 284, 288
- [18] TOD, KENNETH P. The $SU(\infty)$ -Toda field equation and special four-dimensional metrics. *Geometry and physics* (Aarhus, 1995), 307–312. Lecture Notes in Pure and Applied Mathematics, 184. *Dekker, New York*, 1997. MR1423177, Zbl 0876.53026. 283, 284, 294
- [19] TOD, KENNETH P. Scalar-flat Kähler and hyper-Kähler metrics from Painlevé-III. *Classical Quantum Gravity* **12** (1995), no. 6, 1535–1547. MR1344288, Zbl 0828.53061, doi: 10.1088/0264-9381/12/6/018. 297
- [20] TOD, PAUL. A note on Riemannian anti-self-dual Einstein metrics with symmetry. Preprint, 2006. arXiv:hep-th/0609071. 283, 284, 294

- [21] WALKER, MARTIN; PENROSE, ROGER. On quadratic first integrals of the geodesic equations for type {22} spacetimes. *Comm. Math. Phys.* **18** (1970), 265–274. MR0272351, Zbl 0197.26404, doi: 10.1007/BF01649445. 283, 284, 295
- [22] Ward, Richard S. Einstein-Weyl spaces and $SU(\infty)$ Toda fields. *Classical Quantum Gravity* **7** (1990), no. 4, L95–L98. MR1045295, Zbl 0687.53044 doi: 10.1088/0264-9381/7/4/003. 282, 283, 284, 291, 292

(Paul Tod) MATHEMATICAL INSTITUTE, OXFORD UNIVERSITY, OXFORD OX1 2JD, UNITED KINGDOM
paul.tod@sjc.ox.ac.uk

This paper is available via <http://nyjm.albany.edu/j/2026/32-12.html>.