

Multiplier modules of Hilbert C*-modules revisited

Michael Frank

Dedicated to David Royal Larson

ABSTRACT. The theory of multiplier modules of Hilbert C*-modules is reconsidered to obtain more properties of these special Hilbert C*-modules. The property of a Hilbert C*-module to be a multiplier C*-module is shown to be invariant with respect to consideration as a left or right Hilbert C*-module in the sense of imprimitivity bimodules in strong Morita equivalence theory. The interrelation of the C*-algebras of “compact” operators, the Banach algebras of bounded module operators and the Banach spaces of bounded module operators of a Hilbert C*-module to its C*-dual Banach C*-module, are characterized for pairs of Hilbert C*-modules and their respective multiplier modules. The structures on the latter are always isometrically embedded into the respective structures on the former. Examples are given for which continuation of these kinds of bounded module operators from the initial Hilbert C*-module to its multiplier module fails. However, existing continuations turn out to be always unique. Similarly, bounded modular functionals from both kinds of Hilbert C*-modules to their respective C*-algebras of coefficients are compared, and eventually existing continuations are shown to be unique.

CONTENTS

1. Introduction	222
2. On multiplier modules	226
3. On modular operators and functionals	230
References	235

Multiplier modules of (full) Hilbert C*-modules appeared in the literature during investigations of extensions of Hilbert C*-modules in terms of short exact sequences. There are several equivalent approaches to the subject, e.g. the double centralizer type approach [16, 46, 13, 14, 12, 15] or a pure Hilbert C*-module approach [5, 6], cf. [17, 47] for the link between them. We follow the approach in [5, 6]. The goal was to generalize the extension theory of C*-algebras

Received May 15, 2025.

2010 *Mathematics Subject Classification.* Primary 46L08; Secondary 46L05, 46H10, 47B48.

Key words and phrases. Hilbert C*-modules; multiplier modules; multiplier algebras; bounded modular maps and operators.

to the context of full Hilbert C^* -modules. The notion of a multiplier module of a full Hilbert A -module over a C^* -algebra A is justified by [5, Thm. 1.2]: It is the largest essential extension of X up to unitary modular isomorphism of Hilbert $M(A)$ -modules, where $M(A)$ is the multiplier C^* -algebra of A . For details see section 2. In the sequel to [5, 6], large parts of the extension theory of Hilbert C^* -modules have been described, e.g. in [4, 10, 1, 32, 3, 28, 24]. Recently, J. Taylor described local multiplier modules over local multiplier algebras in case of imprimitivity modules between (non-unital, in general) strongly Morita equivalent C^* -algebras, cf. [48, 49].

The aim of the present note is to fill in some missing facts from the point of view of classical Hilbert C^* -module theory and to describe pairs of Hilbert C^* -modules and their multiplier modules from the point of view of their common and partially even differing properties. We get some results that contradict habits and opinions from Hilbert and Banach space theory. So, some new examples complement the existing points of view on the theory of Hilbert C^* -modules.

Considering multiplier modules as full left Hilbert C^* -modules X over a C^* -algebra A , we obtain that they are at the same time full right Hilbert C^* -modules and multiplier modules over the respective C^* -algebras $K_A(X)$. This reminds one of special imprimitivity bimodules, however no new type of Morita equivalence can be derived. Considering pairs $(X, M(X))$ of Hilbert C^* -modules and their multiplier modules, we consider the interrelations of comparable types of operator algebras of bounded module operators over them and continuation problems of operators from X to $M(X)$, the same for bounded modular functionals. So, for “compact” operator algebras $K_A(X)$ is $*$ -isometrically embedded into $K_{M(A)}(M(X))$, but the former might be smaller than the latter one. Furthermore, the Banach algebra of bounded module operators $\text{End}_{M(A)}(M(X))$ is isometrically embedded into $\text{End}_A(X)$, as well as the Banach space $\text{End}_{M(A)}(M(X), M(X)')$ into $\text{End}_A(X, X')$, and again the former might be smaller than the latter one. Consequently, not any operator of the smaller spaces on X can be continued to a bounded operator on the larger spaces on $M(X)$ obeying strict convergence. If such a continuation exists, it is unique. The same picture can be obtained for bounded modular functionals on $M(X)$ and on X , respectively, so we found examples of pairs of Hilbert C^* -modules $(X, M(X))$ with $X^\perp = \{0\}$ for which no general Hahn-Banach type theorem can be obtained. Remarkably, there does not exist any non-trivial bounded module map from $M(X)$ to $M(A)$ vanishing on X .

1. Introduction

We denote C^* -algebras by A, B . In case a C^* -algebra A is non-unital, general C^* -multiplier theory provides us with some derived structures like multiplier algebras $M(A)$, left and right multiplier algebras $LM(A)$ and $RM(A)$, resp., and quasi-multiplier spaces $QM(A)$. To calculate these linear spaces, any faithful

$*$ -representation of A on a Hilbert space H can be used. The calculation environment is the von Neumann algebra generated by the faithfully $*$ -represented C^* -algebra A , or $B(H)$ itself, cf. [38, Ch. 3], [42]. The unital C^* -algebra $M(A)$ is defined as

$$M(A) = \{m \in B(H) : ma, am \in A \text{ for any } a \in A\}.$$

The Banach algebras $LM(A) = RM(A)^*$ can be defined as

$$LM(A) = RM(A)^* = \{m \in B(H) : ma \in A \text{ for any } a \in A\},$$

whereas the involutive Banach space $QM(A)$ can be obtained as

$$QM(A) = \{m \in B(H) : bma \in A \text{ for any } a, b \in A\}.$$

Inventing different kinds of strict topologies, intrinsic characterizations of these structures as certain topological completions of A on bounded sets of A are available. For comprehensive sources, we refer to the book by P. Ara and M. Matthieu [2] and to [35, 36, 8, 18]. Note, that multiplier algebras might admit an entire lattice of non-unital, two-sided, non-isomorphic ideals A_α such that $M(A_\alpha) = M(A_\beta)$ for any two of them, cf. [27]. Also, either $M(A) = LM(A)$ and $M(A) = QM(A)$ at the same time, or $LM(A)$ is strictly larger than $M(A)$ and $QM(A)$ is strictly larger than $LM(A)$, cf. [9, Cor. 4.18].

We would like to consider Hilbert C^* -modules over (non-unital, in general) C^* -algebras and, in particular, their multiplier modules and related structures. By convention, all Hilbert C^* -modules are right C^* -modules at the first glance. However, for full Hilbert A -modules one can obtain an operator-valued inner product turning them into full left Hilbert $K_A(X)$ -modules, and vice versa. So, the point of view decides which of the two Hilbert C^* -module structures is primary and which is secondary. This kind of consideration will be used frequently in the present paper.

A pre-Hilbert A -module over a C^* -algebra A is an A -module X equipped with an A -valued, non-degenerate mapping $\langle \cdot, \cdot \rangle : X \times X \rightarrow A$ being conjugate- A -linear in the first argument and A -linear in the second one, and satisfying $\langle x, x \rangle \geq 0$ for every $x \in X$. The map $\langle \cdot, \cdot \rangle$ is called the A -valued inner product on X . A pre-Hilbert A -module $\{X, \langle \cdot, \cdot \rangle\}$ is Hilbert if and only if it is complete with respect to the norm $\| \cdot \| = \| \langle \cdot, \cdot \rangle \|_A^{1/2}$. We always assume that the complex linear structures of A and X are compatible. A Hilbert A -module $\{X, \langle \cdot, \cdot \rangle\}$ over a C^* -algebra A is full if the norm-closed A -linear hull $\langle X, X \rangle$ of the range of the inner product coincides with A . Two (full) Hilbert A -modules $\{X, \langle \cdot, \cdot \rangle_X\}$ and $\{Y, \langle \cdot, \cdot \rangle_Y\}$ over a fixed C^* -algebra A are unitarily equivalent (or unitarily isomorphic) iff there exists a bounded invertible adjointable map $T : X \rightarrow Y$ such that $\langle \cdot, \cdot \rangle_X = \langle T(\cdot), T(\cdot) \rangle_Y$ on X . The A -dual Banach A -module X' of a Hilbert A -module X is defined as the set of all bounded A -linear maps from X into A . It might not be a Hilbert A -module itself, cf. [18]. But, X is always canonically isometrically embedded into X' as a Banach A -submodule via the identification of $x \in X$ with $\langle x, \cdot \rangle \in X'$. Note, that two A -valued inner products on a Hilbert A -module X inducing equivalent norms on X might not be unitarily isomorphic,

cf. [9, 18]. Thus, full Hilbert C^* -modules are always a triple of the C^* -algebra A of coefficients, the Banach A -module X and the A -valued inner product on X . We omit the explicit reference to the C^* -valued inner product in situations where its definition formula is standard or only its existence is important.

We are interested in properties of sets of bounded A -linear operators between Banach and Hilbert C^* -modules X . The set $\text{End}_A(X)$ of all bounded module operators on Hilbert A -modules X forms a Banach algebra, whereas the set $\text{End}_A^*(X)$ of all bounded module operators which possess an adjoint operator inside $\text{End}_A(X)$ has the structure of a unital C^* -algebra. Note, that these two sets do not coincide in general, cf. [40, 18]. An important subset of $\text{End}_A^*(X)$ is the set $K_A(X)$ of “compact” operators, which is defined as the norm-closure of the set of all finite linear combinations of elementary operators

$$\{\theta_{a,b} \in \text{End}_A(X) : a, b \in X, \theta_{a,b}(c) = b\langle a, c \rangle \text{ for every } c \in X\}.$$

It is a C^* -subalgebra and a two-sided ideal of $\text{End}_A^*(X)$. In contrast to the well-known situation for Hilbert spaces, the properties of an operator of being “compact” or possessing an adjoint depend strongly on the choice of the A -valued inner product on X , i.e., these properties are not invariant for unitarily non-isomorphic C^* -valued inner products on X inducing equivalent norms, cf. [18].

We postpone a detailed introduction to multiplier modules of Hilbert C^* -modules to the next section. Our standard references to Hilbert C^* -module theory are [34, 44, 50].

Searching for intrinsic characterizations of Hilbert C^* -modules, especially over non-unital C^* -algebras, the notions of orthonormal bases and of frames for Hilbert spaces were rediscovered in the modular context of Hilbert C^* -modules by D. R. Larson and the author during 2018-2022, [21, 22, 23] and cf. [25]. The new theory started with the norm-convergent case. Remarkably, there was a shift in significance towards modular frames, since not all Hilbert C^* -modules admit orthogonal bases. Also, certain classes of Hilbert C^* -modules do not possess modular frames, however the most usual classes of Hilbert C^* -modules do. Theory and applications have been developed and extended since then. The type of convergence of the defining series in the modular context has been widened to strict, weak, weak* types or algebraic order type of convergences in cases.

In 2017, Lj. Arambašić and D. Bakić have made a significant progress in the understanding of Hilbert C^* -modules over non-unital C^* -algebras. They used the strict completion picture to multiplier modules of X and introduced so called outer frames of the multiplier modules $M(X)$ to extend the available sets of norm-convergent or strictly convergent modular frames of the related initial Hilbert C^* -modules X . It turned out that all outer and inner frames of countably generated or algebraically finitely generated Hilbert C^* -modules X in the sense of strict convergence can be characterized by surjections of either $\text{End}_A^*(l_2(A), X)$ or $\text{End}_A^*(A^N, X)$ for some $N \in \mathbb{N}$, resp., [3, Thm. 3.18,

Thm. 3.19, Prop. 3.22, Prop. 3.23]. By the way, the notion of countably generatedness of Hilbert C*-modules over non-unital C*-algebras has been formulated more precisely. Later on, M. Naroei Irani and A. Nazari made steps towards inner and outer modular woven frames, cf. [39].

We are partially interested in strong Morita equivalence of C*-algebras and in imprimitivity bimodules realizing those equivalences. A C*-correspondence from a C*-algebra B to a C*-algebra A (or an B - A C*-correspondence) is a Hilbert module X over A together with a nondegenerate $*$ -homomorphism from B to $\text{End}_A^*(X)$, which introduces a left B -module structure on X . For any B - A C*-correspondence X , the actions extend uniquely to multiplier algebras of B and of A , so that we may always treat X as a $M(B)$ - $M(A)$ C*-correspondence. To relate it to the multiplier $M(B)$ - $M(A)$ bimodule $M(X)$, let us note that whenever Y is a Hilbert A -module, then $\text{End}_A^*(Y, X)$ is naturally a C*-correspondence from $M(B)$ to $\text{End}_A^*(Y)$. In particular, $\text{End}_A^*(Y, X)$ contains $\text{K}_A(Y, X)$ as a sub-C*-correspondence. An B - A imprimitivity bimodule is a full B - A C*-correspondence X which is also a full left Hilbert B -module and the two B - resp. A -valued inner products satisfy the equality $\langle x, y \rangle_B z = x \langle y, z \rangle_A$ for an $x, y, z \in X$. In particular, for imprimitivity bimodules X , X is a full B - A C*-correspondence such that the left action $\phi : B \rightarrow \text{End}_A^*(X)$ restricts to a $*$ -isomorphism of B with $\text{K}_A(X)$. Two C*-algebras B and A are said to be strongly Morita equivalent iff there exists an imprimitivity bimodule (or a B - A imprimitivity bimodule) connecting them. It is a real equivalence relation, so one can form classes of strongly Morita equivalent C*-algebras sharing important properties among their elements. The interested reader is referred to [44, 16, 13, 14, 12, 15] to learn more about the theory of strong Morita equivalence of C*-algebras and about C*-correspondences or imprimitivity bimodules. If X and Y are, respectively, B - A and C - B imprimitivity bimodules, then their algebraic tensor product $X \odot_B Y$ is a C - A imprimitivity bimodule, and one can define two inner products on $X \odot_B Y$ by

$$\langle x \otimes z, y \otimes t \rangle_A = \langle \langle y, x \rangle_B z, t \rangle_A, \quad \langle x \otimes z, y \otimes t \rangle_C = \langle x, y \langle t, z \rangle_B \rangle_C$$

for any $x, y \in X$, $z, t \in Y$. This way $X \odot_B Y$ becomes an C - A imprimitivity bimodule. Remarkably, the set of all B - A imprimitivity bimodules connecting two given strongly Morita equivalent C*-algebras A and B can be quite varied. Using the defined tensor product operation, the problem boils down to the consideration of the Picard group $\text{Pic}(A)$ of one C*-algebra A of unitary isomorphism classes of A - A imprimitivity bimodules for an arbitrarily selected element A of a given class of strongly Morita equivalent to A C*-algebras. Note, that the inverse of an imprimitivity bimodule X in $\text{Pic}(A)$ is its dual module \hat{X} . If Y is an B - A imprimitivity bimodule, then the map $X \rightarrow \hat{Y} \otimes_A X \otimes_A Y$ induces an isomorphism of $\text{Pic}(A)$ and $\text{Pic}(B)$, and [7, Theorem 1.2] shows that the Picard groups are stable isomorphism invariants. For a non-trivial example, compare [31] for the Picard groups of irrational rotation algebras. For a comprehensive account, see [11, sec. 6-7].

2. On multiplier modules

Let us show a way to define multiplier modules $M(X)$ of given (full) Hilbert A -modules X as certain related Hilbert $M(A)$ -modules, and let us provide some important properties of them. We start with a class of extensions of a given full Hilbert A -module X as defined in [5]:

Definition 2.1. (cf. [5, Def. 1.1]) Let X be a full Hilbert C^* -module over a given (non-unital, in general) C^* -algebra A . An extension of X is a triple (Y, B, Φ) such that

- (i) B is a C^* -algebra containing A as a two-sided norm-closed ideal.
- (ii) Y is a Hilbert B -module.
- (iii) $\Phi : X \rightarrow Y$ is a bounded module map satisfying $\langle \Phi(x), \Phi(y) \rangle = \langle x, y \rangle$ for any $x, y \in X$.
- (iv) $\text{Im}(\Phi) = YA = \{za : z \in Y, a \in A\} = \{x \in Y : \langle x, x \rangle \in A\}$ (by the Hewitt-Cohen factorization theorem, [43, Thm. 4.1], [44, Prop. 2.31], [26, Thm. 23.22]).

The triple (Y, B, Φ) is an essential extension of X if A is an essential ideal of B .

Note, that Φ is an A -linear isometry of Hilbert A -modules and, hence, in case of a surjective mapping a unitary map, preserving A -valued inner products up to unitary equivalence, cf. [33], [17, Thm. 5], [47, Thm. 1.1]. So, Y and $\Phi(Y)$ are unitarily equivalent Hilbert C^* -modules. In the sequel we consider the C^* -algebras A and B as a Hilbert A -module and as a Hilbert B -module over itself, respectively, setting $\langle a, b \rangle = a^*b$ for any two C^* -algebra elements a, b . Then A and B are $*$ -isometrically isomorphic to the C^* -algebras $K_A(A)$ and $K_B(B)$, respectively. By [50, Prop. 2.2.16], these $*$ -isomorphisms extend to $*$ -isomorphisms of $M(A)$ and of $M(B)$ with the C^* -algebras $\text{End}_{M(A)}^*(A)$ and $\text{End}_{M(B)}^*(B)$, respectively. We shall use these identifications freely.

Definition 2.2. Let X be a (not necessarily full) Hilbert C^* -module over a given (non-unital, in general) C^* -algebra A . Denote by $M(X)$ the set of all adjointable maps from A to X , i.e., $M(X) = \text{End}_A^*(A, X)$. Obviously, $M(X)$ is a Hilbert $M(A)$ -module with the $M(A)$ -valued inner product $\langle z_1, z_2 \rangle = z_1^*z_2$ for $z_1, z_2 \in M(X)$. The resulting Hilbert $M(A)$ -module norm coincides with the operator norm on $M(X)$. We call $M(X)$ the multiplier module of X .

In [5] $M(X)$ is shown to be an essential extension of X identifying X and $K_A(A, X)$ as the subset $\{za : z \in M(X), a \in A\}$ isometrically. In fact, $M(X)$ is the largest essential extension of X , because $M(A)$ is the largest essential extension of A containing $*$ -isomorphic copies of all C^* -algebras B which contain $*$ -isomorphic copies of A as an essential ideal, [5, Thm. 1.2]. This justifies the point of view on $M(X)$ as a Hilbert C^* -module analog of the multiplier algebra in C^* -theory. For a given Hilbert A -module $\{X, \langle ., . \rangle_X\}$ the respective multiplier module $M(X)$ is unique up to unitary isomorphism of Hilbert $M(A)$ -modules.

Let us explain the definition of multiplier modules $M(X)$ for non-full Hilbert A -modules X . $\langle X, X \rangle$ is a two-sided norm-closed ideal of A . If X is a non-full

Hilbert A -module form the derived full Hilbert A -module X_c by adding a copy of A as an orthogonal direct summand to X , $X_c = X \oplus A$, $\langle \cdot, \cdot \rangle_c = \langle \cdot, \cdot \rangle_X + \langle \cdot, \cdot \rangle_A$. Then construct its multiplier module $M(X_c)$. Using the mapping picture of Definition 2.2 of $M(X_c)$, we see that $M(X_1 \oplus X_2) = M(X_1) \oplus M(X_2)$ using modular projection operators onto each of the orthogonal summands. So, we have an orthogonal decomposition $M(X_c) = M(X) \oplus M(A)$, taking the first orthogonal direct summand and Hilbert $M(A)$ -module as the definition of the multiplier module $M(X)$ of X . Clearly, the (non-unital, in general) C*-algebra $\langle X, X \rangle$ is a two-sided norm-closed ideal of A , and the C*-algebra $\langle M(X), M(X) \rangle$ is a two-sided norm-closed ideal of $M(A)$ containing $\langle X, X \rangle$. However, the latter might be non-unital, so we should be more careful with their C*-algebraic interrelations. In general, $\langle X, X \rangle$ is a two-sided norm-closed ideal in $\langle M(X), M(X) \rangle$. The uniqueness results for the pairings $(X, M(X))$ are preserved. It is important to realize that $M(X)$ might depend on the choice of the C*-algebra acting on X , e.g., for $X = A$ with A a non-unital C*-algebra we have $M_A(A) = \text{End}_A^*(A, A) = M(A)$, but $M_{M(A)}(A) = \text{End}_{M(A)}^*(M(A), A) = A$. Last but not least, if the C*-algebra A is non-unital then for $X = l_2(A)$ the multiplier module $M(l_2(A))$ equals the set

$$\{\{x_n\}_{n \in \mathbb{N}} : x_n \in M(A), \Sigma_n x_n^* x_n \text{ converges strictly w.r.t. } A \text{ in } M(A)\},$$

cf. [5, Thm. 2.1]. This gives a good non-trivial example, in particular, for certain non- σ -unital C*-algebras A , cf. [5, Ex. 2.2]. In particular, $l_2(M(A))$ is smaller than $M(l_2(A))$, generally speaking. For $A = K(l_2)$ and $X = l_2(A)$ one obtains $M(A) = B(l_2)$ and $M(X) = l_2(M(A))' \equiv M(A) = B(l_2)$, the $M(A)$ -dual Banach $M(A)$ -module of $l_2(M(A))$, where $l_2(M(A))'$ equals the set

$$\left\{ m = \{m_i\}_{i=1}^n : m_i \in M(A), \left\| \sum_{i=1}^n m_i^* m_i \right\| \leq K_m < \infty \text{ for any } n \in \mathbb{N} \right\}$$

– a self-dual Hilbert W^* -module over $M(A) = B(l_2)$, because $A = K(l_2)$ is stable, cf. [5, Ex. 2.2] and [40].

Proposition 2.3.

- (i) *For a given pair of C*-algebras $(A, M(A))$, let X_1, X_2 be two full Hilbert C*-modules over A such that their multiplier modules $M(X_1), M(X_2)$ are unitarily isomorphic as Hilbert $M(A)$ -modules. Then X_1 and X_2 are unitarily isomorphic as Hilbert A -modules to $M(X_1)A \equiv M(X_2)A$. So, the pairings $(X, M(X))$ are bound to each other for given C*-algebras $(A, M(A))$ up to unitary equivalence.*
- (ii) *Suppose we have two non- $*$ -isomorphic C*-algebras A_1 and A_2 such that they admit the same multiplier C*-algebra $M(A)$. Let X_1 be a full Hilbert A_1 -module and X_2 a full Hilbert A_2 -module such that $M(X_1)$ and $M(X_2)$ are unitarily isomorphic as Hilbert $M(A)$ -modules. Then X_1 is not unitarily isomorphic to X_2 as a Hilbert $M(A)$ -module.*

The first assertion follows from (iv) of Definition 2.1 combining Φ_2 with Φ_1^{-1} with respect to the set $\{za : z \in M(X), a \in A\}$ in the largest essential extension $M(X)$, cf. [45, Prop. 1.4]. The second assertion can be illustrated by example (see below), however $M(X_1)A_1$ is obviously not isometrically isomorphic to $M(X_2)A_2$ by supposition.

Example 2.4. Let H be an infinite-dimensional Hilbert space, and $K(H)$ and $B(H)$ the sets of compact linear operators and of bounded linear operators on it, respectively. Consider the three C^* -algebras

$$A_2 = \begin{pmatrix} K(H) & 0 \\ 0 & K(H) \end{pmatrix}, \quad A_1 = \begin{pmatrix} K(H) & 0 \\ 0 & B(H) \end{pmatrix},$$

$$M(A_1) = M(A_2) = \begin{pmatrix} B(H) & 0 \\ 0 & B(H) \end{pmatrix}.$$

Let us describe these C^* -algebras as (extended) Hilbert C^* -modules over A_1 and find their multiplier modules with respect to A_1 .

$$X_2 = \begin{pmatrix} K(H) & 0 \\ 0 & K(H) \end{pmatrix} \oplus \begin{pmatrix} K(H) & 0 \\ 0 & B(H) \end{pmatrix},$$

$$X_1 = \begin{pmatrix} K(H) & 0 \\ 0 & B(H) \end{pmatrix} \oplus \begin{pmatrix} K(H) & 0 \\ 0 & B(H) \end{pmatrix},$$

$$M(X_2) = \begin{pmatrix} B(H) & 0 \\ 0 & K(H) \end{pmatrix} \oplus \begin{pmatrix} B(H) & 0 \\ 0 & B(H) \end{pmatrix}.$$

$$M(X_1) = \begin{pmatrix} B(H) & 0 \\ 0 & B(H) \end{pmatrix} \oplus \begin{pmatrix} B(H) & 0 \\ 0 & B(H) \end{pmatrix}.$$

By [45, Prop. 1.1], both X_1 and X_2 admit a canonical isometric modular embedding into $M(X_1)$ and into $M(X_2)$, respectively, as Hilbert $M(A_1)$ -submodules. However, the multiplier module of A_2 with respect to A_1 , $M_{A_1}(A_2)$, is a non-unital C^* -algebra.

Now, consider X_1 as a full Hilbert A_1 -module and

$$X_3 = \begin{pmatrix} K(H) & 0 \\ 0 & K(H) \end{pmatrix} \oplus \begin{pmatrix} K(H) & 0 \\ 0 & K(H) \end{pmatrix},$$

as a full Hilbert A_2 -module. Then $M(X_1)$ is unitarily isomorphic to $M(X_3)$ as a Hilbert $M(A_1) \equiv M(A_2)$ -module, but X_1 and X_3 are not.

In general, for unital C^* -algebras $A = M(A)$, any Hilbert A -module X is its own multiplier module $M(X) = X$, [5, Remark 1.11]. To see that, an intrinsic topological characterization of multiplier modules for $\langle X, X \rangle \subseteq A$ being an essential ideal of $M(A)$ is useful.

By [5], there exists a suitable variant of a strict topology on multiplier modules $M(X)$: Let A be a C^* -algebra and X be a Hilbert A -module. Let the strict topology on $M(X)$ be induced jointly by the two families of semi-norms $\{\|z \rightarrow za\| : a \in A\}$ and $\{\|\langle z, x \rangle\| : x \in X, \|x\| \leq 1\}$ for $z \in M(X)$. It is a locally convex topology. The multiplier module $M(X)$ turns out to be complete

with respect to this strict topology, and $M(X)$ is the strict completion of X , [5, Thm. 1.8, 1.9]. Moreover, the strict completion is an idempotent operation, i.e., $M_{M(A)}(M_A(X)) = M_A(X)$. Consequently, for unital C*-algebras $A = M(A)$ and Hilbert A -modules X we have $X = M(X)$. The same is true whenever $K_A(X)$ is unital, cf. [5, Cor. 2.9].

For X, Y Hilbert A -modules each operator $T \in \text{End}_A^*(X, Y)$ has an extension $T_M \in \text{End}_{M(A)}^*(M(X), M(Y))$ of the same norm value obtained as the strict continuation of T . Therefore, it is uniquely determined. Moreover, every operator in $\text{End}_{M(A)}^*(M(X), M(Y))$ arises this way, i.e., the C*-algebras $\text{End}_A^*(X, Y)$ and $\text{End}_{M(A)}^*(M(X), M(Y))$ are $*$ -isomorphic, [5, Thm. 2.3]. Since a full Hilbert A -module X is not only a right Hilbert A -module, but also a full left Hilbert $K_A(X)$ -module by the theory of strong Morita equivalence of the C*-algebras A and $K_A(X)$, X can be considered as a C*-correspondence or as an imprimitivity bimodule of these two C*-algebras. By [30], the C*-algebra $\text{End}_A^*(X)$ can be considered as the multiplier algebra of the C*-algebra $K_A(X)$. Consequently, $M(X)$ is a full left Hilbert $M(K_A(X))$ -module. So, for (full) Hilbert $K_A(X)$ -modules X the Hilbert $M(K_A(X))$ -module $M(X)$ can be identified with the (left) multiplier module of X w.r.t. the pairing $(K_A(X), M(K_A(X)))$, too. This makes the property of a (full) Hilbert C*-module to be a multiplier module invariant under the choice of the point of view as a (full) left or right Hilbert C*-module. For similar thoughts, compare with [5, pp. 20-21, (a)-(b)], [16, Props. 13 and 16].

Theorem 2.5. *Let A be a C*-algebra and $M(A)$ be its multiplier algebra. Let X be a full (right) Hilbert A -module and $M(X)$ be its multiplier module, a full (right) Hilbert $M(A)$ -module. Then $M(X)$ is also the full (left) multiplier module of the (left) Hilbert $K_A(X)$ -module X with respect to the pairing of C*-algebras $K_A(X)$ and $M(K_A(X)) = \text{End}_A^*(X) = M(K_{M(A)}(M(X))) = \text{End}_{M(A)}^*(M(X))$, and vice versa.*

Proof. By [5, Thm. 1.8, 1.9], the unit ball of $M(X)$ is complete with respect to the locally convex strict (right) topology induced jointly by the two families of semi-norms $\{\|z \rightarrow za\| : a \in A\}$ and $\{\|\langle z, x \rangle_r\| : x \in X, \|x\| \leq 1\}$ for $z \in M(X)$. Also, the unit ball of X generates the unit ball of $M(X)$ strictly. By [4, Def. 2, Remark 3], the operator strict topology on X defined by the joint family of semi-norms $\{\|x \rightarrow T(x)\| : T \in K_A(X)\}$ and $\{\|x \rightarrow xa\| : a \in A\}$ with $x \in X$ coincides with the strict (right) topology on bounded sets of X and of $M(X)$, and so on unit balls, in particular. Consequently, X is dense in $M(X)$ w.r.t. the operator strict topology on X and on $M(X)$. Note, that the operator strict topology is symmetric for imprimitivity bimodules X in A and $K_A(X)$. By strong Morita equivalence via the imprimitivity bimodule X , we can symmetrically conclude that the unit ball of X is dense w.r.t. the locally convex strict (left) topology induced jointly by the two families of semi-norms $\{\|z \rightarrow T(z)\| : T \in K_A(X)\}$ and $\{\|\langle z, x \rangle_l\| : x \in X, \|x\| \leq 1\}$ for $z \in M(X)$. This gives the argument by [5, Thm. 1.8, 1.9] applied to the operator point of view

on X and on $M(X)$. The $*$ -isomorphisms of the respective operator C^* -algebras are shown in [5, Thm. 2.3]. \square

Remark 2.6. Let us check the idea of a stronger kind of strong Morita equivalence restricting the set to multiplier module imprimativity bimodules. There is a good class of pairwise strongly Morita equivalent C^* -algebras

$$\{\mathbb{C}, M_n(\mathbb{C}), K(H) : n \in \mathbb{N}, H \text{ any infinite-dimensional Hilbert space}\},$$

because it has been well investigated in [4, 5, 24]. One has multiplier imprimativity bimodules whenever the left and/or right Hilbert C^* -module structure of imprimativity bimodules involves $\{\mathbb{C}, M_n(\mathbb{C}) : n \in \mathbb{N}\}$. The C^* -algebras

$$\{K(H) : H \text{ any infinite-dimensional Hilbert space}\}$$

may appear among the strongly Morita equivalent ones at the other end. What about imprimativity bimodules X connecting C^* -algebras of type $K(H)$ for infinite-dimensional Hilbert spaces H ? The minimal requirement to an equivalence relation is that an object has to be equivalent to itself. Take $K(H)$ and a full (left) Hilbert $K(H)$ -module serving as a imprimativity bimodule X of $K(H)$ with itself, i.e., with the structure of a full (right) Hilbert $K(H)$ -module. Then the full multiplier C^* -module $M(X) = B(H)$ of X is a imprimativity bimodule of $B(H)$ with itself and does not belong to the set of $K(H)$ - $K(H)$ imprimativity bimodules any more. So, the set of $K(H)$ - $K(H)$ imprimativity bimodules does not contain any multiplier module, similarly for Hilbert spaces H of pairwise non-isomorphic infinite-dimensional dimensions. The concept does not work.

3. On modular operators and functionals

The aim of this section is the investigation of the Banach algebras of all bounded module maps $\text{End}_A(X)$ and $\text{End}_{M(A)}(M(X))$ and their interrelations, as well as the sets of all bounded module maps X' and $M(X)'$ over pairs of (full) Hilbert A -modules X and their multiplier modules $M(X)$ over $M(A)$. To get non-trivial examples, we need examples of C^* -algebras A such that their multiplier algebras $M(A)$ are strictly smaller than their left/right multiplier algebras. By [9, Cor. 4.18], either $M(A) = LM(A)$ and $M(A) = QM(A)$ at the same time, or $LM(A)$ is strictly larger than $M(A)$ and $QM(A)$ is strictly larger than $LM(A)$. For theory and examples, see the existing literature on different types of multiplier algebras and local multiplier algebras, e.g. [8]. We give a simple example following [35, pp. 165-166].

Recall the C^* -algebras c_0 , c and l_∞ of all complex-valued sequences converging to zero, converging at all and being bounded in norm, respectively. Change the target C^* -algebra \mathbb{C} to the C^* -algebra of all two-by-two valued matrices $M_2(\mathbb{C})$. Consider the C^* -algebra of all $M_2(\mathbb{C})$ -valued sequences with the sequence in the upper left corner converging at all and with the sequences derived from the other three positions being sequences converging to zero. We

write A as a symbol

$$A = \begin{pmatrix} c & c_0 \\ c_0 & c_0 \end{pmatrix}.$$

Then we can find the derived (left/right/two-sided) multiplier algebras / spaces:

$$M(A) = \begin{pmatrix} c & c_0 \\ c_0 & l_\infty \end{pmatrix}, \quad LM(A) = RM(A)^* = \begin{pmatrix} c & l_\infty \\ c_0 & l_\infty \end{pmatrix},$$

$$QM(A) = \begin{pmatrix} c & l_\infty \\ l_\infty & l_\infty \end{pmatrix}.$$

This can be calculated in the W^* -algebra of all bounded $M_2(\mathbb{C})$ -valued sequences. Note, that $QM(A) = LM(A) + RM(A) \neq RM(A) \circ LM(A)$ for this particular example, a non-general situation.

Theorem 3.1. *Let A be a C^* -algebra with multiplier algebra $M(A)$. Let X be a full Hilbert A -module and $M(X)$ be its full multiplier module.*

- (i) *The C^* -algebra $K_A(X)$ of all “compact” operators on X admits a $*$ -isomorphic embedding into the C^* -algebra $K_{M(A)}(M(X))$ of all “compact” operators on $M(X)$. If $X \neq M(X)$ then $K_A(X)$ is smaller than $K_{M(A)}(M(X))$. Nevertheless, their multiplier algebras are $*$ -isomorphic, i.e., $\text{End}_A(X)^* \cong \text{End}_{M(A)}(M(X))^*$. If $X \neq M(X)$ then the embedding is not a surjection.*
- (ii) *There does not exist any bounded $M(A)$ -linear map $T_0 : M(X) \rightarrow M(X)$ such that $T_0 \neq 0$ on $M(X)$, but $T_0 = 0$ on $X \subseteq M(X)$.*
- (iii) *The Banach algebra $\text{End}_{M(A)}(M(X))$ admits an isometric embedding into the Banach algebra $\text{End}_A(X)$ by restricting an element on the domain from $M(X)$ to $X \subseteq M(X)$. If the left multiplier algebra of $K_A(X)$ is larger than the multiplier algebra of it, then $\text{End}_{M(A)}(M(X))$ can be smaller than $\text{End}_A(X)$, i.e., not every bounded module operator on X might admit a bounded module operator continuation on $M(X)$.*

Proof. Since $X \subseteq M(X)$, elementary “compact” operators on X can be extended to $M(X)$ preserving their operator norm by the strict density of X in $M(X)$. The C^* -algebras of “compact” operators on Hilbert C^* -modules are generated linearly by elementary operators w.r.t. the operator norm. So the isometric $*$ -isomorphic embedding of the C^* -algebra $K_A(X)$ into the C^* -algebra $K_{M(A)}(M(X))$ follows. However, the multiplier C^* -algebras $M(K_A(X)) = \text{End}_A(X)^*$ and $M(K_{M(A)}(M(X))) = \text{End}_{M(A)}(M(X))^*$ are always $*$ -isomorphic by [5, Thm. 2.3].

Suppose there exists a bounded $M(A)$ -linear operator T_0 on $M(X)$ such that $T_0 = 0$ on $X \subseteq M(X)$, but $T_0(m) \neq 0$ for some $m \in M(X)$. Let $\{x_\alpha : \alpha \in I\}$ be a net of elements of X converging strictly to m , i.e., the nets $\{x_\alpha a : \alpha \in I\}$ converge to $ma \in X$ in norm for any $a \in A$. Consider the set $\{\langle n, T_0(ma) \rangle : a \in A, n \in M(X)\}$. All these values are equal to zero by supposition. Since $\langle n, T_0(ma) \rangle = \langle n, T_0(m) \rangle a = 0$ for any $a \in A$ and A is an essential ideal of $M(A)$ we conclude $\langle n, T_0(m) \rangle = 0$ for any $n \in M(X)$, forcing $T_0(m) = 0$, a contradiction to our assumption.

Restricting a bounded $M(A)$ -linear operator on $M(X)$ on the domain from $M(X)$ to $X \subseteq M(X)$, one obtains a bounded A -linear operator on X . The norm is preserved since X is strictly dense in $M(X)$. There does not exist any non-trivial bounded module operator on $M(X)$ vanishing on $X \subseteq M(X)$ and the norm is preserved on the subalgebra $\text{End}_{(M(A))}(M(X))^* \equiv \text{End}_A(X)^*$ by [5, Thm. 2.3]. So, the restriction of a non-adjointable operator is again a non-adjointable operator. In case the left multiplier algebra $LM(K_A(X)) = \text{End}_A(X)$ is larger than the multiplier algebra $M(K_A(X)) = \text{End}_A(X)^*$ (see example above), all elements of $LM(K_A(X)) \setminus M(K_A(X))$ are not extendable to bounded module operators on $M(X)$: Indeed, if $A = X$ is a C^* -algebra with $LM(A) \supset M(A)$ then $M(X) = M(A)$ and all elements of $LM(A) \setminus M(A)$ preserve $X \subset M(X)$, but do not preserve $M(X)$, cf. [36, Thm. 1.5, 1.6], [9, Cor. 4.18]. \square

We found that not any bounded module operator on a Hilbert C^* -module might admit a continuation to a bounded module operator of the same operator norm value on its multiplier module. However, if such a continuation (with the same operator norm value) exists, it is unique.

Remark 3.2. We demonstrate by example that the choice of the A -valued inner product on Hilbert A -modules X within the class of A -valued inner products on X inducing equivalent norms on X may lead to other unitarily non-equivalent multiplier $M(A)$ -modules. Return to the example at the beginning of the present section. If the C^* -algebra A defined there is equipped with the standard A -valued inner product as a Hilbert A -module X , then $M(X) = M(A)$. Now, modify this A -valued inner product setting $\langle \cdot, \cdot \rangle_1 := \langle T(\cdot), T(\cdot) \rangle_A$ for

$$T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in LM(A) \setminus M(A),$$

where 0 is the zero sequence and 1 is the identity sequence. Clearly, T is a non-adjointable invertible bounded A -linear operator on A , and $\langle \cdot, \cdot \rangle_1$ is an A -valued inner product on A inducing an equivalent Hilbert module norm on $A = X$. A simple calculation for elements of $M(A)$ inside the C^* -valued inner product $\langle \cdot, \cdot \rangle_1$ yields

$$\left\langle \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \circ \begin{pmatrix} c & c_0 \\ c_0 & l_\infty \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \circ \begin{pmatrix} c & c_0 \\ c_0 & l_\infty \end{pmatrix} \right\rangle_A = \begin{pmatrix} c & c \\ c & l_\infty \end{pmatrix},$$

which is not a subset of $M(A)$. Consequently, the A -valued inner product $\langle \cdot, \cdot \rangle_1$ cannot be extended to $M(A)$, and $M(A)$ is not the multiplier module of the Hilbert A -module $X_1 = \{A, \langle \cdot, \cdot \rangle_1\}$. So, the Banach A -module A of the concrete example does not determine its multiplier module alone, one has to take into account the particular A -valued inner product on it. To calculate the multiplier

module of X_1 , one can use the von Neumann algebra of all bounded $M_2(\mathbb{C})$ -valued sequences as an environment. Then

$$\begin{aligned}
 \langle S(x), a \rangle_A &= \langle x, S^*(a) \rangle_1 \\
 &= \langle T(x), T(S^*(a)) \rangle_X \\
 &= \langle T(x), (T^{-1})^* T^*(T(S^*(a))) \rangle_X \\
 &= \langle T^{-1}T(x), T^*(T(S^*(a))) \rangle_X \\
 &= \langle x, (T^*TS^*)(a) \rangle_X
 \end{aligned}$$

for any $S \in \text{End}_{A,1}^*(X, A)$, any $x \in X$, $a \in A$. Thus, $T^*TS^* \in \text{End}_A^*(A, X) = \{M(A), \langle \cdot, \cdot \rangle_{M(A)}\}$.

In 2022, J. Kaad and M. Skeide published an example of a singular extension of the zero bounded C^* -linear functional on a Hilbert C^* -submodule Y in a Hilbert C^* -module X where the orthogonal complement of Y in X was supposed to be the zero element of X , cf. [29]. The author proved that such phenomena cannot appear for Hilbert C^* -modules over monotone complete C^* -algebras and for maximal one-sided ideals of C^* -algebras, cf. [20]. We should evaluate the pairs $(X, M(X))$ under consideration.

Theorem 3.3. *Let A be a C^* -algebra with multiplier algebra $M(A)$. Let X be a full Hilbert A -module and $M(X)$ be its full multiplier module.*

- (i) *There does not exist any bounded $M(A)$ -linear map $f_0 : M(X) \rightarrow M(A)$ such that $f_0 \neq 0$ on $M(X)$, but $f_0 = 0$ on $X \subseteq M(X)$.*
- (ii) *The Banach $M(A)$ -module $M(X)'_{M(A)}$ admits an isometric modular embedding into the Banach A -module X'_A by restricting an element on the domain from $M(X)$ to $X \subseteq M(X)$. There exist examples such that X'_A is strictly larger than the embedded copy of $M(X)'_{M(A)}$.*

Proof. Suppose there exists a bounded $M(A)$ -linear functional $f_0 : M(X) \rightarrow M(A)$ such that $f_0 = 0$ on $X \subseteq M(X)$, but $f_0(m) \neq 0$ for some $m \in M(X)$. Let $\{x_\alpha : \alpha \in I\}$ be a net of elements of X converging strictly to m , i.e., the nets $\{x_\alpha a : \alpha \in I\}$ converge to $ma \in X$ in norm for any $a \in A$. Consider the set $\{f_0(ma) : a \in A\}$. All these values are equal to zero by supposition. Since $f_0(ma) = f_0(m)a = 0$ for any $a \in A$ and A is an essential ideal of $M(A)$ we conclude $f_0(m) = 0$, a contradiction to our assumption.

Restricting $f \in M(X)'$ to $X \subseteq M(X)$ we obtain a bounded A -linear functional of X' . The norm is preserved, since X is strictly dense in $M(X)$ and there does not exist any non-trivial bounded $M(A)$ -linear functional on $M(X)$ vanishing on $X \subseteq M(X)$. The example in the beginning of the present section can be read as follows: Let A be a C^* -algebra such that $LM(A) \supset M(A)$. Setting $A = X$ we get $X' = LM(A)$ and $M(X)' = M(A)' = M(A)$ since $M(X) = M(X)'$ is self-dual. So $X' \supset M(X)'$ and an A -valued bounded functional in $LM(A) \setminus M(A)$ cannot be continued from $X \subseteq M(X)$ to $M(X)$. \square

Consequently, there does not exist any general Hahn-Banach type theorem for bounded C^* -linear functionals for pairs $(X, M(X))$ of full Hilbert C^* -modules X and their multiplier modules $M(X)$, cf. [19, 20]. However, if a bounded A -linear functional from a full Hilbert A -module to the C^* -algebra A admits a continuation to a bounded $M(A)$ -linear functional from its multiplier module to $M(A)$ of the same norm value, then it is unique.

Theorem 3.4. *Let A be a non-unital C^* -algebra with multiplier algebra $M(A)$. Let X be a full Hilbert A -module and $M(X)$ be its full multiplier module.*

- (i) *There does not exist any bounded $M(A)$ -linear map $T_0 : M(X) \rightarrow M(X)'$ such that $T_0 \neq 0$ on $M(X)$, but $T_0 = 0$ on $X \subseteq M(X)$.*
- (ii) *The Banach space $\text{End}_{M(A)}(M(X), M(X)')$ admits an isometric embedding into the Banach space $\text{End}_A(X, X')$ by restricting an element on the domain from $M(X)$ to $X \subseteq M(X)$. There exist examples such that $\text{End}_A(X, X')$ is strictly larger than the embedded copy of $\text{End}_{M(A)}(M(X), M(X)')$.*

Proof. Assume there exists a bounded $M(A)$ -linear map $T_0 : M(X) \rightarrow M(X)'$ such that $T_0 \neq 0$ on $M(X)$, but $T_0 = 0$ on $X \subseteq M(X)$. Then there exists a non-zero element $m \in M(X)$ such that $T_0(m) \in M(X)'$, $T_0(m) \neq 0$ on $M(X)$, but $T_0(m) = 0$ on $M(X)A = X$. This was excluded by Theorem 3.3, (i), a contradiction.

If we restrict an element $T \in \text{End}_{M(A)}(M(X), M(X)')$ to $M(X)A = X$ we obtain an element $T \in \text{End}_A(X, X')$ of the same operator norm, since X is strictly dense in $M(X)$ and $M(X)A = X'$. The algebraic operations are preserved.

The example in the beginning of the present section shows: Let A be a C^* -algebra such that $LM(A) \supset M(A)$. Setting $A = X$ we get $\text{End}_A(X, X') = QM(K_A(X))$ and $\text{End}_{M(A)}(M(X), M(X)') = QM(K_{M(A)}(M(X)))$, cf. [36, Thm. 1.6]. Since $QM(K_A(X)) \neq LM(K_A(X))$, but $QM(K_{M(A)}(M(X))) = M(K_{M(A)}(M(X))) = M(M(A)) = M(A)$ by [9, Cor. 4.18], the assertion is demonstrated. \square

Continuing Remark 3.2, the quasi-multiplier T^*T of A of the example induces a bounded modular map from $X = A$ to $X' = LM(A)$ that cannot be extended to a bounded modular map of the same norm from $M(X) = M(A)$ to $M(X)' = M(A)$.

Remark 3.5. One natural question is whether multiplier modules might be C^* -reflexive with respect to certain C^* -algebras B with $A \subseteq B \subseteq M(A)$, or not. The background for this consists of Definition 2.1, and a result by W. L. Paschke that the C^* -valued inner product on a full Hilbert C^* -module can be always continued to its C^* -reflexive C^* -bidual Banach C^* -module preserving the isometric embedding of the former into the latter, cf. [41]. One could hope to find intrinsic topological or order-algebraical characterizations of some C^* -reflexive Hilbert C^* -modules, cf. the $l_2(K(l_2))$ -example above in the context of [40]. In general, C^* -reflexivity of Hilbert C^* -modules highly depends on both the inner structures of the C^* -algebra of coefficients and of the Hilbert C^* -module under consideration.

To test this point of view select a non-unital C^* -algebra A with $M(A) = LM(A)$. First, if one considers A -reflexivity of $X = A$ with the standard A -valued inner product then $M_A(X) = M(A)$, $X'_A = M(A)$ and $X''_A = A$, so $M_A(X)$ is not A -reflexive. Furthermore, considering $X = A$ as a Hilbert $M(A)$ -module, $M_{M(A)}(X) = A$, $X'_{M(A)} = M(A)$ and $X''_{M(A)} = M(A)$, so it is not $M(A)$ -reflexive, either. The deeper reason could be that condition (iv) of Definition 2.1 might be violated for the C^* -reflexive Hilbert C^* -module extension of X .

Furthermore, instead of two-sided norm-closed ideals A in their multiplier algebras $M(A)$ (and hence, multiplier modules) one-sided norm-closed ideals in C^* -algebras and their biorthogonal complements in the host C^* -algebra can be considered. Here several notions of density of the respective ideal in the arising C^* -subalgebra of the host C^* -algebra have to be compared, and it is far from obvious that they coincide like in the two-sided norm-closed ideal case, cf. [37]. These considerations provide a wider variety of examples of Hilbert C^* -modules and their multiplier modules.

Acknowledgement

I am grateful to David R. Larson for the years of fruitful collaboration during 1998-2002. David R. Larson from Texas A&M University made outstanding contributions to various mathematics subjects including operator theory, operator algebras and applied harmonic analysis, in particular wavelet and frame theory. As well, he was engaged in organizing the highly impacted annual Great Plains Operator Theory Symposium (GPOTS) during many years.

I thank Bartosz Kwaśniewski, who attracted my attention to the double centralizer type approach to multiplier modules using only Banach C^* -module constructions, and to problems about the equivalence of both these approaches in the case of Hilbert C^* -modules, cf. [13, 14, 12, 17, 47].

References

- [1] AMYARI, MARYAM; CHAKOSHI, MAHNNAZ. Pullback diagram of Hilbert C^* -modules. *Math. Commun.* **16** (2011), no. 2, 569–575. [MR2900777](#), [Zbl 1250.46037](#). 222
- [2] ARA, PERE; MATHIEU, MARTIN. Local multipliers of C^* -Algebras. Springer Monographs in Mathematics. Springer-Verlag, Ltd., London, 2003. xii+319 pp. ISBN:1-85233-237-9. [MR1940428](#), [Zbl 1015.46001](#), doi: [10.1007/978-1-4471-0045-4](#). 223
- [3] ARAMBAŠIĆ, LJILJANA; BAKIĆ, DAMIR. Frames and outer frames for Hilbert C^* -modules. *Linear Multilinear Algebra* **65** (2017), no. 2, 381–431. [MR3577457](#), [Zbl 1367.46048](#), [arXiv:1507.04101](#), doi: [10.1080/03081087.2016.1186588](#). 222, 225
- [4] BAKIĆ, DAMIR. A class of strictly complete Hilbert C^* -modules. Preprint, 2025. 8 pp. https://www.researchgate.net/publication/242688089_A_class_of_strictly_complete_Hilbert_C-modules. 222, 229, 230
- [5] BAKIĆ, DAMIR; GULJAŠ, BORIS. Extensions of Hilbert C^* -modules. *Houston J. Math.* **30** (2004), no. 2, 537–558. [MR2084917](#), [Zbl 1069.46032](#). 221, 222, 226, 227, 228, 229, 230, 231, 232
- [6] BAKIĆ, DAMIR; GULJAŠ, BORIS. Extensions of Hilbert C^* -modules. II. *Glasnik Matematički* **38(58)** (2003), no. 2, 343–359. [MR2052751](#), [Zbl 1057.46046](#), doi: [10.3336/gm.38.2.12](#). 221, 222

- [7] BROWN, LAWRENCE G.; GREEN, PHILIP; RIEFFEL, MARC A. Stable isomorphism and strong Morita equivalence of C^* -algebras. *Pacific J. Math.* **71** (1977), no. 2, 349–363. [MR0463928](#), [Zbl 0362.46043](#), doi: [10.2140/pjm.1977.71.349](#). 225
- [8] BROWN, LAWRENCE G.; MINGO, JAMES A.; SHEN, NIEN-TSU. Quasi-multipliers and embeddings of Hilbert C^* -bimodules. *Canad. J. Math.* **46** (1994), no. 6, 1150–1174. [MR1304338](#), [Zbl 0846.46031](#), doi: [10.4153/CJM-1994-065-5](#). 223, 230
- [9] BROWN, LAWRENCE G. Close hereditary C^* -subalgebras and the strucure of quasi-multipliers. *Proc. Royal Society of Edinburgh Sect A* **147** (2017), no. 2, 263–292. [MR3627950](#), [Zbl 1378.46038](#), [arXiv:1501.07613](#), doi: [10.1017/S0308210516000172](#). 223, 224, 230, 232, 234
- [10] BRÜCKLER, FRANKA MIRIAM. A note on extensions of Hilbert C^* -modules and their morphisms. *Glas. Mat. Ser. III* **39(59)** (2004), no. 2, 313–326. [MR2109673](#), [Zbl 1086.46043](#), doi: [10.3336/gm.39.2.12](#). 222
- [11] BURSZTYN, HENRIQUE; WALDMANN, STEFAN. Completely positive inner products and strong Morita equivalence. *Pacific J. Math.* **222** (2005), no. 2, 201–236. [MR2225070](#), [Zbl 1111.53071](#), [arXiv:math/0309402](#), doi: [10.2140/pjm.2005.222.201](#). 225
- [12] BUSS, ALCIDE; KWAŚNIEWSKI, BARTOSZ; MCKEE, ANDREW; SKALSKI, ADAM. Fourier-Stieltjes category for twisted groupoid actions. Preprint, 2024. [arXiv:2405.15653](#). 221, 225, 235
- [13] DAWS, MATTHEW. Multipliers, self-induced and dual Banach algebras. *Dissertationes Math.* **470** (2010), 62pp. [MR2681109](#), [Zbl 1214.43004](#), [arXiv:1001.1633](#), doi: [10.4064/dm470-0-1](#). 221, 225, 235
- [14] DAWS, MATTHEW. Multipliers of locally compact quantum groups via Hilbert C^* -modules. *J. Lond. Math. Soc. (2)* **84** (2011), no. 2, 385–407. [MR2835336](#), [Zbl 1235.43004](#), [arXiv:1004.0215](#). 221, 225, 235
- [15] DELFÍN, ALFONSO. Representations of C^* -correspondences on pairs of Hilbert spaces. *J. Operator Theory* **92** (2024), no. 1, 167–188. [MR4804331](#), [Zbl 1568.46029](#), [arXiv:2208.14605](#), doi: [10.7900/jot.2022sep02.2431](#). 221, 225
- [16] ECHTERHOFF, SIEGFRIED; RAEURN, IAIN. Multipliers of imprimitivity bimodules and Morita equivalence of crossed products. *Math. Scand.* **76** (1995), no. 2, 289–309. [MR1354585](#), [Zbl 0843.46049](#), doi: [10.7146/math.scand.a-12543](#). 221, 225, 229
- [17] FRANK, MICHAEL. A multiplier approach to the Lance–Blecher theorem. *Z. Anal. Anwendungen* **16** (1997), no. 3, 565–573. [MR1472718](#), [Zbl 0881.46040](#), [arXiv:funct-an/9701001](#), doi: [10.4171/zaa/778](#). 221, 226, 235
- [18] FRANK, MICHAEL. Geometrical aspects of Hilbert C^* -modules. *Positivity* **3** (1999), no. 3, 215–243. [MR1708656](#), [Zbl 0945.46043](#), doi: [10.1023/A:1009729204027](#). 223, 224
- [19] FRANK, MICHAEL. On Hahn–Banach type theorems for Hilbert C^* -modules. *Internat. J. Math.* **13** (2002), no. 7, 675–693. [MR1921507](#), [Zbl 1059.46035](#), [arXiv:funct-an/9609003](#), doi: [10.1142/S0129167X02001356](#). 234
- [20] FRANK, MICHAEL. Regularity results for classes of Hilbert C^* -modules with respect to special bounded modular functionals. *Ann. Funct. Anal.* **15** (2024), no. 2, Paper No. 19, 18 pp. [MR4707465](#), [Zbl 1546.46050](#), [arXiv:2207.13164](#), doi: [10.1007/s43034-024-00320-5](#). 233, 234
- [21] FRANK, MICHAEL; LARSON, DAVID R. A module frame concept for Hilbert C^* -modules. *The functional and harmonic analysis of wavelets and frames* (San Antonio, TX, 1999), 207–233. *Contemporary Mathematics*, 247. *American Mathematical Society, Providence, RI, 1999*. ISBN:0-8218-1957-7. [MR1738091](#), [Zbl 0949.46027](#), [arXiv:math/0011184](#). 224
- [22] FRANK, MICHAEL; LARSON, DAVID R. Modular frames for Hilbert C^* -modules and symmetric approximation of frames. *International Symposium on Optical Science and Technology* (San Diego, 2000). *Wavelet Applications in Signal and Image Processing, VIII. Proceedings of the Society of Photo-Optical Instrumentation Engineers (SPIE)* **4119** (2000), 325–336. [arXiv:math/0010115](#), doi: [10.1117/12.408617](#). 224

- [23] FRANK, MICHAEL; LARSON, DAVID R. Frames in Hilbert C^* -modules and C^* -algebras. *J. Operator Theory* **48** (2002), no. 2, 273–314. [MR1938798](#), [Zbl 1029.46087](#), [arXiv:math/0010189](#). 224
- [24] GULJAŠ, BORIS. Hilbert C^* -modules in which all relatively strictly closed submodules are complemented. *Glas. Mat. Ser. III* **56(76)** (2021), no. 2, 343–374. [MR4400848](#), [Zbl 1503.46047](#). 222, 230
- [25] HAN, DEGUANG; JING, WU; LARSON, DAVID R.; MOHAPATRA, RAM N.. Riesz bases and their dual modular frames in Hilbert C^* -modules. *J. Math. Anal. Appl.* **343** (2008), no. 1, 246–256. [MR2412125](#), [Zbl 1185.46040](#), doi: [10.1016/j.jmaa.2008.01.013](#). 224
- [26] HEWITT, EDWIN; ROSS, KENNETH A. Abstract harmonic analysis. II: Structure and analysis for compact groups. Analysis on compact abelian groups. Die Grundlehren der mathematischen Wissenschaften, Band 152. *Springer-Verlag, New York-Berlin*, 1970. ix+771 pp. [MR0262773](#), [Zbl 0830.43001](#), doi: [10.1007/978-3-662-26755-4](#). 226
- [27] HINES, TAYLOR; WALSBERG, ERIK. Nontrivially Noetherian C^* -algebras. *Math. Scand.* **111** (2012), no. 1, 135–146. [MR3001364](#), [Zbl 1272.46050](#), doi: [10.7146/math.scand.a-15219](#). 223
- [28] JINGMING, ZHU. Geometric description of multiplier modules for Hilbert C^* -modules in simple cases. *Ann. Funct. Anal.* **8** (2017), no. 1, 51–62. [MR3566890](#), [Zbl 1352.39021](#), doi: [10.1215/2008752-3749995](#). 222
- [29] KAAD, JENS; SKEIDE, MICHAEL. Kernels of Hilbert module maps: a counterexample. *J. Oper. Theory*, **89** (2023), no. 2, 343–348. [MR4591644](#), [Zbl 1563.46088](#), [arXiv:2101.03030](#), doi: [10.7900/jot.2021jun11.2338](#). 233
- [30] KASPAROV, G. G. Hilbert C^* -modules: theorems of Stinespring and Voiculescu. *J. Operator Theory* **4** (1980), no. 1, 133–150. [MR0587371](#), [Zbl 0456.46059](#). 229
- [31] KODAKA, KAZUNORI. Picard groups of irrational rotation C^* -algebras. *J. London Math. Soc.* (2) **56** (1997), no. 1, 179–188. [MR1462834](#), [Zbl 0892.46067](#), doi: [10.1112/S0024610797005243](#). 225
- [32] KOLAREC, BISERKA. A survey on extensions of Hilbert C^* -modules. *Quantum Probability and Related Topics*, 209–221. QP-PQ: Quantum Probability and White Noise Analysis, 29. *World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ*, 2013 ISBN:978-981-4447-53-9. [MR3288048](#), [Zbl 1327.81227](#), doi: [10.1142/9789814447546_0013](#). 222
- [33] LANCE, E. CHRISTOPHER. Unitary operators on Hilbert C^* -modules. *Bull. London Math. Soc.* **26** (1994), no. 4, 363–366. [MR1302069](#), [Zbl 0821.46073](#), doi: [10.1112/blms/26.4.363](#). 226
- [34] LANCE, E. CHRISTOPHER. Hilbert C^* -modules. A toolkit for operator algebraists. London Mathematical Society Lecture Note Series, 210. *Cambridge University Press, Cambridge, England*, 1995. x+130 pp. ISBN:0-521-47910-X. [MR1325694](#), [Zbl 0822.46080](#), doi: [10.1017/CBO9780511526206](#). 224
- [35] LIN, HUA XIN. The structure of quasimultipliers of C^* -algebras. *Trans. Amer. Math. Soc.* **315** (1989), no. 1, 147–172. [MR0937248](#), [Zbl 0684.46047](#), doi: [10.1090/s0002-9947-1989-0937248-3](#). 223, 230
- [36] LIN, HUA XIN. Bounded module maps and pure completely positive maps. *J. Operator Theory* **26** (1991), no. 1, 121–138. [MR1214924](#), [Zbl 0791.46032](#). 223, 232, 234
- [37] MANUILOV, VLADIMIR M.. On large submodules in Hilbert C^* -modules. *J. Math. Anal. Appl.* **542** (2025), no. 2, Paper No. 128781, 13pp. [MR4791358](#), [Zbl 1554.46038](#), [arXiv:2402.07288](#), doi: [10.1016/j.jmaa.2024.128781](#). 235
- [38] MURPHY, GERARD J. C^* -Algebras and operator theory. *Academic Press Inc., Boston, MA*, 1990. x+286 pp. ISBN:0-12-511360-9. [MR1074574](#), [Zbl 0714.46041](#), doi: [10.1016/C2009-0-22289-6](#). 223
- [39] NAROEI IRANI, MONA; NAZARI, AKBAR. The woven frame of multipliers in Hilbert C^* -modules. *Commun. Korean Math. Soc.* **36** (2021), no. 2, 257–266. [MR4252839](#), [Zbl 1479.42083](#), doi: [10.4134/CKMS.c200179](#). 225
- [40] PASCHKE, WILLIAM L. Inner product modules over B^* -algebras. *Trans. Amer. Math. Soc.* **182** (1973), 443–468. [MR0355613](#), [Zbl 0239.46062](#), doi: [10.2307/1996542](#). 224, 227, 234

- [41] PASCHKE, WILLIAM L. The double B -dual of an inner product module over a C^* -algebra B . *Can. J. Math.* **26** (1974), no. 5, 1272–1280. [MR0470687](#), [Zbl 0288.46058](#), doi: [10.4153/CJM-1974-121-0](#). 234
- [42] PEDERSEN, GERT K. Multipliers of AW^* -algebras. *Math. Z.* **187** (1984), no. 1, 23–24. [MR0753416](#), [Zbl 0547.46037](#), doi: [10.1007/BF01163162](#). 223
- [43] PEDERSEN, GERT K. Factorization in C^* -algebras. *Exposition. Math.* **16** (1998), no. 2, 145–156. [MR1630695](#), [Zbl 0912.46054](#). 226
- [44] RAEBURN, IAIN; WILLIAMS, DANA P. Morita equivalence and continuous trace C^* -algebras. Mathematical Surveys and Monographs 60. *American Mathematical Society, Providence, RI*, 1998. xiv+327 pp. ISBN:0-8218-0860-5. [MR1634408](#), [Zbl 0922.46050](#), doi: [10.1090/surv/060.224, 225, 226](#)
- [45] RAEBURN, IAIN; THOMPSON, SHAUN J.. Countably generated Hilbert modules, the Kasparov stabilization theorem, and frames with Hilbert modules. *Proc. Amer. Math. Soc.* **131** (2003), no. 5, 1557–1564. [MR1949886](#), [Zbl 1015.46034](#), doi: [10.1090/S0002-9939-02-06787-4](#). 228
- [46] SCHWEIZER, JÜRGEN. Hilbert C^* -modules with a predual. *J. Operator Theory* **48** (2002), no. 3, suppl., 621–632. [MR1962475](#), [Zbl 1029.46088](#). 221
- [47] SOLEL, BARUCH. Isometries of Hilbert C^* -modules. *Trans. Amer. Math. Soc.* **353** (2001), no. 11, 4637–4660. [MR1851186](#), [Zbl 0985.46035](#), [arXiv:math/0104188](#), doi: [10.1090/S0002-9947-01-02874-4](#). 221, 226, 235
- [48] TAYLOR, JONATHAN P. Aperiodic dynamical inclusions of C^* -algebras. Ph.D. Thesis, Georg-August-Universität Göttingen, Göttingen, Germany, (2022). doi: [10.53846/goediss-9727.222](#)
- [49] TAYLOR, JONATHAN P. Aperiodic dynamical inclusions of C^* -algebras. Preprint, 2023. [arXiv:2303.10905](#). 222
- [50] WEGGE-OLSEN, NIELS ERIK. K -theory and C^* -algebras: a friendly approach. Oxford Science Publications. *The Clarendon Press, Oxford University Press, New York*, 1993. xii+370 pp. ISBN:0-19-859694-4. [MR1222415](#), [Zbl 0780.46038](#), doi: [10.1093/oso/9780198596943.001.0001](#). 224, 226

(Michael Frank) HOCHSCHULE FÜR TECHNIK, WIRTSCHAFT UND KULTUR (HTWK) LEIPZIG, FAKULTÄT INFORMATIK UND MEDIEN, PF 301166, D-04251 LEIPZIG, GERMANY, ORCID 0000-0001-8972-2154

michael.frank@htwk-leipzig.de, michael.frank.leipzig@gmx.de

This paper is available via <http://nyjm.albany.edu/j/2026/32-8.html>.