New York Journal of Mathematics

New York J. Math. **12** (2006) 219–233.

Endomorphism rings of almost full formal groups

David J. Schmitz

Abstract. Let $\mathfrak{o}_K$ be the integral closure of $\mathbb{Z}_p$ in a finite field extension K of $\mathbb{Q}_p$, and let F be a one-dimensional full formal group defined over $\mathfrak{o}_K$. We study certain finite subgroups C of F and prove a conjecture of Jonathan Lubin concerning the absolute endomorphism ring of the quotient F/C when F has height 2. We also investigate ways in which this result can be generalized to p-adic formal groups of higher height.

Contents

Introduction 219
1. p-adic formal groups and isogenies 220
2. Points of finite order of a full formal group 224
3. Deflated subgroups 227
4. Generalizations of Conjecture 1 228
5. Free Tate modules of rank 1 230
6. Special results for height 2 formal groups 232
References 233

Introduction

In September, 2000, Jonathan Lubin conveyed to me the following two conjectures of his describing the quotients of full and almost full height 2 p-adic formal groups by certain finite subgroups:

Conjecture 1. *Let F be a full p-adic formal group of height 2, and let C be a cyclic subgroup of F having order p^n. Assume that* $\mathrm{End}(F)$*, the absolute endomorphism ring of F, is isomorphic to the ring of integers $\mathfrak{o}_K$ in a quadratic p-adic number field K; assume further that if $K/\mathbb{Q}_p$ is totally ramified, then C does not contain* $\ker[\pi]_F$*, where π is a uniformizer of $\mathfrak{o}_K$. Then* $\mathrm{End}(F/C) \cong \mathbb{Z}_p + p^n\,\mathfrak{o}_K$.

Conjecture 2. *Suppose G is an almost full p-adic formal group of height 2 with* $\mathrm{End}(G) \cong \mathbb{Z}_p + p^n\mathfrak{o}$*, where $\mathfrak{o}$ is some p-adic integer ring. Then there is a cyclic subgroup D of G of order p^n, canonical somehow, such that G/D is full.*

Received February 28, 2006.
Mathematics Subject Classification. 11S31, 14L05.
Key words and phrases. p-adic formal groups, endomorphism rings.

ISSN 1076-9803/06

We prove the first of these conjectures in this paper as Theorem 6.3. Furthermore, as we describe below, we are able to generalize this result in a couple of ways to p-adic formal groups of arbitrary (finite) height. The proofs of Conjecture 2 and some its generalizations are left for a subsequent paper. (See [S].)

If F is a p-adic formal group with $\mathrm{End}(F)$ integrally closed, then $c : g \mapsto g'(0)$ defines an isomorphism from $\mathrm{End}(F)$ onto a p-adic integer ring $\mathfrak{o}$. Via this association, we can view the torsion subgroup $\Lambda(F)$ of F as an $\mathfrak{o}$-module. For a finite subgroup C of $\Lambda(F)$, we denote by $\mathcal{I}(C)$ the annihilator of C in $\mathfrak{o}$. We prove the following as Theorem 4.3:

Theorem 1. *Let F be a p-adic formal group such that $\mathrm{End}(F)$ is integrally closed. If C is a finite cyclic subgroup of $\Lambda(F)$, then $c\big(\mathrm{End}(F/C)\big) = \mathbb{Z}_p + \mathcal{I}(C)$.*

This generalizes Conjecture 1 since $\mathcal{I}(C) = p^n\mathfrak{o}$ for the finite subgroups described there.

We are also able to say something about $c\big(\mathrm{End}(F/C)\big)$ when C is not necessarily cyclic. We will say that a finite subgroup C of the torsion subgroup of a p-adic formal group F is *a deflated subgroup of F* if there is no finite subgroup D of $\Lambda(F)$ having fewer elements than C such that $F/D \cong F/C$. We show in Section 3 that if F is full, then C is a deflated subgroup of F if and only if C does not contain the kernel of any noninvertible F-endomorphism. Lubin proves in [Lu2] that if F is full, then for any finite subgroup C of $\Lambda(F)$, $c(\mathrm{End}(F/C))$ is a subring of $c(\mathrm{End}(F))$. More specifically, we prove as Theorem 4.4:

Theorem 2. *Let F be a full p-adic formal group, and let C be a deflated subgroup of F. The conductor of $c\big(\mathrm{End}(F)\big)$ with respect to $c\big(\mathrm{End}(F/C)\big)$ is $\mathcal{I}(C)$.*

In Section 1, we review the basic theory of p-adic formal groups, paying particular attention to the integer rings over which certain homomorphisms are defined; we point out when some of the theorems from [Lu2] can be extended in this respect. In Section 2, we use the Tate module of F to study the $\mathrm{End}(F)$-module structure of the torsion subgroup of F. After describing the basic properties of deflated subgroups in Section 3, we prove in the final sections several theorems concerning almost full p-adic formal groups, including Theorem 1, Theorem 2, and Conjecture 1. We also see what other conclusions can be drawn in the height 2 case using our general theorems.

1. *p*-adic formal groups and isogenies

Fix a prime p. Let $\mathbb{C}_p$ be the completion of a fixed algebraic closure $\overline{\mathbb{Q}}_p$ of $\mathbb{Q}_p$ with respect to the unique extension of the p-adic valuation v on $\mathbb{Q}_p$ normalized so that $v(p) = 1$. Then v extends uniquely to a rational valuation on $\mathbb{C}_p$, and we denote this valuation by v as well. Let $\overline{\mathbb{Z}}_p$ (resp., $\mathfrak{O}$) be the set of elements in $\overline{\mathbb{Q}}_p$ (resp., in $\mathbb{C}_p$) with nonnegative valuation, and let $\overline{\mathfrak{m}}$ (resp., $\mathfrak{M}$) be the maximal ideal of $\overline{\mathbb{Z}}_p$ (resp., of $\mathfrak{O}$). For any subfield K of $\mathbb{C}_p$, we denote by $\mathfrak{o}_K$ the integer ring of K, i.e., $\mathfrak{o}_K = K \cap \mathfrak{O}$. Subfields of $\mathbb{C}_p$ which are finite extensions of $\mathbb{Q}_p$ are called *p-adic number fields*, and their integer rings are called *p-adic integer rings*. We define a *p-adic formal group* to be a one-dimensional formal group of finite height defined over a p-adic integer ring.

We will first review some of the basic results from the theory of p-adic formal groups. Proofs and more detailed discussions of these facts can be found in [F],

[Lu2], [Lu3], and [Laz]. Our purpose here is not merely to be expository. In many of the published works on p-adic formal groups, the theorems refer only to homomorphisms defined over a p-adic integer ring. Our methods will sometimes involve homomorphisms which are defined over the completion of a discretely-valued, infinite extension field of $\mathbb{Q}_p$. In this section, we will point out where the standard results can be extended to cover these "nonalgebraic" cases.

If F and G are two p-adic formal groups, then we define $\mathrm{Hom}(F,G)$ to be the abelian group of all homomorphisms from F to G defined over $\mathfrak{O}$. If there is some $g \in \mathrm{Hom}(F,G)$ with invertible linear coefficient, then F *is isomorphic to* G, written $F \cong G$, and g is called an *isomorphism from* F *to* G. It is easily shown that the compositional inverse g^{-1} of an isomorphism $g : F \to G$ belongs to $\mathrm{Hom}(G,F)$. If $F = G$, then we write $\mathrm{End}(F)$ instead of $\mathrm{Hom}(F,F)$, and we refer to it as the *absolute endomorphism ring of* F. The *automorphism group of* F, denoted by $\mathrm{Aut}(F)$, is the group of units of $\mathrm{End}(F)$.

For p-adic formal groups F and G, the map $c : \mathrm{Hom}(F,G) \to \mathfrak{O}$ sending a homomorphism $g : F \to G$ to its linear coefficient is an injective group homomorphism with closed image [Lu3, §2]. When $F = G$, c is a map of commutative $\mathbb{Z}_p$-algebras, for if $[n]_F$ is the multiplication-by-n endomorphism of F, then $c([n]_F) = n$. Following Lubin, we denote by $[a]_F$ *the* element of $\mathrm{End}(F)$ such that $c([a]_F) = a$, provided such an endomorphism exists. Another consequence of the injectivity of c is that if H is another p-adic formal group and if $0 \neq g \in \mathrm{Hom}(F,G)$ and $0 \neq j \in \mathrm{Hom}(G,H)$, then $0 \neq j \circ g \in \mathrm{Hom}(F,H)$. Furthermore, if $g \in \mathrm{Hom}(F,G)$ is an isomorphism, then $j \mapsto g \circ j \circ g^{-1}$ defines a ring isomorphism from $\mathrm{End}(F)$ onto $\mathrm{End}(G)$, and so $c\big(\mathrm{End}(F)\big) = c\big(\mathrm{End}(G)\big)$.

Lubin [Lu3, p 470] showed that if F is a p-adic formal group of height h, and if K is a p-adic number field containing the coefficients of F and all p-adic number fields of degree h over $\mathbb{Q}_p$, then $\mathrm{End}(F) \subset \mathfrak{o}_K[[T]]$. This is equivalent to stating $c\big(\mathrm{End}(F)\big) \subseteq \mathfrak{o}_K$ because each coefficient of $g \in \mathrm{Hom}(F,G)$ is a polynomial function of $c(g)$ with coefficients in any field containing the coefficients of F and G [F, p 98]. We denote by Σ_F the fraction field of $c\big(\mathrm{End}(F)\big)$. Since $\mathbb{Z}_p \subseteq c\big(\mathrm{End}(F)\big) \subseteq \mathfrak{o}_{\Sigma_F}$, we see that $c\big(\mathrm{End}(F)\big)$ is a $\mathbb{Z}_p$-order in Σ_F; moreover, $[\Sigma_F : \mathbb{Q}_p]$ is a divisor of h [Lu3, 2.3.2].

Definition 1.1. A p-adic formal group F of height h is *full* if $[\Sigma_F : \mathbb{Q}_p] = h$ and $c\big(\mathrm{End}(F)\big) = \mathfrak{o}_{\Sigma_F}$. We say F is *almost full* if $[\Sigma_F : \mathbb{Q}_p] = h$ but $c\big(\mathrm{End}(F)\big) \neq \mathfrak{o}_{\Sigma_F}$.

For any p-adic number field K, Lubin and Tate [LT] give a way of constructing full p-adic formal groups F defined over $\mathfrak{o}_K$ such that $c\big(\mathrm{End}(F)\big) = \mathfrak{o}_K$.

Whereas the endomorphisms of a p-adic formal group are all defined over a single p-adic integer ring, the same cannot be said of the homomorphisms between different p-adic formal groups. (See [Lu3, 4.3.2].) We will say that $g : F \to G$ is an *isogeny* if g is defined over some $\mathfrak{o}_L$ (or, equivalently, if $c(g) \in \mathfrak{o}_L$), where L is a complete, discretely-valued subfield of $\mathbb{C}_p$ containing the coefficients of F and G. We write $\mathrm{Isog}(F,G)$ for the set of all isogenies from F to G, and we say that F *is isogenous to* G if $\mathrm{Isog}(F,G) \neq 0$. We show later that $\mathrm{Isog}(F,G)$ is a subgroup of $\mathrm{Hom}(F,G)$. It is clear that every endomorphism of a p-adic formal group is an isogeny. In [Lu2] and [F], for example, an isogeny is assumed to be defined over the integers in a finite extension of the field over which the p-adic formal groups are

defined. We will show that those homomorphisms which satisfy our more general definition of isogeny share many of the properties exhibited by "p-adic isogenies".

A p-adic formal group F can be used to define an abelian group law on $\mathfrak{M}$ by setting $\alpha +_F \beta = F(\alpha, \beta)$ for $\alpha, \beta \in \mathfrak{M}$. We denote this group by $F(\mathfrak{O})$, and refer to it as *the points of* F. From the definition of a p-adic formal group, we see that for $\alpha, \beta \in F(\mathfrak{O})$, $v\left(\alpha +_F \beta\right) \geq \min\{v(\alpha), v(\beta)\}$, with equality if $v(\alpha) \neq v(\beta)$. For any $g \in \mathrm{Hom}(F, G)$, the association $\alpha \mapsto g(\alpha)$ defines a group homomorphism from $F(\mathfrak{O})$ to $G(\mathfrak{O})$, which we also denote by g. In particular, if the integer m is prime to p, then $[m]_F$ maps $F(\mathfrak{O})$ isomorphically onto itself, and so the order of an element of $F(\mathfrak{O})$ of finite order is necessarily a power of p. Therefore the torsion subgroup $\Lambda(F)$ of the points of F can be expressed as

$$\Lambda(F) = \bigcup_{n \in \mathbb{N}} \ker [p^n]_F.$$

Proposition 1.2. *If $g \in \mathrm{Hom}(F, G)$ and $\alpha \in F(\mathfrak{O})$, then $v\big(g(\alpha)\big) \geq v(\alpha)$, with equality if and only if either $\alpha = 0$ or $c(g) \in \mathfrak{O}^\times$.*

Proof. Writing $g(T) = T \cdot j(T)$, where $j(T) \in \mathfrak{O}[[T]]$, we see that $v\big(g(\alpha)\big) \geq v(\alpha)$ because $j(\alpha) \in \mathfrak{O}$. Furthermore, if $\alpha \neq 0$, then $v\big(g(\alpha)\big) = v(\alpha)$ if and only if $v\big(j(\alpha)\big) = 0$, which holds if and only if $j(0) = c(g)$ is a unit in $\mathfrak{O}$ because $v(\alpha) > 0$. □

If g is a nonzero isogeny defined over the complete discretely-valued subring $\mathfrak{o}_L$ of $\mathfrak{O}$, then the Weierstrass Preparation Theorem [Lang, V.11.2] implies that there is a monic polynomial $P(T) \equiv T^d \pmod{\mathfrak{m}_L}$ of degree $d = \mathrm{wdeg}\,(g)$, the *Weierstrass degree of* g, and a power series $U(T) \in \mathfrak{o}_L[[T]]$ with $U(0) \notin \mathfrak{m}_L$ such that $g = P \cdot U$. The elements of $\ker(g)$ are the roots of $P(T)$; they belong to $\mathfrak{M}$ and have multiplicity one [Lu2, §1.2]. Thus, the kernel of any nonzero isogeny $g : F \to G$ is a finite subgroup of $F(\mathfrak{O})$ of order $\mathrm{wdeg}\,(g)$. In particular, $\ker [p]_F$ has order p^h, where h is the height of F. The elements of $\Lambda(F)$ are all integral over $\mathbb{Z}_p$: indeed, for every $n \in \mathbb{N}$, $[p^n]_F$, is defined over any p-adic integer ring $\mathfrak{o}_K$ containing the coefficients of F, and so the polynomial $P(T) \in \mathfrak{o}_K[T]$ arising from the Weierstrass Preparation Theorem has roots in $\overline{\mathfrak{m}}$.

If $g \in \mathrm{Hom}(F, G)$, then for every $m \in \mathbb{Z}$, $[m]_G \circ g = g \circ [m]_F$, and therefore $g\left(\Lambda(F)\right) \subseteq \Lambda(G)$. A slight modification of the argument in [Lu2, §1.2] will show that $g : \Lambda(F) \to \Lambda(G)$ is surjective whenever g is a nonzero isogeny. Suppose that g is defined over $\mathfrak{o}_L$, where L is a complete, discretely-valued subfield of $\mathbb{C}_p$. For any $\alpha \in \Lambda(G)$, the power series $g(T) - \alpha$ is defined over the ring of integers in $L(\alpha)$ (which is also a complete discretely-valued subfield of $\mathbb{C}_p$ because α is integral over $\mathbb{Z}_p$), and $\mathrm{wdeg}\big(g(T) - \alpha\big) = \mathrm{wdeg}\,(g) \geq 1$. The Weierstrass Preparation Theorem implies that $g(T) - \alpha$ has $\mathrm{wdeg}\,(g)$ zeros in $F(\mathfrak{O})$ all belonging to $\Lambda(F)$ since $\alpha \in \Lambda(G)$ and g is a homomorphism of p-adic formal groups having a finite kernel.

If C is a finite subgroup of $F(\mathfrak{O})$, Lubin [Lu2, 1.4] proved that the power series

$$\varphi_C(T) = \prod_{\gamma \in C} F(T, \gamma)$$

is a p-adic isogeny from F to the p-adic formal group $\varphi_C\left(F\big({\varphi_C}^{-1}(X), {\varphi_C}^{-1}(Y)\big)\right)$, which we denote by F/C and refer to as *the quotient of F by C*. It is clear that $\ker(\varphi_C) = C$. Lubin showed that any p-adic isogeny $j : F \to H$ vanishing on C

factors uniquely through F/C. Using nearly the same proof, one can show that this fact holds for any such isogeny j. One needs only to observe (as above) that if K is a complete discretely-valued subfield of $\mathbb{C}_p$ and if $C = \{\alpha_1, \ldots, \alpha_n\}$ is a finite subgroup of $\Lambda(F)$, then $K(\alpha_1, \ldots, \alpha_n)$ is also a complete discretely-valued subfield of $\mathbb{C}_p$. We record the precise result here.

Theorem 1.3 ([Lu2, 1.5])**.** *Let F, G, H be p-adic formal groups and let L be a complete discretely-valued subfield of $\mathbb{C}_p$ containing the coefficients of F, G, and H. If $g_1 : F \to G$, $g_1 \neq 0$, and $g_2 : F \to H$ are isogenies defined over $\mathfrak{o}_L$ such that $\ker(g_1) \subseteq \ker(g_2)$, then there is a unique isogeny $j : G \to H$ defined over $\mathfrak{o}_L$ such that $j \circ g_1 = g_2$. If $\ker(g_1) = \ker(g_2)$, then j is an isomorphism.*

We can interpret Theorem 1.3 in terms of divisibility in the ring $c\big(\mathrm{End}(F)\big)$.

Corollary 1.4. *Let F be a p-adic formal group, and let $\zeta_1, \zeta_2 \in c\big(\mathrm{End}(F)\big)$. Then ζ_1 divides ζ_2 in $c\big(\mathrm{End}(F)\big)$ if and only if $\ker [\zeta_1]_F \subseteq \ker [\zeta_2]_F$. In particular, ζ_1 and ζ_2 are associates in $c\big(\mathrm{End}(F)\big)$ if and only if $\ker [\zeta_1]_F = \ker [\zeta_2]_F$.*

Proof. If there is an $\eta \in c\big(\mathrm{End}(F)\big)$ such that $\eta \cdot \zeta_1 = \zeta_2$, then $[\eta]_F \circ [\zeta_1]_F = [\zeta_2]_F$, and so $\ker [\zeta_1]_F$ is contained in $\ker [\zeta_2]_F$. Conversely, if $\ker [\zeta_1]_F \subseteq \ker [\zeta_2]_F$, then we may apply Theorem 1.3 to find $j \in \mathrm{End}(F)$ such that $j \circ [\zeta_1]_F = [\zeta_2]_F$. Therefore, $c(j) \cdot \zeta_1 = \zeta_2$. □

The next result shows that, like endomorphisms of a p-adic formal group, all homomorphisms between isogenous p-adic formal groups are defined over a single complete discretely-valued subring of $\mathbb{C}_p$.

Proposition 1.5. *Let F and G be p-adic formal groups, and assume $g : F \to G$ is a nonzero isogeny defined over the integers $\mathfrak{o}_L$ in a complete discretely-valued subfield L of $\mathbb{C}_p$ containing Σ_F and the coefficients of F and G. Then $\mathrm{Isog}(F, G) = \mathrm{Hom}(F, G) \subset \mathfrak{o}_L[[T]]$.*

Proof. By [Lu2, §1.6], there exists a nonzero isogeny $\widetilde{g} : G \to F$ defined over $\mathfrak{o}_L$. Post-composition with $\widetilde{g}$ defines an injective group homomorphism from $\mathrm{Hom}(F, G)$ to $\mathrm{End}(F)$. So, for any $j \in \mathrm{Hom}(F, G)$, $c(\widetilde{g}) \cdot c(j) \in c\big(\mathrm{End}(F)\big) \subset L$, whence $c(j) \in \mathfrak{O} \cap L = \mathfrak{o}_L$. □

Corollary 1.6. *For p-adic formal groups F and G, either $\mathrm{Isog}(F, G) = 0$ or $\mathrm{Isog}(F, G) = \mathrm{Hom}(F, G)$. In either case, $\mathrm{Isog}(F, G)$ is a group.*

The next corollary is essentially a generalization of a result in [Lu2, §3.2] which states that an almost full p-adic formal group is isogenous to a full p-adic formal group.

Corollary 1.7. *Let $\{G_i\}$ $(i = 1, ..., n)$ be full or almost full p-adic formal groups such that $\Sigma_{G_1} = \cdots = \Sigma_{G_n} = \Sigma$. Then there is a complete discretely-valued subfield L of $\mathbb{C}_p$ such that $0 \neq \mathrm{Isog}(G_i, G_j) = \mathrm{Hom}(G_i, G_j) \subset \mathfrak{o}_L[[T]]$ for every $1 \leq i, j \leq n$.*

Proof. According to [Lu2, §3.2], for each $i = 1, \ldots n$, there is a full p-adic formal group F_i and nonzero p-adic isogenies $g_i : F_i \to G_i$ and $\widetilde{g}_i : G_i \to F_i$. Let K be a p-adic number field containing Σ and the coefficients of all of these p-adic formal groups and isogenies. For each $1 \leq i, j \leq n$, $\Sigma_{F_i} = \Sigma_{G_i} = \Sigma_{G_j} = \Sigma_{F_j}$ [Lu2, §3.0], and so there is an isomorphism $u_{ij} : F_i \to F_j$ defined over $\mathfrak{o}_L$, where L is the

completion of the maximal unramified extension K^{nr} of K [Lu3, 4.3.2]. Because K^{nr} is discretely-valued, so is L. Therefore,

$$0 \neq g_j \circ u_{ij} \circ \widetilde{g_i} \in \mathrm{Hom}(G_i, G_j) \cap \mathfrak{o}_L[[T]] \subseteq \mathrm{Isog}(G_i, G_j).$$

The corollary now follows from Proposition 1.5. □

We conclude with our main tool for investigating almost full p-adic formal groups.

Corollary 1.8. *Let G be an almost full p-adic formal group. Then there is a full p-adic formal group F and a finite subgroup C of $\Lambda(F)$ such that G is isomorphic to F/C over a p-adic integer ring.*

Proof. As in the proof of Corollary 1.7, we can find a full p-adic formal group F with $\Sigma_F = \Sigma_G$ and a nonzero isogeny $g : F \to G$ defined over a p-adic integer ring. If $C = \ker(g)$, then $\ker(g) = \ker(\varphi_C)$, and so G and F/C are isomorphic over a p-adic integer ring by Theorem 1.3. □

The main focus of the rest of this article will be to see how the structure of the subgroup C influences that of the ring $\mathrm{End}(F/C)$.

2. Points of finite order of a full formal group

In this section, we investigate certain structures within and on the torsion subgroup a full p-adic formal group F. We are primarily interested in the F-endomorphism kernels and the cyclic subgroups contained in $\Lambda(F)$, two kinds of subgroups mentioned in Conjecture 1. Furthermore, a study of the $c\big(\mathrm{End}(F)\big)$-module structure on $\Lambda(F)$ will provide the key to our proof of Conjecture 1. We first review some facts concerning the Tate module of F.

For any p-adic formal group F of height h, the *Tate module* of F is defined to be

$$T(F) = \varprojlim \ker [p^n]_F$$

where the inverse limit is taken with respect to the surjective homomorphisms $[p]_F : \ker [p^{n+1}]_F \to \ker [p^n]_F$. If G is another p-adic formal group, then any homomorphism $g : F \to G$ defines a group homomorphism $T(g) : T(F) \to T(G)$ by $T(g)\big((a_0, a_1, \dots)\big) = \big(g(a_0), g(a_1), \dots\big)$. If $0 \neq g \in \mathrm{Isog}(F, G)$, then $\ker(g)$ is finite, and hence $T(g)$ is injective. In particular, $T(F)$ is a torsion-free $c\big(\mathrm{End}(F)\big)$-module and a free $\mathbb{Z}_p$-module of rank h [F, IV §4]. If $c\big(\mathrm{End}(F)\big)$ is integrally closed (and thus a PID) of rank d over $\mathbb{Z}_p$, then $T(F)$ is a free $c\big(\mathrm{End}(F)\big)$-module of rank $\frac{h}{d}$. Therefore, when F is full, $T(F)$ is free of rank 1 over $c\big(\mathrm{End}(F)\big)$. In Proposition 5.1, we derive a condition for determining when the Tate module of an almost full p-adic formal group G is free of rank 1 over $c\big(\mathrm{End}(G)\big)$.

We denote by $V(F)$ the set of sequences $(a_0, a_1, \dots)$ such that for all $n \geq 0$, $a_n \in \Lambda(F)$ and $[p]_F(a_{n+1}) = a_n$. It is not difficult to see that $V(F) \cong T(F) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$, whence $V(F)$ is an h-dimensional $\mathbb{Q}_p$-vector space, called the *Tate vector space of* F. If $g \in \mathrm{Hom}(F, G)$, the $\mathbb{Z}_p$-module homomorphism $T(g) : T(F) \to T(G)$ extends to a linear map $V(g) : V(F) \to V(G)$ of $\mathbb{Q}_p$-vector spaces which is injective if g is a nonzero isogeny. In fact, the existence of such a g implies that F and G have equal heights [Lu3, 2.2.3 and 2.3.1], and therefore $V(g)$ is an isomorphism. Since $\Sigma_F = c\big(\mathrm{End}(F)\big) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$, the $c\big(\mathrm{End}(F)\big)$-module structure on $T(F)$ induces a Σ_F-vector space structure on $V(F)$. If $[\Sigma_F : \mathbb{Q}_p] = d$, then $V(F)$ is an $\frac{h}{d}$-dimensional

Σ_F-vector space; in particular, when F is full or almost full, $V(F)$ is 1-dimensional over Σ_F. Finally, if $0 \neq g \in \mathrm{Isog}(F,G)$, then $\Sigma_F = \Sigma_G$ and $V(g) : V(F) \to V(G)$ is a Σ_F-isomorphism.

Proposition 2.1. *If $g, j \in \mathrm{Isog}(F,G)$, then $V(g) = V(j)$ if and only if $g = j$.*

Proof. Indeed, if $V(g) = V(j)$, then $g(\alpha) = j(\alpha)$ for all $\alpha \in \Lambda(F)$, which implies that $g - j$ is identically 0 on $\Lambda(F)$. Since $\mathrm{Isog}(F,G)$ is a group, $g - j \in \mathrm{Isog}(F,G)$, and so its kernel is finite unless $g - j = 0$. □

Throughout the remainder of this section, we denote by F a p-adic formal group of height h with $\mathrm{End}(F)$ integrally closed, and we let π be a fixed uniformizer of $c\big(\mathrm{End}(F)\big)$. Moreover, we denote by e (resp., f) the ramification index (resp., the residue field degree) of the extension $\Sigma_F/\mathbb{Q}_p$.

The group $\Lambda(F)$ is the union of the kernels of the endomorphisms $[p^n]_F$ $(n \geq 0)$. If g is any nonzero endomorphism of F, then $\ker(g)$ is also a finite subgroup of $\Lambda(F)$, not necessarily equal to the kernel of one of the multiplication-by-p^n endomorphisms. However, $c(g)$ is an associate of π^m in the ring $c\big(\mathrm{End}(F)\big)$, where $m = e \cdot v\big(c(g)\big)$, and so by Corollary 1.4, $\ker(g) = \ker[\pi^m]_F$. Therefore, $\big\{\ker[\pi^m]_F\big\}_{m\geq 0}$ is the set of kernels of the nonzero F-endomorphisms, and

$$\Lambda(F) = \bigcup_{n\geq 0} \ker[p^n]_F = \bigcup_{m\geq 0} \ker[\pi^m]_F.$$

Moreover, because $\ker[\pi^{m-1}]_F \subset \ker[\pi^m]_F$, the family $\big\{\ker[\pi^m]_F\big\}_{m\geq 0}$ is a filtration of subgroups of $\Lambda(F)$, with $\ker[\pi]_F$ being the smallest kernel of any noninvertible F-endomorphism.

Proposition 2.2. *The kernel of $[\pi^m]_F$ has $p^{m(h/e)}$ elements. In particular, if F is full, then $\big|\ker[\pi^m]_F\big| = p^{mf}$.*

Proof. If $\big|\ker[\pi]_F\big| = p^s$, then the surjectivity of $[\pi]_F : \Lambda(F) \to \Lambda(F)$ implies inductively that $\big|\ker[\pi^m]_F\big| = p^{sm}$. Therefore $p^h = \big|\ker[p]_F\big| = \big|\ker[\pi^e]_F\big| = p^{se}$, and so $s = h/e$. Finally, when F is full, we note that $h = [\Sigma_F : \mathbb{Q}_p] = ef$. □

We can interpret the endomorphism kernels in terms of annihilators.

Definition 2.3. The *annihilator* $\mathcal{I}(X)$ of a subset X of $\Lambda(F)$ is the set

$$\big\{\zeta \in c\big(\mathrm{End}(F)\big) \;\big|\; \forall \alpha \in X, [\zeta]_F(\alpha) = 0\big\}.$$

If $\gamma \in \Lambda(F)$, we will write $\mathcal{I}(\gamma)$ instead of $\mathcal{I}(\{\gamma\})$.

Remarks 2.4.

(i) Because $\mathfrak{o} = c\big(\mathrm{End}(F)\big)$ is a commutative ring, $\mathcal{I}(X)$ is an ideal of $\mathfrak{o}$. Therefore $\mathcal{I}(X) = \pi^m\mathfrak{o}$ for some integer $m \geq 0$. In fact, for each $m \in \mathbb{N}$,

$$\big\{\alpha \in \Lambda(F) \;\big|\; \mathcal{I}(\alpha) = \pi^m\mathfrak{o}\big\} = \ker[\pi^m]_F - \ker[\pi^{m-1}]_F\,.$$

(ii) If C is the cyclic subgroup generated by $\gamma \in \Lambda(F)$, then $\mathcal{I}(C) = \mathcal{I}(\gamma)$. More generally, it follows from Lemma 2.5 below that if C is any finite subgroup of $\Lambda(F)$, where F is a full p-adic formal group, then $\mathcal{I}(C) = \mathcal{I}(\gamma)$, where $\gamma \in C$ is an element of minimal valuation.

We have seen (Corollary 1.8) that any almost full p-adic formal group is isomorphic over a p-adic integer ring to the quotient of a full p-adic formal group F by a finite subgroup C of $\Lambda(F)$. The quotient is much easier to study when the subgroup C can be chosen to be cyclic; this is always possible in height 2 (see §6). In Corollary 6.4, we will use this fact to prove that the isomorphism class of a height 2 almost full p-adic formal group depends only on its absolute endomorphism ring. A key step in our proof is the result given below in Corollary 2.8, which describes when two cyclic subgroups of $\Lambda(F)$ are isomorphic to each other via an automorphism of F. We begin, however, with the following lemma, the proof of which uses the fact that $T(F)$ is free of rank 1 over $c\big(\mathrm{End}(F)\big)$.

Lemma 2.5. *Let F be a full p-adic formal group. For any pair $\gamma, \delta \in \Lambda(F)$, $v(\gamma) \leq v(\delta)$ if and only if there exists some $\zeta \in c\big(\mathrm{End}(F)\big)$ such that $[\zeta]_F(\gamma) = \delta$.*

Proof. Without loss of generality, we may assume that both γ and δ are nonzero. The implication ($\Leftarrow$) follows from Proposition 1.2. Conversely, suppose $v(\gamma) \leq v(\delta)$, and choose n large enough so that $\gamma, \delta \in \ker [p^n]_F$. Then there exist $c, d \in T(F)$ such that $c_n = \gamma$ and $d_n = \delta$. If $b = (b_0, b_1, \dots)$ is any basis of $T(F)$ over $c\big(\mathrm{End}(F)\big)$, then there are (unique) elements $\eta, \theta \in c\big(\mathrm{End}(F)\big)$ such that $\eta \cdot b = c$ and $\theta \cdot b = d$. Assume $v(\eta) \leq v(\theta)$. Then $\zeta = \theta\, \eta^{-1} \in \mathfrak{o}_{\Sigma_F} = c\big(\mathrm{End}(F)\big)$ and $\delta = [\theta]_F(b_n) = [\theta\, \eta^{-1}]_F\big([\eta]_F(b_n)\big) = [\zeta]_F(\gamma)$, which proves the lemma in this case. If, on the other hand, $v(\eta) > v(\theta)$, then a similar calculation would show that $[\eta\, \theta^{-1}]_F(\delta) = \gamma$, which contradicts Proposition 1.2 since $\eta\, \theta^{-1}$ is not a unit in $c\big(\mathrm{End}(F)\big)$. □

If C is any subgroup of $F(\mathfrak{O})$ and if $\lambda \in \mathbb{R}$, then $C_\lambda = \{\gamma \in C |\, v(\gamma) \geq \lambda\}$ is a subgroup of C. Using Lemma 2.5 and Proposition 1.2, we can obtain a description of the cyclic $\mathrm{End}(F)$-submodules of $\Lambda(F)$ when F is full. For any $\alpha \in \Lambda(F)$,

$$\mathrm{End}(F) \cdot \alpha = \big\{\beta \in \Lambda(F) \;\big|\; v(\beta) \geq v(\alpha)\big\} = \Lambda(F)_{v(\alpha)}\,.$$

The subsets $\Lambda(F)_{v(\alpha)}$ are examples of *congruence-torsion subgroups of F* (see [Lu1]). These turn out to be the so-called "canonical subgroups" mentioned in Conjecture 2.

Theorem 2.6. *Let F be a full p-adic formal group. The following are equivalent for elements $\gamma, \delta \in \Lambda(F)$:*

(i) $v(\gamma) = v(\delta)$.
(ii) *There exists some $u \in \mathrm{Aut}(F)$ such that $u(\gamma) = \delta$.*
(iii) $\mathcal{I}(\gamma) = \mathcal{I}(\delta)$.

Proof. (i) ⇒ (ii): This follows immediately from Lemma 2.5 and Proposition 1.2.

(ii) ⇒ (iii): If $\epsilon = c(u) \in c\big(\mathrm{End}(F)\big)^\times$, then $\zeta \mapsto \zeta \cdot \epsilon$ is a bijection from $\mathcal{I}(\delta)$ onto $\mathcal{I}(\gamma)$. Because these two sets are ideals of $c\big(\mathrm{End}(F)\big)$, they are equal.

(iii) ⇒ (i): Without loss of generality, we may assume that $v(\gamma) < v(\delta)$. Choose $\zeta \in c\big(\mathrm{End}(F)\big)$ such that $[\zeta]_F(\gamma) = \delta$ and suppose π^m ($m \geq 1$) generates $\mathcal{I}(\gamma)$. Since ζ is not a unit in $c\big(\mathrm{End}(F)\big)$ (Proposition 1.2), $\zeta = \pi\, \eta$ for some $\eta \in c\big(\mathrm{End}(F)\big)$. Then $\pi^{m-1} \in \mathcal{I}(\delta)$ because $[\pi^{m-1}]_F(\delta) = [\pi^m]_F\big([\eta]_F(\gamma)\big) = [\eta]_F\big([\pi^m]_F(\gamma)\big) = 0$. Therefore, $\mathcal{I}(\gamma) \neq \mathcal{I}(\delta)$. □

Corollary 2.7. *Let F be a full p-adic formal group. For any $m \in \mathbb{N}$, $\mathrm{Aut}(F)$ acts transitively on the set* $\ker [\pi^m]_F - \ker [\pi^{m-1}]_F$.

Proof. Using Remark 2.4(i) and Theorem 2.6 (iii) $\Rightarrow$ (i), we see that all the elements of $\ker[\pi^m]_F - \ker[\pi^{m-1}]_F$ have the same valuation, which, in light of Lemma 2.5, is less than the valuation of any of the elements of $\ker[\pi^{m-1}]_F$. The corollary now follows from Theorem 2.6 (i) $\Rightarrow$ (ii). □

Corollary 2.8. *Let F be a full p-adic formal group and let C_1 and C_2 be finite cyclic subgroups of $\Lambda(F)$. Then there exists some $u \in \mathrm{Aut}(F)$ such that $C_1 = u(C_2)$ if and only if $\mathcal{I}(C_1) = \mathcal{I}(C_2)$.*

Proof. This follows from Remark 2.4(ii) and Theorem 2.6. □

3. Deflated subgroups

When expressing a full or almost full p-adic formal group G as being isomorphic to the quotient of a full p-adic formal group F by a finite subgroup C of $\Lambda(F)$, F is uniquely determined up to isomorphism. Indeed, if $F/C \cong F'/C'$, where F and F' are full, then $\Sigma_F = \Sigma_{F/C} = \Sigma_{F'/C'} = \Sigma_{F'}$ (see the proof of Corollary 1.7), whence $F \cong F'$ via an isogeny [Lu3, 4.3.2]. However, the subgroup C is by no means unique (not even up to isomorphism).

Proposition 3.1. *Let F be any p-adic formal group. If C is a finite subgroup of $\Lambda(F)$ and $0 \neq g \in \mathrm{End}(F)$, then $F/g^{-1}(C) \cong F/C$ over a p-adic integer ring.*

Proof. Since $g^{-1}(C)$ is the kernel of the p-adic isogenies $\varphi_{g^{-1}(C)} : F \to F/g^{-1}(C)$ and $\varphi_C \circ g : F \to F/C$, we can use Theorem 1.3. □

Taking $g = [p^n]_F$ for various $n \in \mathbb{N}$, we see that there are infinitely many nonisomorphic finite subgroups of $\Lambda(F)$ which yield isomorphic quotients. This prompts the following.

Definition 3.2. Let F be a p-adic formal group. For finite subgroups C_1, C_2 of $\Lambda(F)$, we write $C_1 \sim C_2$ if $F/C_1 \cong F/C_2$.

It is clear that $\sim$ is an equivalence relation on the set of finite subgroups of $\Lambda(F)$. If C and D are two subgroups of $\Lambda(F)$ such that $C \sim D$, then we will say that C and D are *equivalent*. We now show that when F is a full p-adic formal group, then the converse of Proposition 3.1 is true.

Proposition 3.3. *Let F be a full p-adic formal group and let C, D be equivalent finite subgroups of $\Lambda(F)$. If $|C| \geq |D|$, then there exists $0 \neq g \in \mathrm{End}(F)$ such that $C = g^{-1}(D)$.*

Proof. By assumption, there is an isomorphism $u : F/C \to F/D$, and according to Proposition 1.5, the homomorphism $u \circ \varphi_C$ is a nonzero isogeny (since φ_D is). Thus, the maps $V(u\circ\varphi_C), V(\varphi_D) : V(F) \to V(F/D)$ are isomorphisms of Σ_F-vector spaces (see §2). Also, since F is full, F/D must be full or almost full [Lu2, 3.0], and so $V(F)$ and $V(F/D)$ are one-dimensional over Σ_F. Consequently, $V(u\circ\varphi_C)$ $\big($resp., $V(\varphi_D)\big)$ is scalar multiplication by some nonzero element α (resp., β) of Σ_F. Assume now that $\beta^{-1}\alpha \in c\big(\mathrm{End}(F)\big)$, and let $g = [\beta^{-1}\alpha]_F$. Then $V(g)$ operates on $V(F)$ via scalar multiplication by $\beta^{-1}\alpha$, and so $V(u \circ \varphi_C) = V(\varphi_D) \circ V(g) = V(\varphi_D \circ g)$. Therefore, $u \circ \varphi_C = \varphi_D \circ g$ by Proposition 2.1. Comparing kernels, we see that $C = g^{-1}(D)$.

We now show that $\beta^{-1}\alpha$ must be in $c\big(\mathrm{End}(F)\big)$. If $\beta^{-1}\alpha \notin c\big(\mathrm{End}(F)\big)$, then because $c\big(\mathrm{End}(F)\big)$ is a valuation ring, it follows that $\alpha^{-1}\beta \in c\big(\mathrm{End}(F)\big)$, but it is not a unit. The same reasoning as above shows that $\varphi_D = (u \circ \varphi_C) \circ \widetilde{g}$, where $\widetilde{g} = [\alpha^{-1}\beta]_F$. This implies that $\widetilde{g}^{-1}(C) = D$, and since $\ker\big(\widetilde{g}\big) \neq \{0\}$, we arrive at $|D| > |C|$, a contradiction. □

If F is a full p-adic formal group and C a finite subgroup of $\Lambda(F)$, then many properties of $\Lambda(F/C)$ and $\mathrm{End}(F/C)$ depend on the element(s) of minimal size in the equivalence class of C. We now name these subgroups.

Definition 3.4. Let F be a p-adic formal group. A finite subgroup D of $\Lambda(F)$ is a *deflated subgroup of* F if $D \sim C$ implies $|D| \leq |C|$.

There may be multiple deflated subgroups of F belonging to the same equivalence class. Indeed, if $u \in \mathrm{Aut}(F)$ and if D is a deflated subgroup of F, then $u^{-1}(D) \sim D$ and $u^{-1}(D)$ is deflated since $|u^{-1}(D)| = |D|$. On the other hand, if $\ker(g) \subseteq D$ for some $0 \neq g \in \mathrm{End}(F) - \mathrm{Aut}(F)$, then D is not deflated. To see this, we notice that $g(D) \sim D$ because $g^{-1}\big(g(D)\big) = D$, and $|g(D)| < |D|$ because $\ker(g) \neq \{0\}$. In the next theorem, we show that when F is full, this property characterizes the nondeflated subgroups of F.

Theorem 3.5. *Let F be a full p-adic formal group. A finite subgroup C of $\Lambda(F)$ is a deflated subgroup of F if and only if $\ker [\pi]_F \nsubseteq C$.*

Proof. We have already shown why C is not a deflated subgroup of F if it contains $\ker [\pi]_F$. Conversely, if C is not a deflated subgroup of F, then there is a finite subgroup D of $\Lambda(F)$ such that $D \sim C$ and $|D| < |C|$. By Proposition 3.3, there is some $0 \neq g \in \mathrm{End}(F)$ such that $C = g^{-1}(D)$; in particular, $\ker(g) \subseteq C$. Also, $\ker(g) \neq \{0\}$ because $|C| \neq |D|$. The result now follows since the kernels of the endomorphisms of F are totally ordered with respect to inclusion, with $\ker [\pi]_F$ the smallest nonzero subgroup among them. □

If F is a p-adic formal group of height 1, then $c\big(\mathrm{End}(F)\big) = \mathbb{Z}_p$, and F is necessarily full. We can take p to be a uniformizer of $c\big(\mathrm{End}(F)\big)$, and $\ker [p]_F$ has order p. It follows that *every* nonzero finite subgroup C of $\Lambda(F)$ is cyclic and contains $\ker [p]_F$; therefore, by Theorem 3.5, C is not a deflated subgroup of F. However, for full p-adic formal groups F of height $h > 1$, nondeflated cyclic subgroups are more the exception than the rule. According to Theorem 3.5, F has nondeflated cyclic subgroups if and only if $\ker [\pi]_F$ is cyclic, where π is a uniformizer of $c\big(\mathrm{End}(F)\big)$. Using Proposition 2.2, plus the fact that $\ker [\pi]_F \subseteq \ker [p]_F$, we see that $\ker [\pi]_F$ is cyclic if and only if $\Sigma_F / \mathbb{Q}_p$ is totally ramified.

We can now restate Conjecture 1 more concisely using the terminology and notation we have developed so far:

Conjecture 1. *Let F be a full p-adic formal group of height 2, and let C be a deflated cyclic subgroup of F of order p^n. Then $c\big(\mathrm{End}(F/C)\big) = \mathbb{Z}_p + p^n\mathfrak{o}$, where $\mathfrak{o} = c\big(\mathrm{End}(F)\big)$.*

4. Generalizations of Conjecture 1

We now prove a couple of theorems which generalize Conjecture 1 to p-adic formal groups of arbitrary height. First we look at the situation where the finite

subgroup C is cyclic, but not necessarily deflated, and then where C is deflated, but not necessarily cyclic. Our main tool is Lemma 4.1, which is a special case of [Lu2, 3.1].

Lemma 4.1. *Let F be a p-adic formal group such that $\mathrm{End}(F)$ is integrally closed. If C is a finite subgroup of $\Lambda(F)$, then*

$$c\big(\mathrm{End}(F/C)\big) = \big\{\zeta \in c\big(\mathrm{End}(F)\big) \;\big|\; [\zeta]_F(C) \subseteq C\big\}.$$

Proof. Let L be the lattice in $V(F)$ consisting of all elements $(a_0, a_1, \dots)$ with $a_0 \in C$. Then L is the lattice corresponding to $\varphi_C : F \to F/C$ as described in [Lu2, §2.2]. Therefore, according to [Lu2, 3.1],

$$c\big(\mathrm{End}(F/C)\big) = \big\{\zeta \in \Sigma_F \,\big|\, \zeta L \subseteq L\big\}.$$

Because $c\big(\mathrm{End}(F/C)\big)$ is a $\mathbb{Z}_p$-order in Σ_F, $c\big(\mathrm{End}(F/C)\big) \subseteq \mathfrak{o}_{\Sigma_F} = c\big(\mathrm{End}(F)\big)$. Thus

$$c\big(\mathrm{End}(F/C)\big) = \big\{\zeta \in c\big(\mathrm{End}(F)\big) \,\big|\, \zeta L \subseteq L\big\}.$$

But for $\zeta \in c\big(\mathrm{End}(F)\big)$ and $a = (a_0, a_1, \dots) \in V(F)$, $\zeta \cdot a = \big([\zeta]_F(a_0), [\zeta]_F(a_1), \dots\big)$. Hence $\zeta L \subseteq L$ if and only if $[\zeta]_F(C) \subseteq C$. □

Remark 4.2. If G is a p-adic formal group where $c\big(\mathrm{End}(G)\big)$ is not integrally closed, then there is some $n \in \mathbb{N}$ such that $p^n\, \mathfrak{o}_{\Sigma_G} \subseteq c\big(\mathrm{End}(G)\big)$. In this case, recall that for $\zeta \in \mathfrak{o}_{\Sigma_G}$ and $a = (a_0, a_1, \dots) \in V(G)$,

$$\zeta \cdot a = \big([p^n\zeta]_G(a_n), [p^n\zeta]_G(a_{n+1}), \dots\big).$$

A modification of the proof of Lemma 4.1 yields

$$c\big(\mathrm{End}(G/C)\big) = \Big\{\zeta \in \mathfrak{o}_{\Sigma_G} \,\Big|\, [p^n\zeta]_G\big([p^n]_G^{-1}(C)\big) \subseteq C\Big\}.$$

When F is a full p-adic formal group and C is a cyclic subgroup of $\Lambda(F)$, then the ring $c\big(\mathrm{End}(F/C)\big)$ has a rather simple description in terms of the annihilator of C in $\mathfrak{o} = c\big(\mathrm{End}(F)\big)$. We note that Theorem 4.3 is a generalization of Conjecture 1 since, as we will show, $\mathcal{I}(C) = p^n\mathfrak{o}$ for the subgroups C considered there.

Theorem 4.3. *Let F be a p-adic formal group such that $\mathrm{End}(F)$ is integrally closed. If C is a finite cyclic subgroup of $\Lambda(F)$, then $c\big(\mathrm{End}(F/C)\big) = \mathbb{Z}_p + \mathcal{I}(C)$.*

Proof. Let γ be a generator of C. By Remark 2.4(ii), $\mathcal{I}(C) = \mathcal{I}(\gamma)$. If $\zeta \in \mathcal{I}(C)$, then $[\zeta]_F(C) = \{0\} \subseteq C$, and so by Lemma 4.1, $\zeta \in c\big(\mathrm{End}(F/C)\big)$. It is now clear that $\mathbb{Z}_p + \mathcal{I}(C) \subseteq c\big(\mathrm{End}(F/C)\big)$. Conversely, take any $\zeta \in c\big(\mathrm{End}(F/C)\big)$. Then by Lemma 4.1, $[\zeta]_F(\gamma) \in C$, and so there is an $m \in \mathbb{Z}$ such that $[\zeta]_F(\gamma) = [m]_F(\gamma)$. Hence, $\zeta - m \in \mathcal{I}(\gamma) = \mathcal{I}(C)$, i.e., $\zeta \in \mathbb{Z}_p + \mathcal{I}(C)$. □

When C is a deflated (but not necessarily cyclic) subgroup of a full p-adic formal group F, we can determine the conductor of $c\big(\mathrm{End}(F)\big)$ with respect to $c\big(\mathrm{End}(F/C)\big)$. Recall that if $A \subseteq B$ are commutative unitary rings, then the *conductor of B with respect to A* is the ideal $\mathfrak{c} = \{b \in B \mid bB \subseteq A\}$.

Theorem 4.4. *Let F be a full p-adic formal group, and let C be a deflated subgroup of F. The conductor $\mathfrak{c}$ of $c\big(\mathrm{End}(F)\big)$ with respect to $c\big(\mathrm{End}(F/C)\big)$ is $\mathcal{I}(C)$.*

Proof. Let π be a uniformizer of $\mathfrak{o} = c\big(\mathrm{End}(F)\big)$. As the result is trivial if $C = \{0\}$, we may assume that $\mathcal{I}(C) = \pi^m\mathfrak{o}$ for some $m \geq 1$. Then $\mathfrak{c} = \pi^k\mathfrak{o}$, where k is the smallest nonnegative integer for which $\pi^k\mathfrak{o} \subseteq c\big(\mathrm{End}(F/C)\big)$. Now, if $\zeta \in \mathfrak{o}$, then $[\pi^m\zeta]_F(C) = [\zeta]_F\big([\pi^m]_F(C)\big) = \{0\} \subseteq C$. By Lemma 4.1, $\pi^m\zeta \in c\big(\mathrm{End}(F/C)\big)$, and so $k \leq m$. Suppose that $\pi^{m-1}\mathfrak{o} \subseteq c\big(\mathrm{End}(F/C)\big)$. Then for every $\epsilon \in \mathfrak{o}^\times$, $[\epsilon]_F\big([\pi^{m-1}]_F(C)\big) \subseteq C$. Since $\{0\} \neq [\pi^{m-1}]_F(C) \subseteq \ker[\pi]_F$, Corollary 2.7 implies that

$$\bigcup_{u\in\mathrm{Aut}(F)} u\big([\pi^{m-1}]_F(C)\big) = \ker[\pi]_F,$$

whence $\ker[\pi]_F \subseteq C$. According to Theorem 3.5, this contradicts the assumption that C is a deflated subgroup of F, and so $k = m$. Therefore, $\mathfrak{c} = \mathcal{I}(C)$. □

5. Free Tate modules of rank 1

Lubin [Lu2, §3.2] showed that if R is a $\mathbb{Z}_p$-order in a finite extension K of $\mathbb{Q}_p$ with $R \neq \mathfrak{o}_K$, then there exists an almost full p-adic formal group G with $c\big(\mathrm{End}(G)\big) = R$. However, unlike the situation for full p-adic formal groups, there do exist nonisomorphic almost full p-adic formal groups which have isomorphic absolute endomorphism rings. (We show in §6, however, that such formal groups cannot have height 2.) Waterhouse [W] proves that two almost full p-adic formal groups G_1 and G_2 are isomorphic if and only if $c\big(\mathrm{End}(G_1)\big) = c\big(\mathrm{End}(G_2)\big) = R$ and $T(G_1) \cong T(G_2)$ as R-modules. A key lemma in his proof asserts that there is an almost full p-adic formal group H with $c\big(\mathrm{End}(H)\big) = R$ such that $T(H)$ is free of rank 1 over R. In our next proposition, we use our results to derive a necessary and sufficient condition on a finite subgroup C of the points of a full p-adic formal group F which guarantees that $T(F/C)$ is free of rank 1 over $c\big(\mathrm{End}(F/C)\big)$. In the proof, we use the fact that if G is a p-adic formal group, then an element $(a_0, a_1, a_2, \dots)$ of $V(G)$ belongs to $T(G)$ if and only if $a_0 = 0$.

Proposition 5.1. *Let F be a full p-adic formal group and let C be a finite nonzero subgroup of $\Lambda(F)$. Then $T(F/C)$ is free of rank one over $c\big(\mathrm{End}(F/C)\big)$ if and only if there exists a $\gamma \in C$ satisfying the following two properties:*

(P1) *γ has minimal valuation among the elements of C.*
(P2) *If $g \in \mathrm{End}(F)$ and $g(\gamma) \in C$, then $g(C) \subseteq C$.*

Proof. Assume that $\gamma \in C$ satisfies (P1) and (P2); note that $\gamma \neq 0$ because $C \neq \{0\}$. Choose any $b \in V(F)$ such that $b_0 = \gamma$, and define $b' = V(\varphi_C)(b)$. We will show that $T(F/C) = c\big(\mathrm{End}(F/C)\big)\cdot b'$. If $\zeta \in c\big(\mathrm{End}(F/C)\big)$, then $[\zeta]_F(C) \subseteq C$ by Lemma 4.1, and hence

$$\begin{aligned}\zeta \cdot b' &= \zeta \cdot V(\varphi_C)(b)\\ &= \big([\zeta]_{F/C}(\varphi_C(b_0)), [\zeta]_{F/C}(\varphi_C(b_1)), \dots\big)\\ &= \big(\varphi_C([\zeta]_F(\gamma)), \varphi_C([\zeta]_F(b_1)), \dots\big)\\ &= (0, \dots) \in T(F/C).\end{aligned}$$

Therefore, $c\big(\mathrm{End}(F/C)\big)\cdot b' \subseteq T(F/C)$. Conversely, take any $a \in T(F/C)$, and let ζ be the unique element of Σ_F such that $a = \zeta \cdot b'$. Choose an integer n large

enough so that $p^n\zeta \in c\big(\mathrm{End}(F)\big)$. Then

$$\begin{aligned} a &= V(\varphi_C)(\zeta \cdot b) \\ &= V(\varphi_C)(p^{-n} \cdot p^n\zeta \cdot b) \\ &= \big(\varphi_C([p^n\zeta]_F(b_n)), \varphi_C([p^n\zeta]_F(b_{n+1})), \dots\big), \end{aligned}$$

which implies that $[p^n\zeta]_F(b_n) \in C$ since $a_0 = 0$. By (P1), $v(\gamma) \le v\big([p^n\zeta]_F(b_n)\big)$, and so by Lemma 2.5 there is an $\eta \in c\big(\mathrm{End}(F)\big)$ such that $[\eta]_F(\gamma) = [p^n\zeta]_F(b_n)$. Therefore, because $\gamma = [p^n]_F(b_n)$, we know that $p^n(\zeta - \eta) \in \mathcal{I}(b_n)$. However, $p^n \notin \mathcal{I}(b_n)$ (since $\gamma \neq 0$) and so $v\big(p^n(\zeta-\eta)\big) > v(p^n)$. This in turn implies that $v(\zeta - \eta) > 0$, which proves that $\zeta \in c\big(\mathrm{End}(F)\big)$. We see now that

$$\begin{aligned} a = \zeta \cdot V(\varphi_C)(b) \quad &\Longrightarrow \quad \varphi_C\big([\zeta]_F(\gamma)\big) = 0 \\ &\Longrightarrow \quad [\zeta]_F(\gamma) \in C \\ &\Longrightarrow \quad [\zeta]_F(C) \subseteq C \quad \big(\text{from (P2)}\big) \end{aligned}$$

which shows that $\zeta \in c\big(\mathrm{End}(F/C)\big)$ according to Lemma 4.1.

Now, suppose that $T(F/C)$ is free of rank 1 over $c\big(\mathrm{End}(F/C)\big)$ and choose any $b \in V(F)$ such that $\big\{V(\varphi_C)(b)\big\}$ is a $c\big(\mathrm{End}(F/C)\big)$-basis for $T(F/C)$. Because $V(\varphi_C)(b) \in T(F/C)$, it follows that $\varphi_C(b_0) = 0$, i.e., $b_0 \in C$. We will show that $\gamma = b_0$ satisfies (P1) and (P2). Take any $\delta \in C$ and $d \in V(F)$ with $d_0 = \delta$. As $V(\varphi_C)(d) \in T(F/C)$, there is a unique $\zeta \in c\big(\mathrm{End}(F/C)\big) \subseteq c\big(\mathrm{End}(F)\big)$ such that $V(\varphi_C)(d) = \zeta \cdot V(\varphi_C)(b) = V(\varphi_C)(\zeta \cdot b)$. Because $V(\varphi_C)$ is an isomorphism, $\zeta \cdot b = d$, and so $[\zeta]_F(\gamma) = \delta$. Proposition 1.2 shows that $v(\delta) \ge v(\gamma)$, which establishes (P1). Finally, if $g \in \mathrm{End}(F)$ and $g(\gamma) \in C$, then

$$c(g) \cdot V(\varphi_C)(b) = V(\varphi_C \circ g)(b) = \big(\varphi_C(g(\gamma)), \dots\big) = (0, \dots) \in T(F/C).$$

This implies that $c(g) \in c\big(\mathrm{End}(F/C)\big)$, i.e., $g(C) \subseteq C$, and so (P2) holds as well. □

Corollary 5.2. *If F is a full p-adic formal group and if C is a finite cyclic subgroup of $\Lambda(F)$, then $T(F/C)$ is free of rank 1 over $c\big(\mathrm{End}(F/C)\big)$.*

Proof. The result is clear if $C = \{0\}$. Otherwise, if $C = \langle\gamma\rangle \neq \{0\}$, then the pair (C, γ) satisfies (P1) (use Proposition 1.2) and (P2) of Proposition 5.1. □

The converse of Corollary 5.2 is not true in general, even if we require the subgroup to be deflated. Let F be a full p-adic formal group and let π be a uniformizer of $\mathfrak{o} = c\big(\mathrm{End}(F)\big)$. Fix any $0 \neq \gamma \in \Lambda(F)$ and let C be a finite subgroup of $\Lambda(F)$ containing γ as an element of minimal valuation. By Remark 2.4(ii), $\mathcal{I}(C) = \mathcal{I}(\gamma) = \pi^k\mathfrak{o}$ for some $k \in \mathbb{N}$. The set

$$\mathcal{S}_C = \big\{\zeta \in \mathfrak{o} \,\big|\, [\zeta]_F(C) \subseteq C\big\} = c\big(\mathrm{End}(F/C)\big)$$

is a subring of $\mathfrak{o}$ containing $\mathcal{I}(\gamma)$, and the set

$$\mathcal{T}_{C,\gamma} = \big\{\zeta \in \mathfrak{o} \,\big|\, [\zeta]_F(\gamma) \in C\big\}$$

is a subgroup of $\mathfrak{o}$ containing $\mathcal{S}_C$. Moreover, the evaluation map $\zeta \mapsto [\zeta]_F(\gamma)$ induces a group isomorphism $\mathcal{T}_{C,\gamma}/\mathcal{I}(\gamma) \to C$ (see Lemma 2.5). Therefore the pair (C, γ) satisfies (P1) and (P2) if and only if $\mathcal{S}_C = \mathcal{T}_{C,\gamma}$, i.e., if and only if $\overline{\mathcal{S}_C} = \mathcal{S}_C/\pi^k\mathfrak{o}$ and C have the same order. Conversely, if $\mathcal{S}$ is any subring of $\mathfrak{o}$ which contains $\mathcal{I}(\gamma)$, then we can consider the submodule $C_{\mathcal{S}} = \mathcal{S} \cdot \gamma = \big\{[\zeta]_F(\gamma) \,\big|\, \zeta \in \mathcal{S}\big\}$ of the finite $\mathcal{S}$-module $\ker[\pi^k]_F$. According to Proposition 1.2, the pair $(C_{\mathcal{S}}, \gamma)$ satisfies

(P1). Furthermore, $\mathcal{S} \subseteq \mathcal{S}_{C_\mathcal{S}} \subseteq \mathcal{T}_{C_\mathcal{S},\gamma} \subseteq \mathcal{S}$, which shows that $(C_\mathcal{S}, \gamma)$ satisfies (P2) as well. We note also that if (C, γ) satisfies (P1) and (P2), then $C_{\mathcal{S}_C} = C$. Indeed, it is clear that $C_{\mathcal{S}_C} \subseteq C$, and $C \subseteq C_{\mathcal{S}_C}$ according to Lemma 2.5 plus the fact that $\mathcal{S}_C = \mathcal{T}_{C,\gamma}$. This proves the following.

Corollary 5.3. *Let F be a full p-adic formal group. For each $0 \neq \gamma \in \Lambda(F)$, the association $C \mapsto \mathcal{S}_C$ defines a one-to-one correspondence between finite subgroups C of $\Lambda(F)$ for which the pair (C, γ) satisfies properties* (P1) *and* (P2) *of Proposition* 5.1 *and subrings of $\mathfrak{o}_{\Sigma_F}$ containing the ideal $\mathcal{I}(\gamma)$.*

In the special case where $\mathcal{I}(\gamma) = \pi\mathfrak{o}$, for any subgroup C of $\ker[\pi]_F$ containing γ, $\overline{\mathcal{S}_C}$ is a subfield of the residue field $\mathfrak{o}/\pi\mathfrak{o} = \mathbb{F}_{p^f}$. For each divisor r of f, one can use Corollary 5.3 to construct a (unique) subgroup C_r of $\ker[\pi]_F$ of order p^r such that (C_r, γ) satisfies (P1) and (P2); more specifically, $\overline{\mathcal{S}_{C_r}}$ is the subfield of $\mathbb{F}_{p^f}$ of order p^r. If f is composite and $r \neq 1$ or f, then C_r is a noncyclic deflated subgroup of F such that $T(F/C_r)$ is a free $c\big(\mathrm{End}(F/C_r)\big)$-module of rank 1.

6. Special results for height 2 formal groups

Our general results from §4 and §5 yield a wealth of information about p-adic formal groups of height 2 because of the following.

Proposition 6.1. *If F is a p-adic formal group of height* 2*, then every deflated subgroup of F is cyclic.*

Proof. Because $\ker[p]_F$ has p^2 elements, C is a product of at most two cyclic subgroups. But as C is deflated, $\ker[p]_F \nsubseteq C$. Hence $C \cap \ker[p]_F$ has at most p elements which proves that C is cyclic. □

The discussion after Theorem 3.5 shows that the converse of Proposition 6.1 is not true.

Corollary 6.2. *If G is an almost full p-adic formal group of height* 2*, then $T(G)$ is a free* $\mathrm{End}(G)$*-module of rank* 1*.*

We now give a proof of Conjecture 1.

Theorem 6.3. *Let F be a full p-adic formal group of height* 2*, and let C be a deflated (and hence cyclic) subgroup of F of order p^n. If $\mathfrak{o} = c\big(\mathrm{End}(F)\big)$, then $c\big(\mathrm{End}(F/C)\big) = \mathbb{Z}_p + p^n\mathfrak{o}$.*

Proof. The result is obvious if $C = \{0\}$, so we may assume that $n \geq 1$. By Theorem 4.3 and Remark 2.4(ii), it suffices to show that $\mathcal{I}(\gamma) = p^n\mathfrak{o}$, where γ is a generator of C. Clearly, $[p^n]_F(\gamma) = 0$ and $[p^{n-1}]_F(\gamma) \neq 0$. If $\Sigma_F/\mathbb{Q}_p$ is unramified, then p is a uniformizer of $\mathfrak{o}_{\Sigma_F}$, which shows that $\mathcal{I}(\gamma) = p^n\mathfrak{o}$ in this case. On the other hand, if $\Sigma_F/\mathbb{Q}_p$ is totally ramified and if π is a uniformizer of $\mathfrak{o}$, then either p^n or πp^{n-1} generates $\mathcal{I}(\gamma)$. If $[\pi p^{n-1}]_F(\gamma) = 0$, then $[p^{n-1}]_F(\gamma)$ would be a nonzero element of $\ker[\pi]_F \cap C$, which would imply that $\ker[\pi]_F \subseteq C$ because $\ker[\pi]_F$ is cyclic. This contradicts the assumption that C is a deflated subgroup of F, and so $\mathcal{I}(\gamma) = p^n\mathfrak{o}$ in this case as well. □

Finally, as an application, we use our results to show that the isomorphism class of an almost full p-adic formal group of height 2 depends only on its absolute endomorphism ring. This is a generalization in height 2 of [Lu3, 4.3.2].

Corollary 6.4. *Let G_1 and G_2 be almost full p-adic formal groups of height 2 such that $c\big(\mathrm{End}(G_1)\big) = c\big(\mathrm{End}(G_2)\big)$. Then G_1 and G_2 are isomorphic via an isogeny.*

Proof. Using Corollary 1.8 and the results in §3, we can find full p-adic formal groups F_1 and F_2 and deflated subgroups C_1 and C_2 of F_1 and F_2 respectively such that $F_1/C_1 \cong G_1$ and $F_2/C_2 \cong G_2$. Since $\Sigma_{F_1} = \Sigma_{G_1} = \Sigma_{G_2} = \Sigma_{F_2}$, we may assume without loss of generality that $F_1 = F_2 = F$ [Lu3, 4.3.2]. Then, according to Theorem 4.4, the fact that $c\big(\mathrm{End}(G_1)\big) = c\big(\mathrm{End}(G_2)\big)$ implies that $\mathcal{I}(C_1) = \mathcal{I}(C_2)$. Since C_1 and C_2 are cyclic, it follows from Corollary 2.8 that there exists some $u \in \mathrm{Aut}(F)$ such that $C_1 = u(C_2)$. Therefore, $C_1 \sim C_2$ by Proposition 3.1, whence $G_1 \cong G_2$ by definition. That this isomorphism is an isogeny follows from Corollary 1.7. □

Remark 6.5. We could have instead used the main theorem in [W] to prove Corollary 6.4. Indeed, for $i = 1, 2$, C_i is cyclic, and therefore the Tate module $T(G_i)$ is free of rank 1 over $R = c\big(\mathrm{End}(G_i)\big)$, according to Corollary 5.2. So, certainly $T(G_1)$ and $T(G_2)$ are isomorphic as R-modules.

References

[F] Fröhlich, A. Formal groups. Lecture Notes in Mathematics, 74. *Springer-Verlag, Berlin-New York*, 1968. MR0242837 (39 #4164), Zbl 0177.04801.

[Lang] Lang, Serge. Algebra, Second edition. *Addison-Wesley Publishing Company, Advanced Book Program, Reading, MA*, 1984. MR0783636 (86j:00003), Zbl 0712.00001.

[Laz] Lazard, Michel. Sur les groupes de Lie formels à un paramétre. *Bull. Soc. Math. France* **83** (1955), 251–274. MR0073925 (17,508e), Zbl 0068.25703.

[Lu1] Lubin, Jonathan. Canonical subgroups of formal groups. *Trans. Amer. Math. Soc.* **251** (1979), 103–127, MR0531971 (80j:14039), Zbl 0431.14014.

[Lu2] Lubin, Jonathan. Finite subgroups and isogenies of one-parameter formal Lie groups. *Ann. of Math.* **85** (1967), 296–302. MR0209287 (35 #189), Zbl 0166.02803.

[Lu3] Lubin, Jonathan. One-parameter formal Lie groups over $\mathfrak{p}$-adic integer rings. *Ann. of Math.* **80** (1964), 464–484. MR0168567 (29 #5827), Zbl 0135.07003.

[LT] Lubin, Jonathan; Tate, John. Formal complex multiplication in local fields. *Ann. of Math.* **81** (1965), 380–387, MR0172878 (30 #3094), Zbl 0128.26501.

[S] Schmitz, David. Canonical and filling subgroups of formal groups. *New York J. Math.* **12** (2006), 235–247.

[W] Waterhouse, William C. A classification of almost full formal groups. *Proc. Amer. Math. Soc.* **20** (1969), 426–428. MR0236189 (38 #4487), Zbl 0176.30303.

DEPARTMENT OF MATHEMATICS, NORTH CENTRAL COLLEGE, 30 N BRAINARD ST, NAPERVILLE, IL 60540

djschmitz@noctrl.edu

This paper is available via http://nyjm.albany.edu/j/2006/12-13.html.