

# Unbounded strongly irreducible operators and transitive representations of quivers on infinite-dimensional Hilbert spaces

Masatoshi Enomoto and Yasuo Watatani

ABSTRACT. We introduce unbounded strongly irreducible operators and transitive operators. These operators are related to a certain class of indecomposable Hilbert representations of quivers on infinite-dimensional Hilbert spaces. We regard the theory of Hilbert representations of quivers as a generalization of the theory of unbounded operators. A non-zero Hilbert representation of a quiver is said to be transitive if the endomorphism algebra is trivial. If a Hilbert representation of a quiver is transitive, then it is indecomposable. But the converse is not true. Let  $\Gamma$  be a quiver whose underlying undirected graph is an extended Dynkin diagram. Then there exists an infinite-dimensional transitive Hilbert representation of  $\Gamma$  if and only if  $\Gamma$  is not an oriented cyclic quiver.

## CONTENTS

1. Introduction	975
2. Hilbert representations of quivers	978
3. Unbounded strongly irreducible operators	981
4. Extended Dynkin diagrams and transitive Hilbert representations	988
References	1013

## 1. Introduction

A bounded linear operator  $T$  on a Hilbert space  $H$  is called strongly irreducible if  $T$  cannot be decomposed to a non-trivial (not necessarily orthogonal) direct sum of two operators, that is, if there exist no non-trivial invariant closed subspaces  $M$  and  $N$  of  $T$  such that  $M \cap N = 0$  and  $M + N = H$ . A strongly irreducible operator is an infinite-dimensional generalization of

---

Received April 10, 2017.

2010 *Mathematics Subject Classification*. Primary 47A65, Secondary 46C07, 47A15, 16G20.

*Key words and phrases*. unbounded strongly irreducible operators, transitive operators, quiver, indecomposable representation, Hilbert space.

This work was supported by JSPS KAKENHI Grant Numbers 23654053 and 25287019.

a Jordan block. F. Gilfeather [Gi] introduced the notion of strongly irreducible operators. We refer to excellent books [JiW1] and [JiW2] by Jiang and Wang on strongly irreducible operators.

In [EW1, EW2] we studied the relative positions of subspaces in a separable infinite-dimensional Hilbert space after Nazarova [Na1], Gelfand and Ponomarev [GeP]. We think that relative positions of subspaces have a close relation with subfactor theory [Jo, GoHJ].

Let  $H$  be a Hilbert space and  $E_1, \dots, E_n$  be  $n$  subspaces in  $H$ . Then it is said that  $\mathcal{S} = (H; E_1, \dots, E_n)$  is a system of  $n$  subspaces in  $H$  or an  $n$ -subspace system in  $H$ . Two systems  $\mathcal{S} = (H; E_1, \dots, E_n)$  and  $\mathcal{T} = (K; F_1, \dots, F_n)$  are isomorphic if there exists an invertible operator  $\varphi : H \rightarrow K$  such that  $\varphi(E_i) = F_i$  for  $i = 1, 2, \dots, n$ . A non-zero system  $\mathcal{S} = (H; E_1, \dots, E_n)$  is said to be indecomposable if it cannot be decomposed to a non-trivial direct sum of two systems up to isomorphism. We recall that strongly irreducible operators contribute an important role to construct indecomposable systems of four subspaces [EW1].

On the other hand, Gabriel [Ga] introduced a finite-dimensional (linear) representation of quivers by attaching vector spaces and linear maps for vertices and edges of quivers respectively. A finite-dimensional indecomposable representation of a quiver is a direct graph generalization of a Jordan block. Historically, Kronecker [Kro] solved the indecomposable representations of  $\tilde{A}_1$ , the so called matrix pencils in 1890. Nazarova [Na1] and Gelfand-Ponomarev [GeP] treated the four-subspace situation  $\tilde{D}_4$ . Donovan-Freislich [DoF] and Nazarova [Na2] classified the indecomposable representations of the tame quivers. About these topics we also refer to Bernstein-Gelfand-Ponomarev [BGP], V. Dlab-Ringel [DIR], Ringel [Ri2], Gabriel-Roiter [GaR], Kac [Ka], and so on.

We recall infinite-dimensional representations in purely algebraic setting. In [Au] Auslander found that if a finite-dimensional algebra is not of finite representation type, then there exist indecomposable modules which are not of finite length. These are trivially infinite-dimensional. Several works about infinite-dimensional Kronecker modules have been done by N. Aronszjan, A. Dean, U. Fixman, F. Okoh and F. Zorzitto in [Ar, DeZ1, Fi, FiO, FiZ, Ok]. A. Dean and F. Zorzitto [DeZ2] constructed a family of infinite-dimensional indecomposable representations of  $\tilde{D}_4$ . K. Ringel [Ri1] founded a general theory of infinite-dimensional representations of tame, hereditary algebra (see also [Ri3, KrR]).

In [EW3, E] we started to investigate the representation theory of quivers on Hilbert spaces. We asked the existence of an indecomposable infinite-dimensional Hilbert representation for any quiver whose underlying undirected graph is one of extended Dynkin diagrams. And we solved it affirmatively using the unilateral shift  $S$ . The argument works even if we replace the unilateral shift  $S$  with any strongly irreducible operator. From this,

it is suggested that strongly irreducible operators are useful to construct indecomposable Hilbert representations of quivers [EW4].

From the analogy of a transitive lattice (see P.R. Halmos [H] and K.J. Harrison, H. Radjavi and P. Rosenthal [HRR]), we called an indecomposable Hilbert representation  $(H, f)$  of a quiver such that  $\text{End}(H, f) = \mathbb{C}I$  transitive. If a Hilbert representation of a quiver is transitive, then it is indecomposable. But the converse is not true. Therefore, it is important to investigate the existence problem of an transitive infinite-dimensional Hilbert representation for any quiver whose underlying undirected graph is one of extended Dynkin diagrams. In this direction, we [EW4] showed two kinds of constructions of quite non-trivial transitive Hilbert representations  $(H, f)$  of the Kronecker quiver.

In the purely algebraic setting, a representation of a quiver is called a brick if its endomorphism ring is a division ring. But for a Hilbert representation  $(H, f)$ ,  $\text{End}(H, f)$  is a Banach algebra and not isomorphic to its purely algebraic endomorphism ring in general, because we only consider bounded endomorphisms. By the Gelfand-Mazur theorem, any Banach algebra over  $\mathbb{C}$  which is a division ring must be isomorphic to  $\mathbb{C}$ .

We remark that locally scalar representations of quivers were introduced by Kruglyak and Roiter [KrRo]. But their subject is different from ours. We also refer to S. Kruglyak, V. Rabanovich, and Y. Samoilenko [KrRS] and Y. P. Moskaleva and Y. S. Samoilenko [MS].

We consider finite-dimensional indecomposable representations of quivers whose underlying graph is a Dynkin diagram. They are transitive (cf.[As]). But it is extremely difficult to solve the existence problem for infinite-dimensional indecomposable (also transitive) Hilbert representations of quivers whose underlying undirected graph is a Dynkin diagram. The existence is not known even for quivers whose underlying undirected graph is  $D_4$ .

In this paper we introduce unbounded strongly irreducible operators and transitive operators. It is known that any unbounded closed operator  $T$  on a Hilbert space can be realized as a quotient  $BA^{-1}$  of bounded operators  $A$  and  $B$  on  $H$ . This fact is related with operator ranges and intersections of domains of unbounded operators. See, for example, P. Fillmore and J. Williams [FiW], W.E. Kaufman [Kau] and H. Kosaki [Ko]. We point out that the study of an unbounded closed operator  $T = BA^{-1}$  can be translated to the study of a Hilbert representation given by  $A$  and  $B$  of the Kronecker quiver. We show that some transitive operators are constructed by a certain transitive Hilbert representation of the Kronecker quiver. We regard the theory of Hilbert representations of quivers as a generalization of the theory of unbounded operators. We also solve completely the existence problem of infinite-dimensional transitive Hilbert representations of quivers whose underlying undirected graphs are the extended Dynkin diagrams.

Let  $\Gamma$  be a quiver whose underlying undirected graph is an extended Dynkin diagram. If the underlying undirected graph of  $\Gamma$  is not  $\widetilde{A}_n$ , then there exists an infinite-dimensional transitive Hilbert representation of  $\Gamma$ . If the underlying undirected graph of  $\Gamma$  is  $\widetilde{A}_n$ , then there exists an infinite-dimensional transitive Hilbert representation of  $\Gamma$  if and only if  $\Gamma$  is not an oriented cyclic quiver. We used unbounded transitive operators based on an idea of a transitive lattice by K.J. Harrison, H. Radjavi and P. Rosenthal ([HRR],[RR]).

## 2. Hilbert representations of quivers

A quiver  $\Gamma = (V, E, s, r)$  is a quadruple consisting of the set  $V$  of vertices, the set  $E$  of arrows, and two maps  $s, r : E \rightarrow V$  which associate with each arrow  $\alpha \in E$  its support  $s(\alpha)$  and range  $r(\alpha)$ . In this paper we assume that  $\Gamma$  is a finite quiver.

We denote by  $\alpha : x \rightarrow y$  an arrow with  $x = s(\alpha)$  and  $y = r(\alpha)$ . Thus a quiver is a directed graph. We denote by  $|\Gamma|$  the underlying undirected graph of a quiver  $\Gamma$ . We say that a quiver  $\Gamma$  is connected if  $|\Gamma|$  is a connected graph. A quiver  $\Gamma$  is called finite if both  $V$  and  $E$  are finite sets. A path of length  $m$  is a finite sequence  $\alpha = (\alpha_1, \dots, \alpha_m)$  of arrows such that  $r(\alpha_k) = s(\alpha_{k+1})$  for  $k = 1, \dots, m-1$ . Its support is  $s(\alpha) = s(\alpha_1)$  and its range is  $r(\alpha) = r(\alpha_m)$ . A path of length  $m \geq 1$  is called a cycle if its support and range coincide. A cycle of length one is called a loop. A quiver which is a loop is also called the Jordan quiver  $L$ . A quiver which is a cycle of length  $m \geq 1$  is also called the oriented cyclic quiver  $C_m$  with length  $m \geq 1$ . A quiver is said to be acyclic if it contains no cycles.

**Definition.** Let  $\Gamma = (V, E, s, r)$  be a finite quiver. It is said that  $(H, f)$  is a Hilbert representation of  $\Gamma$  if  $H = (H_v)_{v \in V}$  is a family of Hilbert spaces and  $f = (f_\alpha)_{\alpha \in E}$  is a family of bounded linear operators with  $f_\alpha : H_{s(\alpha)} \rightarrow H_{r(\alpha)}$ .

**Definition.** Let  $\Gamma = (V, E, s, r)$  be a finite quiver. Let  $(H, f)$  and  $(K, g)$  be Hilbert representations of  $\Gamma$ . A homomorphism  $T : (H, f) \rightarrow (K, g)$  is a family  $T = (T_v)_{v \in V}$  of bounded operators  $T_v : H_v \rightarrow K_v$  satisfying  $T_{r(\alpha)} f_\alpha = g_\alpha T_{s(\alpha)}$  for any arrow  $\alpha \in E$ .

The composition  $T \circ S$  of homomorphisms  $T$  and  $S$  is defined by  $(T \circ S)_v = T_v \circ S_v$  for  $v \in V$ . In this way we have obtained a category  $\text{HRep}(\Gamma)$  of Hilbert representations of  $\Gamma$ . We denote by  $\text{Hom}((H, f), (K, g))$  the set of homomorphisms  $T : (H, f) \rightarrow (K, g)$ . We denote by  $\text{End}(H, f) := \text{Hom}((H, f), (H, f))$  the set of endomorphisms. We can regard  $\text{End}(H, f)$  as a subalgebra of  $\bigoplus_{v \in V} B(H_v)$ .

In the paper we distinguish the following two classes of operators. A bounded operator  $A$  is said to be a projection (resp. an idempotent) if  $A^2 =$

$A = A^*$  (resp.  $A^2 = A$ ). We denote by

$$\begin{aligned} Idem(H, f) &:= \{T \in End(H, f) \mid T^2 = T\} \\ &= \{T = (T_v)_{v \in V} \in End(H, f) \mid T_v^2 = T_v \text{ (for any } v \in V)\} \end{aligned}$$

the set of all idempotents of  $End(H, f)$ .

Let  $0 = (0_v)_{v \in V}$  be a family of zero endomorphisms and  $I = (I_v)_{v \in V}$  be a family of identity endomorphisms. It is said that  $(H, f)$  and  $(K, g)$  are isomorphic, denoted by  $(H, f) \cong (K, g)$ , if there exists an isomorphism  $\varphi : (H, f) \rightarrow (K, g)$ , that is, there exists a family  $\varphi = (\varphi_v)_{v \in V}$  of bounded invertible operators  $\varphi_v \in B(H_v, K_v)$  such that  $\varphi_{r(\alpha)} f_\alpha = g_\alpha \varphi_{s(\alpha)}$  for any arrow  $\alpha \in E$ . We say that  $(H, f)$  is a finite-dimensional representation if  $H_v$  is finite-dimensional for all  $v \in V$ . And  $(H, f)$  is an infinite-dimensional representation if  $H_v$  is infinite-dimensional for some  $v \in V$ .

We recall a notion of indecomposable representation in [EW3], that is, a representation which cannot be decomposed into a direct sum of smaller representations anymore.

**Definition.** Let  $\Gamma = (V, E, s, r)$  be a finite quiver. Let  $(K, g)$  and  $(K', g')$  be Hilbert representations of  $\Gamma$ . We define the direct sum  $(H, f) = (K, g) \oplus (K', g')$  by  $H_v = K_v \oplus K'_v$  for  $v \in V$  and  $f_\alpha = g_\alpha \oplus g'_\alpha$  for  $\alpha \in E$ . It is said that a Hilbert representation  $(H, f)$  is zero, denoted by  $(H, f) = 0$  if  $H_v = 0$  for any  $v \in V$ .

**Definition.** A Hilbert representation  $(H, f)$  of  $\Gamma$  is said to be decomposable if  $(H, f)$  is isomorphic to a direct sum of two non-zero Hilbert representations. A non-zero Hilbert representation  $(H, f)$  of  $\Gamma$  is called indecomposable if it is not decomposable, that is, if  $(H, f) \cong (K, g) \oplus (K', g')$  then  $(K, g) \cong 0$  or  $(K', g') \cong 0$ .

The following proposition is useful to show the indecomposability in concrete examples.

**Proposition 2.1.** [EW3, Proposition 3.1.] *Let  $(H, f)$  be a Hilbert representation of a quiver  $\Gamma$ . Then the following conditions are equivalent:*

- (1)  $(H, f)$  is indecomposable.
- (2)  $Idem(H, f) = \{0, I\}$ .

**Remark.** The indecomposability of Hilbert representations of a quiver is an isomorphic invariant, but it is not a unitary invariant. Hence we cannot replace the set  $Idem(H, f)$  of idempotents of endomorphisms by the subset of idempotents of endomorphisms which consists of projections to show the indecomposability.

**Definition.** ([EW4, page 569]) A Hilbert representation  $(H, f)$  of a quiver  $\Gamma$  is said to be *transitive* if  $End(H, f) = \mathbb{C}I$ .

If a Hilbert representation  $(H, f)$  of  $\Gamma$  is transitive, then  $(H, f)$  is indecomposable. In fact, since  $End(H, f) = \mathbb{C}I$ , any idempotent endomorphism  $T$  is 0 or  $I$ . In purely algebraic setting, a representation of a quiver is said

to be a *brick* if its endomorphism ring is a division ring (see for example, cf. [As]).

Let  $H$  be a Hilbert space and  $E_1, \dots, E_n$  be  $n$  subspaces in  $H$ . Then it is said that  $\mathcal{S} = (H; E_1, \dots, E_n)$  is a system of  $n$  subspaces in  $H$ . Let  $\mathcal{T} = (K; F_1, \dots, F_n)$  be another system of  $n$  subspaces in a Hilbert space  $K$ . Then we say that  $\varphi : \mathcal{S} \rightarrow \mathcal{T}$  is a homomorphism if  $\varphi : H \rightarrow K$  is a bounded linear operator satisfying that  $\varphi(E_i) \subset F_i$  for  $i = 1, \dots, n$ . We say that  $\varphi : \mathcal{S} \rightarrow \mathcal{T}$  is an isomorphism if  $\varphi : H \rightarrow K$  is an invertible (i.e., bounded bijective) linear operator satisfying that  $\varphi(E_i) = F_i$  for  $i = 1, \dots, n$ . It is said that systems  $\mathcal{S}$  and  $\mathcal{T}$  are *isomorphic* if there is an isomorphism  $\varphi : \mathcal{S} \rightarrow \mathcal{T}$ . This means that the relative positions of  $n$  subspaces  $(E_1, \dots, E_n)$  in  $H$  and  $(F_1, \dots, F_n)$  in  $K$  are same under disregarding angles. Let us denote by  $Hom(\mathcal{S}, \mathcal{T})$  the set of homomorphisms of  $\mathcal{S}$  to  $\mathcal{T}$  and  $End(\mathcal{S}) := Hom(\mathcal{S}, \mathcal{S})$  the set of endomorphisms on  $\mathcal{S}$ . Let  $\mathcal{S} = (H; E_1, \dots, E_n)$  and  $\mathcal{S}' = (H'; E'_1, \dots, E'_n)$  be systems of  $n$  subspaces in Hilbert spaces  $H$  and  $H'$ . Then their direct sum  $\mathcal{S} \oplus \mathcal{S}'$  is defined by

$$\mathcal{S} \oplus \mathcal{S}' := (H \oplus H'; E_1 \oplus E'_1, \dots, E_n \oplus E'_n).$$

A system  $\mathcal{S} = (H; E_1, \dots, E_n)$  of  $n$  subspaces is said to be *decomposable* if the system  $\mathcal{S}$  is isomorphic to a direct sum of two non-zero systems. A non-zero system  $\mathcal{S} = (H; E_1, \dots, E_n)$  of  $n$  subspaces is called *indecomposable* if it is not decomposable.

We recall that strongly irreducible operators  $A$  play an extremely important role to construct indecomposable systems of four subspaces. Moreover the commutant  $\{A\}'$  corresponds to the endomorphism ring.

For any single operator  $A \in B(K)$  on a Hilbert space  $K$ , let  $\mathcal{S}_A = (H; E_1, E_2, E_3, E_4)$  be the associated operator system such that  $H = K \oplus K$  and

$$E_1 = K \oplus 0, E_2 = 0 \oplus K, E_3 = \{(x, Ax); x \in K\}, E_4 = \{(y, y); y \in K\}.$$

It follows that

$$End(\mathcal{S}_A) = \{T \oplus T \in B(H); T \in B(K), AT = TA\}$$

is isomorphic to the commutant  $\{A\}'$ . The associated system  $\mathcal{S}_A$  of four subspaces is indecomposable if and only if  $A$  is strongly irreducible. Moreover for any operators  $A, B \in B(K)$  on a Hilbert space  $K$ , the associated systems  $\mathcal{S}_A$  and  $\mathcal{S}_B$  are isomorphic if and only if  $A$  and  $B$  are similar.

Following [H] and [HRR], we [EW1, page 272] introduced a transitive system of subspaces. A system  $\mathcal{S} = (H; E_1, E_2, \dots, E_n)$  of  $n$  subspaces in a Hilbert space is called transitive if the endomorphism algebra is trivial, that is,

$$End(\mathcal{S}) = \{A \in B(H); A(E_i) \subset E_i \text{ for any } i = 1, 2, \dots, n\} = \mathbb{C}I.$$

### 3. Unbounded strongly irreducible operators

In this section we shall introduce unbounded strongly irreducible operators and transitive operators. These operators are related to a certain class of indecomposable Hilbert representations of quivers on infinite-dimensional Hilbert spaces and four- subspace systems. Let  $H$  be a Hilbert space and  $A$  a bounded linear operator on  $H$ . We denote the image of  $A$  by  $\text{Im}(A)$  and the graph of  $A$  by  $G(A)$ , that is,  $G(A) = \{(x, Ax); x \in H\}$ . For elements  $x, y \in H$ , we define a rank-one operator  $\theta_{x,y}$  by  $\theta_{x,y}(z) = (z|y)x$  for  $z \in H$ .

P.R. Halmos [H] initiated the study of transitive lattices. A lattice  $\mathcal{L}$  of subspaces of a Hilbert space  $H$  containing  $0$  and  $H$  is called a transitive lattice if

$$\{A \in B(H); AM \subset M \text{ for any } M \in \mathcal{L}\} = \mathbb{C}I.$$

K.J. Harrison, H. Radjavi and P. Rosenthal ([HRR]) constructed a transitive subspace lattice using an unbounded weighted shift as follows: Let  $K = \ell^2(\mathbb{Z})$  be a Hilbert space with an orthogonal basis  $\{e_i\}_{i=-\infty}^{+\infty}$ . Let

$$w_n = 1 \quad (n \leq 0), \quad w_n = \exp((-1)^n n!) \quad (n > 0).$$

Let  $T$  be the bilateral weighted shift defined by  $Te_n = w_n e_{n+1}$ , with the domain

$$D(T) = \left\{x = \sum_{i=-\infty}^{+\infty} \alpha_i e_i; \sum_{i=-\infty}^{+\infty} |\alpha_i w_i|^2 < +\infty\right\}.$$

Put  $E_1 = K \oplus 0, E_2 = 0 \oplus K, E_3 = G(T), E_4 = \{(x, x); x \in K\}$ . Their transitive lattice is  $\mathcal{L} = \{0, H = K \oplus K, E_1, E_2, E_3, E_4\}$ . See also the book of Radjavi-Rosenthal [RR, 4.7. page 78].

We [EW4] considered a finite subspace lattice as a Hilbert representation of a quiver  $\Gamma$  as follows. Let  $\mathcal{L} = \{0, M_1, M_2, \dots, M_n, H\}$  be a finite lattice. Consider an  $n$  subspace quiver  $R_n = (V, E, s, r)$ , that is,  $V = \{1, 2, \dots, n, n+1\}$  and  $E = \{\alpha_k; k = 1, \dots, n\}$  with  $s(\alpha_k) = k$  and  $r(\alpha_k) = n+1$  for  $k = 1, \dots, n$ . Then there exists a Hilbert representation  $(K, f)$  of  $R_n$  such that  $K_k = M_k, K_{n+1} = H$  and  $f_{\alpha_k} : M_k \rightarrow H$  is an inclusion for  $k = 1, \dots, n$ . The lattice  $\mathcal{L}$  is transitive if and only if the corresponding Hilbert representation  $(K, f)$  is transitive. By this fact we may use the terminology “transitive” in the Hilbert representation case.

We recall some facts on strongly irreducible operators for convenience.

**Lemma 3.1.** *Let  $A$  be a bounded operator on a Hilbert space  $H$ . Then the following three conditions are equivalent:*

- (0) *For any closed subspaces  $M$  and  $N$  of  $H$  with  $H = M + N$  and  $M \cap N = 0$ , if  $AM \subset M$  and  $AN \subset N$ , then  $M = 0$  or  $N = 0$ .*
- (1) *If  $T \in B(H)$  is an idempotent in the commutant  $\{A\}'$  of  $A$ , then  $T = 0$  or  $T = I$ .*
- (2) *If  $T \in B(H)$  is an idempotent such that  $(T \oplus T)(G(A)) \subset G(A)$ , then  $T = 0$  or  $T = I$ .*

**Proof.** Let  $M$  and  $N$  be closed subspaces of  $H$  such that  $H = M + N$  and  $M \cap N = 0$ . Then there exists an idempotent  $E$  such that  $M = E(H)$  and  $N = (I - E)H$ . Hence (0) is equivalent to (1).

We shall show that (1) is equivalent to (2). Assume that (1) holds. Let  $T \in B(H)$  be an idempotent such that  $(T \oplus T)(G(A)) \subset G(A)$ . Then for any  $x \in H$ , there exists  $y \in H$  such that  $(T \oplus T)((x, Ax)) = (y, Ay)$ . Hence  $Tx = y$  and  $TAx = Ay$ . Thus  $TA = AT$ . Hence  $T \in \{A\}'$ . Since  $T$  is an idempotent,  $T = 0$  or  $T = I$ . Hence (2) holds. Next we assume that (2) holds. Take an idempotent  $T \in \{A\}' \cap B(H)$ . Then

$$(T \oplus T)((x, Ax)) = (Tx, TAx) = (Tx, ATx).$$

Thus  $(T \oplus T)(G(A)) \subset G(A)$ . We have  $T = 0$  or  $T = I$ . Hence (1) holds.  $\square$

**Definition.** A bounded operator  $A \in B(H)$  is said to be strongly irreducible if  $A$  satisfies one of the three conditions of the above lemma.

Inspired by the example of K.J. Harrison, H. Radjavi and P. Rosenthal, we introduce unbounded strongly irreducible operators and unbounded transitive operators.

**Definition.** Let  $A$  be an unbounded closed operator on a Hilbert space  $H$  with the domain  $D(A) \subset H$ . We define the (bounded) commutant  $\{A\}'$  of  $A$  by  $\{A\}' = \{S \in B(H); S(D(A)) \subset D(A) \text{ and, for any } x \in D(A), ASx = SAx\}$ . See for example [Ak, §17].

Let  $A$  and  $B$  be unbounded closed operators on  $H$ . We say that  $A$  and  $B$  are similar if there exists a bounded invertible operator  $T \in B(H)$  such that  $T(D(A)) = D(B)$  and  $B = TAT^{-1}$ . We say that  $A$  is an orthogonal direct sum  $A_1 \oplus A_2$  of operators  $A_1$  and  $A_2$  on  $H = H_1 \oplus H_2$  if  $D(A) = \{(x_1, x_2); x_1 \in D(A_1), x_2 \in D(A_2)\}$  and  $Ax = (A_1x_1, A_2x_2)$  for  $x = (x_1, x_2) \in D(A)$ .

**Lemma 3.2.** *Let  $A$  be an unbounded closed operator on a Hilbert space  $H$  with the domain  $D(A) \subset H$ . Then the following three conditions are equivalent:*

- (0) *If  $A$  is similar to  $A_1 \oplus A_2$  on  $H = H_1 \oplus H_2$  for some unbounded closed operators  $A_1$  and  $A_2$ , then  $H_1 = 0$  or  $H_2 = 0$ .*
- (1) *For any idempotent  $E \in B(H)$ , if  $E$  is in the commutant  $\{A\}'$ , then  $E = 0$  or  $E = I$ .*
- (2) *For any idempotent  $E \in B(H)$ , if  $(E \oplus E)(G(A)) \subset G(A)$ , then  $E = 0$  or  $E = I$ .*

**Proof.** We shall show that (0) $\Rightarrow$ (1). Let  $E \in \{A\}'$  be an idempotent. We have  $E(D(A)) \subset D(A)$  and  $AEx = EAx$  for  $x \in D(A)$ . There exists an invertible operator  $T \in B(H)$  such that  $T(E(H)) = H_1$  and  $T((I - E)H) = H_2$  and  $H = H_1 \oplus H_2$ . We define  $A_1x = TAT^{-1}x = TAET^{-1}x$  for  $x \in T(E(D(A))) \subset H_1$ . Since  $E(D(A)) \subset D(A)$ ,  $A_1$  is well defined. And  $A_1$  is an operator from  $T(E(D(A)))$  to  $H_1$  by  $AEx = EAx$  for  $x \in D(A)$ . We define  $A_2x = TAT^{-1}x = TA(I - E)T^{-1}x$  for  $x \in T((I - E)(D(A))) \subset H_2$ .



Since  $E(D(A)) \subset D(A)$ ,  $A_2$  is well defined. And  $A_2$  is an operator from  $T((I - E)(D(A)))$  to  $H_2$  by  $AEx = EAx$  for  $x \in D(A)$ . Hence we have

$$TAT^{-1} = TAET^{-1} + TA(I - E)T^{-1} = A_1 \oplus A_2.$$

Hence  $A \cong A_1 \oplus A_2$  on  $H_1 \oplus H_2$ . Since (0) holds, we have  $H_1 = 0$  or  $H_2 = 0$ . Hence  $TE(H) = 0$  or  $T(I - E)(H) = 0$ . So  $E = 0$  or  $E = I$ . Thus we have (0) $\Rightarrow$ (1).

Conversely we shall show that (1) $\Rightarrow$ (0). Assume that  $A \cong A_1 \oplus A_2$  on  $H_1 \oplus H_2$  for some unbounded closed operators  $A_1$  and  $A_2$ . There exists an invertible operator  $T \in B(H)$  such that  $TAT^{-1}x = (A_1 \oplus A_2)x$  for  $x \in D(A_1 \oplus A_2) = T(D(A))$ . There exists an idempotent  $E \in B(H)$  such that  $T^{-1}H_1 = E(H)$  and  $T^{-1}H_2 = (I - E)H$ . We shall show that  $E(D(A)) \subset D(A)$  and  $AE = EA$  on  $D(A)$ . We have  $T^{-1}D(A_1) \subset T^{-1}H_1 = EH$  and  $T^{-1}D(A_2) \subset T^{-1}H_2 = (I - E)H$ .  $D(A) = T^{-1}D(A_1 \oplus A_2) = T^{-1}D(A_1) + T^{-1}D(A_2)$ .

$$\begin{aligned} E(D(A)) &= E(T^{-1}D(A_1) + T^{-1}D(A_2)) = T^{-1}D(A_1) \\ &\subset T^{-1}D(A_1) + T^{-1}D(A_2) = D(A). \end{aligned}$$

For  $x \in D(A)$ , we can write  $x = x_1 + x_2$  with  $x_1 \in T^{-1}D(A_1)$  and  $x_2 \in T^{-1}D(A_2)$ . We have

$$\begin{aligned} AEx &= (T^{-1}(A_1 \oplus A_2)T)E(x_1 + x_2) \\ &= (T^{-1}(A_1 \oplus A_2)T)x_1 = T^{-1}A_1Tx_1 \end{aligned}$$

and

$$\begin{aligned} EAx &= E(T^{-1}(A_1 \oplus A_2)T)(x_1 + x_2) \\ &= E(T^{-1}A_1Tx_1 + T^{-1}A_2Tx_2) \\ &= T^{-1}A_1Tx_1. \end{aligned}$$

Thus we have  $AE = EA$  on  $D(A)$ . Therefore  $E = 0$  or  $E = I$ . Hence  $H_1 = 0$  or  $H_2 = 0$ .

Next, we shall show that (1) $\Rightarrow$ (2). Let  $E \in B(H)$  be an idempotent such that  $(E \oplus E)(G(A)) \subset G(A)$ . Then for any  $x \in D(A)$ , there exists  $y \in D(A)$  such that  $(E \oplus E)(x, Ax) = (y, Ay)$ . Hence

$$(Ex, EAx) = (y, Ay) = (Ex, AEx).$$

Thus  $E \in \{A\}'$ . By (1), then  $E = 0$  or  $E = I$ .

Conversely, we shall show that (2) $\Rightarrow$ (1). Let  $E \in \{A\}'$  be an idempotent. Then  $E(D(A)) \subset D(A)$ ,  $EAx = AEx$  for  $x \in D(A)$ , and

$$(E \oplus E)((x, Ax)) = (Ex, EAx) = (Ex, AEx).$$

Hence  $(E \oplus E)(G(A)) \subset G(A)$ , and  $E = 0$  or  $E = I$ . □

**Definition.** An unbounded closed operator  $A$  is said to be strongly irreducible if  $A$  satisfies one of the three conditions of the above lemma.

The next lemma is proved similarly.

**Lemma 3.3.** *Let  $A$  be an unbounded closed operator on a Hilbert space  $H$  with the domain  $D(A) \subset H$ . Then the following two conditions are equivalent:*

- (1) *For any  $T \in B(H)$ , if  $T$  is in the commutant  $\{A\}'$ , then  $T$  is a scalar operator.*
- (2) *For any  $T \in B(H)$ , if  $(T \oplus T)(G(A)) \subset G(A)$ , then  $T$  is a scalar operator.*

**Definition.** An unbounded closed operator  $A$  is said to be transitive if  $A$  satisfies one of the two conditions of the above lemma.

If an unbounded closed operator  $A$  is transitive, then  $A$  is strongly irreducible. Any bounded strongly irreducible operator  $A$  on a Hilbert space  $H$  with  $\dim H \geq 2$  is not transitive, because  $A \in \{A\}'$ .

By the same argument we have the following lemma.

**Lemma 3.4.** *Let  $A$  be an unbounded closed operator on a Hilbert space  $K$  with the domain  $D(A)$ . Let  $\mathcal{S}_A = (H; E_1, E_2, E_3, E_4)$  be a four-subspace system such that  $H = K \oplus K$ ,  $E_1 = K \oplus 0$ ,  $E_2 = 0 \oplus K$ ,  $E_3 = \{(x, Ax); x \in D(A)\}$ , and  $E_4 = \{(x, x); x \in K\}$ . Then  $\mathcal{S}_A$  is transitive if and only if  $A$  is transitive.*

We shall construct transitive operators using transitive Hilbert representations and quotients of operators.

**Definition.** Let  $A$  and  $B$  be bounded linear operators on a Hilbert space  $H$ . We say that  $B(A|_{\text{Ker}(A)^\perp})^{-1}$  is a quotient of  $B$  by  $A$ . We denote  $(A|_{\text{Ker}(A)^\perp})^{-1}$  briefly by  $A^{-1}$ . If we have an additional condition such that  $\ker A \subset \ker B$ , then the quotient is the mapping  $Ax \mapsto Bx, x \in H$ . In [Kau], Kaufman showed the following useful result about quotient operators.

**Theorem 3.5.** [Kau, Theorem 1, page 531] *Let  $T$  be an unbounded operator on a Hilbert space  $H$ . Then  $T$  is a closed operator if and only if  $T = B(A|_{\text{Ker}(A)^\perp})^{-1}$  for some  $A, B \in B(H)$  such that  $\text{Im}(A^*) + \text{Im}(B^*)$  is closed in  $H$ .*

We show that there is a non-zero surjective algebra homomorphism of the endomorphism algebra of a Hilbert representation of the Kronecker quiver to the endomorphism algebra of a four-subspace system. The Kronecker quiver  $Q$  is a quiver with two vertices  $\{1, 2\}$  and two paralleled arrows  $\{\alpha, \beta\}$ :

$$Q : 1 \begin{array}{c} \xrightarrow{\alpha} \\ \xrightarrow{\beta} \end{array} 2$$

A Hilbert representation  $(H, f)$  of the Kronecker quiver is given by two Hilbert spaces  $H_1, H_2$  and two bounded operators  $f_\alpha, f_\beta : H_1 \rightarrow H_2$ .

**Proposition 3.6.** *Let  $K \neq 0$  be a Hilbert space and  $A, B \in B(K)$ . Let  $(H, f)$  be a Hilbert representation of the Kronecker quiver  $Q$  such that  $H_1 = H_2 = K$ ,  $f_\alpha = A$  and  $f_\beta = B$ . Let  $\mathcal{S} = (E_0; E_1, E_2, E_3, E_4)$  be a four-subspace system such that  $E_0 = K \oplus K$ ,  $E_1 = K \oplus 0$ ,  $E_2 = 0 \oplus K$ ,*

$E_3 = \{(Ax, Bx); x \in K\}$ , and  $E_4 = \{(x, x); x \in K\}$ . Assume that  $E_3$  is closed. Then there exists a non-zero surjective algebra homomorphism  $\Phi$  of  $End(H, f)$  to  $End(\mathcal{S})$ . Moreover, if  $\ker A \cap \ker B = 0$ , then  $\Phi$  is one to one.

**Proof.** Let  $(S, T)$  be in  $End(H, f)$ . We have  $AS = TA$  and  $BS = TB$ . Since

$$(T \oplus T)(Ax, Bx) = (TAx, TBx) = (ASx, BSx),$$

we have  $(T \oplus T)(E_3) \subset E_3$ . Clearly  $(T \oplus T)(E_i) \subset E_i$  for  $i = 1, 2, 4$ . Thus we have that  $T \oplus T$  is in  $End(\mathcal{S})$ . We define a mapping  $\Phi$  of  $End(H, f)$  to  $End(\mathcal{S})$  by  $\Phi(S, T) = T \oplus T$ . The map  $\Phi$  is an algebra homomorphism. We shall show that the map  $\Phi$  is onto.

Take  $C \in End(\mathcal{S})$ . Then there exists  $T \in B(K)$  such that  $C = (T \oplus T)$ . We have that

$$(T \oplus T)\{(Ax, Bx); x \in K\} \subset \{(Ay, By); y \in K\}.$$

Hence, for any  $x \in K$ , there exists  $y \in K$  such that  $TAx = Ay$  and  $TBx = By$ . We put  $L_0 = \ker A \cap \ker B$  and  $L_1 = L_0^\perp \cap K$ . By a decomposition of  $y$  such that  $y = y_0 + y_1, y_0 \in L_0, y_1 \in L_1$ , we have  $TAx = Ay_1, TBx = By_1$ . We define an operator  $S$  by  $Sx = y_1$ . We shall show that  $S$  is well defined. If there exists another  $y' = y'_0 + y'_1 \in K$  for  $y'_0 \in L_0$  and  $y'_1 \in L_1$  such that  $TAx = Ay' = Ay'_1$  and  $TBx = By' = By'_1$ . We have  $Ay_1 = Ay'_1$  and  $By_1 = By'_1$ . Hence  $y_1 - y'_1 \in (\ker A \cap \ker B) = L_0$ . We also have  $y_1 - y'_1 \in L_1$ . Hence  $y_1 - y'_1 \in L_0 \cap L_1 = (0)$ . So  $y_1 = y'_1$ . Thus  $S$  is well defined.

Clearly  $S$  is linear. We shall show that  $S$  is a closed operator. Assume that  $x_n \rightarrow x$  and  $Sx_n = y_{n,1} \rightarrow y_1$ , for  $x_n, x \in K$  and  $y_{n,1}, y_1 \in L_1$ . Since  $Sx_n = y_{n,1}$ , we have that  $TAx_n = Ay_{n,1} \rightarrow Ay_1$  and  $TBx_n = By_{n,1} \rightarrow By_1$ . If  $n \rightarrow \infty$ , then  $TAx = Ay_1$  and  $TBx = By_1$ . It follows that  $Sx = y_1$ . Therefore  $S$  is closed. Hence  $S$  is bounded.

Since  $TAx = Ay_1 = ASx$  and  $TBx = By_1 = BSx$  for  $x \in K$  and  $y_1 \in L_1$ , we have that  $TA = AS$  and  $TB = BS$ . Hence  $(S, T) \in End(H, g)$ . And  $\Phi(S, T) = T \oplus T$ . Hence  $\Phi$  is surjective. We shall show that if  $\ker A \cap \ker B = 0$ , then  $\Phi$  is one-to-one. Suppose that  $\Phi(S, T) = T \oplus T = 0$  for  $(S, T) \in End(H, f)$ . Then  $T = 0$ . We have that for any  $x \in K$ ,

$$ASx = TAx = 0, \quad BSx = TBx = 0.$$

Hence  $Sx \in \ker A \cap \ker B = 0$ . Since  $Sx = 0$  for any  $x \in K$ , we have  $S = 0$ . Thus  $(S, T) = 0$ . Therefore  $\Phi$  is one-to-one. □

**Remark.** Let  $K$  be a Hilbert space and  $A, B \in B(K)$ . We consider

$$Z = \begin{pmatrix} A \\ B \end{pmatrix} : K \rightarrow K \oplus K \text{ and } Zx = (Ax, Bx) \text{ for } x \in K.$$

We have

$$Z^* = (A^*, B^*) : K \oplus K \rightarrow K \text{ and } Z^* \begin{pmatrix} x \\ y \end{pmatrix} = A^*x + B^*y \text{ for } x, y \in K.$$

Since  $\text{Im}(Z)$  is closed if and only if  $\text{Im}(Z^*)$  is closed, we have that  $\{(Ax, Bx); x \in K\}$  is closed if and only if  $\text{Im}(A^*) + \text{Im}(B^*)$  is closed.

**Remark.** The map  $\Phi$  is not one-one in general. We shall give an example  $\Phi$  which is not one to one. Let  $K$  be a Hilbert space and  $A, B$  be operators on  $K \oplus K$  such that  $A = B = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ . Let  $S_1, T_1, S_2, T_2$  be operators on  $K \oplus K$  such that  $S_1 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, T_1 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$  and  $S_2 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, T_2 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ . Then  $(S_1, T_1)$  and  $(S_2, T_2)$  are in  $\text{End}(H, f)$ . And  $(S_1, T_1)$  and  $(S_2, T_2)$  give the same endomorphism  $T_1 \oplus T_1$  of  $\mathcal{S}$ . Thus  $\Phi$  is not one to one.

Under a certain condition we have a correspondence between transitive Hilbert representations of the Kronecker quiver and transitive operators.

**Proposition 3.7.** *Let  $K$  be a Hilbert space and  $A, B \in B(K)$ . Assume that  $\ker A = 0$  and  $\text{Im}A^* + \text{Im}B^*$  is closed in  $K$ . Let  $(H, f)$  be a Hilbert representation of the Kronecker quiver  $Q$  such that  $H_1 = H_2 = K, f_\alpha = A$  and  $f_\beta = B$ . Then  $BA^{-1}$  is transitive if and only if  $(H, f)$  is transitive.*

**Proof.** At first we note that the graph  $G(BA^{-1}) = \{(Ax, Bx); x \in K\}$ , because  $\ker(A) = 0$ . Since  $\text{Im}A^* + \text{Im}B^*$  is closed, the operator  $BA^{-1}$  is closed by the remark after Proposition 3.6 (or Theorem 3.5). Let  $\mathcal{S}_{BA^{-1}} = (E_0; E_1, E_2, E_3, E_4)$  be a four-subspace system such that  $E_0 = K \oplus K, E_1 = K \oplus 0, E_2 = 0 \oplus K, E_3 = \{(Ax, Bx); x \in K\} = G(BA^{-1})$ , and  $E_4 = \{(x, x); x \in K\}$ . Since  $\ker(A) = 0$ , there exists an algebra isomorphism  $\Phi$  of  $\text{End}(H, f)$  onto  $\text{End}(\mathcal{S}_{BA^{-1}})$  by Proposition 3.6. Therefore  $(H, f)$  is transitive if and only if  $\mathcal{S}_{BA^{-1}}$  is transitive. Moreover  $\mathcal{S}_{BA^{-1}}$  is transitive if and only if  $BA^{-1}$  is transitive by Lemma 3.4. This implies the conclusion. □

In the following we shall give some examples of transitive operators.

**Proposition 3.8.** *Let  $Q$  be the Kronecker quiver. Let  $S$  be the unilateral shift on  $H = \ell^2(\mathbb{N})$  with a canonical basis  $\{e_1, e_2, \dots\}$ . For a bounded weight vector  $\lambda = (\lambda_1, \lambda_2, \dots) \in \ell^\infty(\mathbb{N})$  we associate with a diagonal operator  $D_\lambda = \text{diag}(\lambda_1, \lambda_2, \dots)$ , so that  $SD_\lambda$  is a weighted shift operator. We assume that  $\lambda_i \neq \lambda_j$  if  $i \neq j$ . Take a vector  $\bar{w} = (\bar{w}_n)_n \in \ell^2(\mathbb{N})$  such that  $w_n \neq 0$  for any  $n \in \mathbb{N}$ . Put  $A = SD_\lambda + \theta_{e_1, \bar{w}}$  and  $B = S$ . Define a Hilbert representation  $(H^\lambda, f^\lambda)$  of the Kronecker quiver  $Q$  by  $H_1^\lambda = H_2^\lambda = H, f_\alpha^\lambda = A$  and  $f_\beta^\lambda = B$ . Then  $\ker A = 0$  and the quotient  $BA^{-1}$  is a transitive operator. Furthermore, the operator  $BA^{-1}$  is densely defined if and only if  $\lambda_k \neq 0$  for each  $k \in \mathbb{N}$  and  $(\frac{w_k}{\lambda_k})_k \notin \ell^2(\mathbb{N})$ .*

**Proof.** By [EW4, Theorem 3.7.], the Hilbert representation  $(H^\lambda, f^\lambda)$  is transitive.

For  $x = (x_n)_n \in \ell^2(\mathbb{N})$ , assume that

$$Ax = (SD_\lambda + \theta_{e_1, \bar{w}})x = \left(\sum_{n=1}^\infty x_n w_n, \lambda_1 x_1, \lambda_2 x_2, \dots\right) = 0.$$

If  $\lambda_k \neq 0$  for any  $k \in \mathbb{N}$ , then  $x_k = 0$  for any  $k \in \mathbb{N}$ . If there exists a  $k \in \mathbb{N}$  such that  $\lambda_k = 0$ , then  $\lambda_i \neq 0$  for  $i \neq k$ . Hence  $x_i = 0$  for  $i \neq k$ . Since  $\sum_{n=1}^\infty x_n w_n = x_k w_k = 0$ ,  $x_k = 0$  by  $w_k \neq 0$ . Thus we have that  $x = 0$  and  $\ker A = 0$ . We note that  $\text{Im} B^* = \text{Im} S^* = H$  and  $\text{Im} A^* + \text{Im} B^* = H$  is closed in  $H$ . Hence  $BA^{-1}$  is a closed operator.

Next we shall consider the condition such that  $BA^{-1}$  is densely defined. We note that  $\overline{D(BA^{-1})} = \overline{\text{Im} A} = (\ker A^*)^\perp$ . We shall show that  $\ker A^* \neq 0$  if and only if (1)  $\lambda_k = 0$  for some  $k \in \mathbb{N}$  or (2)  $\lambda_k \neq 0$  for any  $k \in \mathbb{N}$  and  $(\frac{w_k}{\lambda_k})_k \in \ell^2(\mathbb{N})$ . We see that  $A^* = D_\lambda^* S^* + \theta_{\bar{w}, e_1}$  and  $x = (x_n)_n$  is in  $\ker A^*$  if and only if

$$(\bar{\lambda}_1 x_2, \bar{\lambda}_2 x_3, \dots) = (-x_1 \bar{w}_1, -x_1 \bar{w}_2, \dots).$$

Assume that (1)  $\lambda_k = 0$  for some  $k \in \mathbb{N}$ . We put  $x = (x_i)$  by

$$x_i = \begin{cases} 0 & (i \neq k + 1), \\ 1 & (i = k + 1). \end{cases}$$

We have that  $x \in \ker A^*$  and  $\ker A^* \neq 0$ .

Assume that (2)  $\lambda_k \neq 0$  for any  $k \in \mathbb{N}$  and  $(\frac{w_k}{\lambda_k})_k \in \ell^2(\mathbb{N})$ . Take an element  $x = (1, -(\frac{w_1}{\lambda_1}), -(\frac{w_2}{\lambda_2}), \dots)$ . We have  $x \in \ker A^*$  and  $\ker A^* \neq 0$ . Conversely, assume that there exists  $x (\neq 0) \in \ker A^*$ . Assume that  $x_1 \neq 0$ . Since

$$(\bar{\lambda}_1 x_2, \bar{\lambda}_2 x_3, \dots) = (-x_1 \bar{w}_1, -x_1 \bar{w}_2, \dots),$$

and  $\bar{w}_k \neq 0$  for any  $k \in \mathbb{N}$ , we have  $\lambda_k \neq 0$  for any  $k \in \mathbb{N}$ .

Since  $(-\frac{x_{k+1}}{x_1})_k \in \ell^2(\mathbb{N})$  and  $(-\frac{x_{k+1}}{x_1})_k = (\frac{w_k}{\lambda_k})_k$ , we have that  $(\frac{w_k}{\lambda_k})_k \in \ell^2(\mathbb{N})$ . Hence we have (2). Assume that  $x_1 = 0$ . Since  $x \neq 0$ , there exists  $k \in \mathbb{N}$  such that  $x_{k+1} \neq 0$ . Hence  $\lambda_k = 0$ . Therefore we have (1). □

**Remark.** The operator  $BA^{-1}$  is densely defined for  $\lambda_n = 1/n, w_n = 1/n$  ( $n \in \mathbb{N}$ ). The operator  $BA^{-1}$  is not densely defined for  $(\lambda_n)_n$  defined by

$$\lambda_n = \begin{cases} 0 & (n = 1), \\ 1/n & (n \neq 1). \end{cases}$$

The operator  $BA^{-1}$  is not densely defined for  $\lambda_n = 1 - (1/2^n), w_n = 1/n$  ( $n \in \mathbb{N}$ ).

We refer to [Sh] for weighted shifts.

**Proposition 3.9.** *Let  $Q$  be the Kronecker quiver and  $H = \ell^2(\mathbb{Z})$ . Let  $a = (a(n))_{n \in \mathbb{Z}}, b = (b(n))_{n \in \mathbb{Z}} \in \ell^\infty(\mathbb{Z})$  such that  $a(n) \neq 0, b(n) \neq 0$  for any  $n \in \mathbb{Z}$ . We put  $w_m = \frac{b(m)}{a(m)}, m \in \mathbb{Z}$ . We put*

$$M_k(m, n) := \frac{w_m w_{m+1} \cdots w_{m+k-1}}{w_n w_{n+1} \cdots w_{n+k-1}} \text{ for } m, n \in \mathbb{Z}, k \geq 1.$$

*Assume that for any  $m \neq n, (M_k(m, n))_k$  is an unbounded sequence. Let  $D_a$  be a diagonal operator with  $a = (a(n))_n$  as diagonal coefficients and  $D_b$  be a diagonal operator with  $b = (b(n))_n$  as diagonal coefficients. Let  $U$  be the bilateral forward shift. Put  $A = D_a$  and  $B = UD_b$ . Define a Hilbert representation  $(H, f)$  of the Kronecker quiver  $Q$  by  $H_1 = H_2 = H, f_\alpha = A$  and  $f_\beta = B$ . Then the Hilbert representation  $(H, f)$  is transitive. We also have  $\ker A = 0$  and  $\ker B = 0$ . And the operator  $BA^{-1}$  is a densely defined transitive operator.*

**Proof.** As in [EW4, Theorem 3.8.], we can similarly prove that the Hilbert representation  $(H, f)$  is transitive. By Proposition 3.7, the operator  $BA^{-1}$  is transitive. □

**Example.** [EW4, Theorem 3.8.] Fix a positive constant  $\lambda > 1$ . Consider two sequences  $a = (a(n))_{n \in \mathbb{Z}}$  and  $b = (b(n))_{n \in \mathbb{Z}}$  by

$$a(n) = \begin{cases} e^{-\lambda n} & (n \geq 1, n \text{ is even}), \\ 1 & (\text{otherwise}), \end{cases} \quad b(n) = \begin{cases} e^{-\lambda n} & (n \geq 1, n \text{ is odd}), \\ 1 & (\text{otherwise}). \end{cases}$$

These two sequences  $a$  and  $b$  satisfy the condition of the proposition above.

The concept of transitive operators are useful because certain transitive Hilbert representations of a quiver are given in terms of transitive operators in the next section.

### 4. Extended Dynkin diagrams and transitive Hilbert representations

We consider transitive Hilbert representations of quivers whose underlying undirected graph is an extended Dynkin diagram  $\widetilde{A}_n$  ( $n \geq 0$ ). In the  $\widetilde{A}_0$  case, the oriented cyclic quiver is also called Jordan quiver. Trivially we have no infinite-dimensional transitive Hilbert representations of quivers whose underlying undirected graph is an extended Dynkin diagram  $\widetilde{A}_0$ .

Next we consider transitive Hilbert representations of quivers whose underlying undirected graph is an extended Dynkin diagram  $\widetilde{A}_n$  ( $n \geq 1$ ). The quiver  $C_n$  with  $n \geq 2$  whose underlying undirected graph is an extended Dynkin diagram  $\widetilde{A}_{n-1}$  is called the oriented cyclic quiver if the quiver has cyclic orientation. The set  $V$  of the vertices of  $C_n$  is  $\{1, 2, \dots, n\}$  and the set  $E$  of the arrows of  $C_n$  is  $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$  such that  $s(\alpha_i) = i, r(\alpha_i) = i + 1$  ( $i = 1, \dots, n - 1$ ) and  $s(\alpha_n) = n, r(\alpha_n) = 1$ . For the  $\widetilde{A}_1$  case, the quivers are the oriented cyclic quiver  $C_2$  and the Kronecker quiver  $Q$ .

**Theorem 4.1.** *Let  $\Gamma$  be a quiver whose underlying undirected graph is an extended Dynkin diagram  $\widetilde{A}_n$ ,  $n \geq 1$ . If  $\Gamma$  is not an oriented cyclic quiver, then there exists an infinite-dimensional transitive Hilbert representation of  $\Gamma$ .*

**Proof.** Assume that  $\Gamma$  is not an oriented cyclic quiver. Then there exist vertices  $i$  and  $j$  and arrows  $\alpha$  and  $\beta$  such that  $s(\alpha) = i$ ,  $r(\alpha) = i + 1$  and  $s(\beta) = j + 1$ ,  $r(\beta) = j \pmod n$ . There exists a transitive Hilbert representation  $(H, f)$  of the Kronecker quiver  $Q$  given by  $A, B \in B(H)$  in [EW4, Theorem 3.8.]. We construct a Hilbert representation  $(H', f')$  of  $\Gamma = (V, E)$  such that  $H'_k = H$  ( $k \in V$ ),  $f'_\gamma = I_H$  for  $\gamma \neq \alpha, \beta$  ( $\gamma \in E$ ),  $f'_\alpha = A$ , and  $f'_\beta = B$ . Then the representation  $(H', f')$  of  $\Gamma = (V, E)$  is transitive.  $\square$

By Theorem 4.1, the remaining case of the problem for  $\widetilde{A}_n$  ( $n \geq 1$ ) is an oriented cyclic quiver. It is enough to consider the case that  $H_i \neq 0$  for any  $i$  by the following lemma.

**Lemma 4.2.** *Let  $(H, f)$  be a Hilbert representation of the oriented cyclic quiver  $C_n$ . Assume that there exists a vertex  $k$  such that  $H_k = 0$  ( $1 \leq k \leq n$ ). Let  $(K, g)$  be a Hilbert representation of the oriented cyclic quiver  $C_{n-1}$  such that  $K_i = H_i$  ( $1 \leq i \leq k - 1$ ),  $K_i = H_{i+1}$  ( $k \leq i \leq n - 1$ ),  $g_{\alpha_i} = f_{\alpha_i}$  ( $1 \leq i \leq k - 2$ ),  $g_{\alpha_{k-1}} = 0$ ,  $g_{\alpha_i} = f_{\alpha_{i+1}}$  ( $k \leq i \leq n - 1$ ). Then  $End(H, f)$  is isomorphic to  $End(K, g)$ .*

**Proof.** Take  $T = (T_i)_i \in End(H, f)$  for  $i = 1, \dots, n$ . Since  $H_k = 0$ ,  $B(H_k) = 0$ . Hence we can associate  $T = (T_i)_i \in End(H, f)$  with  $T' = (T'_i)_i \in End(K, g)$ , by putting  $T_i = T'_i$  for  $1 \leq i \leq k - 1$  and  $T_{i+1} = T'_i$  for  $k \leq i \leq n - 1$ . By this correspondence we have that  $End(H, f)$  is isomorphic to  $End(K, g)$ .  $\square$

For the case that  $H_i = \mathbb{C}$  or  $0$ , we introduce a concept of an equivalence relation for vertices in terms of a Hilbert representation.

**Definition.** Let  $(H, f)$  be a Hilbert representation of the oriented cyclic quiver  $C_n = (V, E)$  such that  $H_i = \mathbb{C}$  or  $0$ . We give an equivalence relation for the set of vertices  $\{i \in V; H_i \neq 0\}$  as follows: Take vertices  $i, j$  such that  $H_i \neq 0$  and  $H_j \neq 0$ . We say that vertices  $i$  and  $j$  are  $(H, f)$ -connected if (1)  $i = j$  or (2)  $i < j$  and  $f_{\alpha_{j-1}} \neq 0, \dots, f_{\alpha_{i+1}} \neq 0, f_{\alpha_i} \neq 0$  or (3)  $i > j$  and  $f_{\alpha_{i-1}} \neq 0, \dots, f_{\alpha_{j+1}} \neq 0, f_{\alpha_j} \neq 0$ .

**Lemma 4.3.** *Let  $(H, f)$  be a Hilbert representation of the oriented cyclic quiver  $C_n$  such that  $H_i = \mathbb{C}$  or  $0$  ( $i = 1, 2, \dots, n$ ). Then  $(H, f)$  is transitive if and only if there exists only one  $(H, f)$ -connected component.*

**Proof.** Assume that  $(H, f)$  is transitive. Assume that there exist two  $(H, f)$ -connected components  $D_1$  and  $D_2$  in the set  $\{i \in V; H_i \neq 0\}$ . Let  $\lambda_1 \in \mathbb{C}$ ,  $\lambda_2 \in \mathbb{C}$  such that  $\lambda_1 \neq \lambda_2$ . We define  $T = (T_i)_{i \in V}$  by  $T_i = \lambda_1$

for  $i \in D_1$  and  $T_j = \lambda_2$  for  $j \in D_2$  and  $T_k = 0$  for  $k$  (otherwise). Then  $T = (T_i)_{i \in V}$  is in  $End(H, f)$ . This is a contradiction.

Conversely, assume that there exists only one  $(H, f)$ -connected component. Then there exist decomposition of  $V$  by  $D_3$  and  $D_4$  such that

$$D_3 \cup D_4 = V, D_3 \cap D_4 = \emptyset, D_3 = \{i; H_i \neq 0\}, D_4 = \{j; H_j = 0\}$$

and  $D_3$  is the  $(H, f)$ -connected component.

Let  $T = (T_i)_{i \in V} \in End(H, f)$ . Then  $T_i = T_j$  for  $i, j \in D_3$ . In fact, if  $i < j$ , then  $f_{\alpha_i} \neq 0, f_{\alpha_{i+1}} \neq 0, \dots, f_{\alpha_{j-1}} \neq 0$  and  $f_{\alpha_i} T_i = T_{i+1} f_{\alpha_i}, \dots, f_{\alpha_{j-1}} T_{j-1} = T_j f_{\alpha_{j-1}}$ . Since  $f_{\alpha_i} \neq 0$  for  $i \in D_3$ ,  $T_i = T_{i+1} = \dots = T_j$ . Hence  $T_i = T_j$  for all  $i, j \in D_3$ . And  $T_i = T_j = 0$  for  $i, j \in D_4$ . Thus  $End(H, f)$  is isomorphic to  $\mathbb{C}$ . Hence  $(H, f)$  is transitive.  $\square$

The next lemma guarantees that we may assume that  $H_i \subset H_j$  if  $\dim H_i \leq \dim H_j$ .

**Lemma 4.4.** *Let  $(H_i)_{i=1}^n$  be a family of nonzero Hilbert spaces. Then there exists a family  $(K(i))_{i=1}^n$  of subspaces in a Hilbert space  $V$ , such that for any  $i$  ( $1 \leq i \leq n$ ), there exists a number  $m(i)$  ( $1 \leq m(i) \leq n$ ) such that  $H_i$  is isomorphic to  $\bigoplus_{j=1}^{m(i)} K(j)$ .*

**Proof.** We arrange a family of Hilbert spaces  $(H_i)_i$  in increasing order of dimension and as a result, we have  $(H_{\ell(1)}), (H_{\ell(2)}), \dots, (H_{\ell(n)})$  in increasing order of dimension. Construct an ambient space  $V$  and its increasing subspaces  $H'_i \cong H_i$  such that  $(H'_{\ell(1)}) \subset (H'_{\ell(2)}) \subset \dots \subset (H'_{\ell(n)}) \subset V$ . Put  $K_1 = H'_{\ell(1)}, K_2 = H'_{\ell(2)} \cap (H'_{\ell(1)})^\perp, \dots, K_n = H'_{\ell(n)} \cap (H'_{\ell(n-1)})^\perp$ . Hence there exists a number  $m(i)$  such that  $H'_i = K(1) \oplus K(2) \oplus \dots \oplus K(m(i))$ . Thus we have that  $H_i$  is isomorphic to  $K(1) \oplus K(2) \oplus \dots \oplus K(m(i))$ .  $\square$

Firstly we investigate transitive Hilbert representations of oriented cyclic quivers  $C_2$  and  $C_3$ . Let  $(H, f)$  be a Hilbert representation of  $C_2$ . In what follows we denote  $f_{\alpha_1}, f_{\alpha_2}$  by  $A_1, A_2$  for short.

**Lemma 4.5.** *Let  $(H, f)$  be a transitive Hilbert representation of  $C_2$ . Assume that  $H_1 = H_2 = K \neq 0, A_1 \in \mathbb{C}$  and  $A_2 \in \mathbb{C}$ . If  $A_1 \neq 0$  or  $A_2 \neq 0$ , then  $K = \mathbb{C}$ .*

**Proof.** Let  $T \in B(K)$ . Then  $(T, T) \in End(H, f)$ . In fact,  $A_1 T = T A_1$  and  $A_2 T = T A_2$ . If  $\dim K > 1, B(K) \neq \mathbb{C}I$ . Since  $(H, f)$  is transitive, this is a contradiction. Thus  $\dim K = 1$ .  $\square$

**Lemma 4.6.** *Let  $(H, f)$  be a Hilbert representation of  $C_2$ . Then  $(H, f)$  is transitive if and only if one of the following conditions holds.*

- (1)  $H_1 = \mathbb{C}, H_2 = 0, A_1 = 0$  and  $A_2 = 0,$
- (2)  $H_1 = 0, H_2 = \mathbb{C}, A_1 = 0$  and  $A_2 = 0,$
- (3)  $H_1 = \mathbb{C}$  and  $H_2 = \mathbb{C}$  and  $(A_1 \neq 0$  or  $A_2 \neq 0).$



**Proof.** If (1), (2) or (3) holds, then  $(H, f)$  is clearly transitive. Conversely, assume that  $(H, f)$  is transitive. Assume that  $\dim H_1 \neq 0$  and  $\dim H_2 = 0$ . If  $\dim H_1 > 1$ , then there exists a non-scalar operator in  $B(H_1)$ . Since  $B(H_1) = \text{End}(H, f)$ , this contradicts the transitivity of  $(H, f)$ . Hence  $\dim H_1 = 1$ . This is the case (1).

Similarly we have the case (2).

Therefore it is sufficient to assume that  $\dim H_1 \neq 0$  and  $\dim H_2 \neq 0$ . By Lemma 4.4 we may assume that  $\dim H_1 \leq \dim H_2$  and  $H_1$  is a subspace of  $H_2$ . We define  $T = (T_1, T_2) = (A_2A_1, A_1A_2)$ . Then  $T \in \text{End}(H, f)$ . In fact,

$$A_1T_1 = A_1(A_2A_1) = (A_1A_2)A_1 = T_2A_1$$

and

$$T_1A_2 = (A_2A_1)A_2 = A_2(A_1A_2) = A_2T_2.$$

By the assumption of transitivity for  $(H, f)$ ,

$$(T_1, T_2) \in \{(\mu I_{H_1}, \mu I_{H_2}) \mid \mu \in \mathbb{C}\}.$$

Hence

$$T_1 = A_2A_1 = \mu I_{H_1}, T_2 = A_1A_2 = \mu I_{H_2} \text{ for some } \mu \in \mathbb{C}.$$

We denote by  $E_1 \in B(H_1, H_2)$  the embedding map of  $H_1$  into  $H_2$  and  $E_2 \in B(H_2, H_1)$  the projection map of  $H_2$  onto  $H_1$ . We define

$$T^{\{1\}} = (T_1^{\{1\}}, T_2^{\{1\}}) = (A_2E_1, E_1A_2).$$

Then  $T^{\{1\}} \in \text{End}(H, f)$ . In fact,

$$\begin{aligned} A_1T_1^{\{1\}} &= A_1(A_2E_1) = (A_1A_2)E_1 = \mu I_{H_2}E_1 \\ &= \mu E_1 = E_1\mu I_{H_1} = (E_1A_2)A_1 = T_2^{\{1\}}A_1 \end{aligned}$$

and

$$T_1^{\{1\}}A_2 = (A_2E_1)A_2 = A_2(E_1A_2) = A_2T_2^{\{1\}}.$$

Thus  $T^{\{1\}} \in \text{End}(H, f)$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{1\}} \in \mathbb{C}$  such that  $A_2E_1 = \mu^{\{1\}}I_{H_1}$  and  $E_1A_2 = \mu^{\{1\}}I_{H_2}$ . We define

$$T^{\{2\}} = (T_1^{\{2\}}, T_2^{\{2\}}) = (E_2A_1, A_1E_2).$$

Then  $T^{\{2\}} \in \text{End}(H, f)$ . In fact,

$$A_1T_1^{\{2\}} = A_1(E_2A_1) = (A_1E_2)A_1 = T_2^{\{2\}}A_1$$

and

$$\begin{aligned} T_1^{\{2\}}A_2 &= (E_2A_1)A_2 = E_2(\mu I_{H_2}) = \mu E_2 \\ &= \mu I_{H_1}E_2 = A_2(A_1E_2) = A_2T_2^{\{2\}}. \end{aligned}$$

Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{2\}} \in \mathbb{C}$  such that  $E_2A_1 = \mu^{\{2\}}I_{H_1}$  and  $A_1E_2 = \mu^{\{2\}}I_{H_2}$ .

We define

$$T^{\{1,2\}} = (T_1^{\{1,2\}}, T_2^{\{1,2\}}) = (E_2E_1, E_1E_2).$$

Then  $T^{\{1,2\}} \in \text{End}(H, f)$ . In fact,

$$\begin{aligned} A_1 T_1^{\{1,2\}} &= A_1(E_2 E_1) = (A_1 E_2) E_1 = \mu^{\{2\}} I_{H_2} E_1 = \mu^{\{2\}} E_1 \\ &= \mu^{\{2\}} E_1 = E_1(\mu^{\{2\}} I_{H_1}) = E_1(E_2 A_1) = T_2^{\{1,2\}} A_1, \end{aligned}$$

and

$$\begin{aligned} T_1^{\{1,2\}} A_2 &= (E_2 E_1) A_2 = E_2(\mu^{\{1\}} I_{H_2}) = \mu^{\{1\}} E_2 \\ &= \mu^{\{1\}} E_2 = (\mu^{\{1\}} I_{H_1}) E_2 = A_2(E_1 E_2) = A_2 T_2^{\{1,2\}}. \end{aligned}$$

Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{1,2\}} \in \mathbb{C}$  such that

$$E_2 E_1 = \mu^{\{1,2\}} I_{H_1}, \quad E_1 E_2 = \mu^{\{1,2\}} I_{H_2}.$$

For  $x(\neq 0) \in H_1$ , we have  $x = E_2 E_1 x = \mu^{\{1,2\}} I_{H_1} x = \mu^{\{1,2\}} x$ . Hence  $\mu^{\{1,2\}} = 1$ . If  $H_1 \neq H_2$ , then  $H_1^\perp \cap H_2 \neq 0$ . Take  $x(\neq 0) \in H_1^\perp \cap H_2$ . Then  $E_1 E_2 x = \mu^{\{1,2\}} I_{H_2} x$ . Hence  $0 = x$ . This is a contradiction. Thus  $H_1 = H_2$  and  $E_1 = E_2$ . Since  $A_1 E_2 = \mu^{\{2\}} I_{H_2}$ ,  $A_1 = \mu^{\{2\}} I_{H_1}$ . And we also have  $E_1 A_2 = A_2 = \mu^{\{1\}} I_{H_1}$ . Since  $(H, f)$  is transitive,  $A_1 \neq 0$  or  $A_2 \neq 0$ . By Lemma 4.5, we have  $H_1 = H_2 = \mathbb{C}$ . Thus  $(H, f)$  is in the case (3).  $\square$

Let  $(H, f)$  be a Hilbert representation of the oriented cyclic quiver  $C_3$ . In the below we denote  $f_{\alpha_1}, f_{\alpha_2}, f_{\alpha_3}$  by  $A_1, A_2, A_3$  for short.

**Lemma 4.7.** *Let  $(H, f)$  be a transitive Hilbert representation of  $C_3$ . Assume that  $H_i = \mathbb{C}$  ( $i = 1, 2, 3$ ). Then  $A_i A_j \neq 0$  for some  $i \neq j$ .*

**Proof.** Assume that  $A_i = A_j = 0$  for some  $i \neq j$ . We may and do assume  $i = 1, j = 2$ . Let  $T = (T_1, T_2, T_3)$  such that  $T_1 = T_3, T_2 \neq T_1, T_1 \neq 0$  and  $T_2 \neq 0$ . Then  $T = (T_1, T_2, T_3)$  is in  $\text{End}(H, f)$ . Since  $(H, f)$  is transitive,  $T_1 = T_2 = T_3 \in \mathbb{C}$ . This is a contradiction. Hence this lemma holds.  $\square$

**Lemma 4.8.** *Let  $(H, f)$  be a Hilbert representation of  $C_3$ . Then  $(H, f)$  is transitive if and only if one of the following holds.*

- (1)  $H_1 = \mathbb{C}$  and  $H_i = 0$  ( $i = 2, 3$ ).
- (2)  $H_2 = \mathbb{C}$  and  $H_i = 0$  ( $i = 1, 3$ ).
- (3)  $H_3 = \mathbb{C}$  and  $H_i = 0$  ( $i = 1, 2$ ).
- (4)  $H_i = \mathbb{C}$  ( $i = 1, 2$ ),  $H_3 = 0$  and  $A_1 \neq 0$ .
- (5)  $H_i = \mathbb{C}$  ( $i = 2, 3$ ),  $H_1 = 0$  and  $A_2 \neq 0$ .
- (6)  $H_i = \mathbb{C}$  ( $i = 1, 3$ ),  $H_2 = 0$  and  $A_3 \neq 0$ .
- (7)  $H_i = \mathbb{C}$  ( $i = 1, 2, 3$ ) and  $A_i A_j \neq 0$  for some  $i \neq j$  ( $i, j = 1, 2, 3$ ).

**Proof.** If a Hilbert representations  $(H, f)$  satisfies (1), (2),  $\dots$  or (7), then the Hilbert representation is obviously transitive. Conversely assume that  $(H, f)$  is transitive. At first we assume that all Hilbert spaces  $H_i \neq 0$  ( $1 \leq i \leq 3$ ) and by Lemma 4.4 a totally ordered set by inclusion order and  $H_1 \subset H_i$  ( $i = 2, 3$ ). We define

$$T_1 = A_3 A_2 A_1, \quad T_2 = A_1 A_3 A_2, \quad T_3 = A_2 A_1 A_3, \quad T = (T_1, T_2, T_3).$$

We define a mapping  $E_i \in B(H_i, H_{i+1})$  by

$$E_i = \begin{cases} \text{the inclusion map of } H_i \text{ into } H_{i+1}, & H_i \subset H_{i+1}, \\ \text{the projection map of } H_i \text{ onto } H_{i+1}, & H_{i+1} \subset H_i. \end{cases}$$

For a subset  $S$  of  $\{1, 2, 3\}$ , we define  $B_i \in B(H_i, H_{i+1})$  by

$$B_i = \begin{cases} A_i & \text{if } i \notin S, \\ E_i & \text{if } i \in S. \end{cases}$$

We also define

$$T_1^S = B_3 B_2 B_1, \quad T_2^S = B_1 B_3 B_2, \quad T_3^S = B_2 B_1 B_3, \quad T^S = (T_1^S, T_2^S, T_3^S).$$

We note that  $T^S = (T_1^S, T_2^S, T_3^S)$  is obtained by replacing each word  $A_i$  in  $T = (T_1, T_2, T_3)$  with  $E_i$  for all  $i \in S$ . We regard  $T = (T_1, T_2, T_3)$  as  $T^\emptyset = (T_1^\emptyset, T_2^\emptyset, T_3^\emptyset)$ . Since

$$A_1 T_1 = A_1 (A_3 A_2 A_1) = T_2 A_1,$$

$$A_2 T_2 = A_2 (A_1 A_3 A_2) = T_3 A_2,$$

$$A_3 T_3 = A_3 (A_2 A_1 A_3) = T_1 A_3,$$

we have that  $T$  is in  $End(H, f)$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu \in \mathbb{C}$  such that

$$A_3 A_2 A_1 = \mu I_{H_1}, \quad A_1 A_3 A_2 = \mu I_{H_2}, \quad A_2 A_1 A_3 = \mu I_{H_3}.$$

For  $S = \{1\}$ , we define  $T^S = T^{\{1\}} = (T_1^{\{1\}}, T_2^{\{1\}}, T_3^{\{1\}})$  by

$$T_1^{\{1\}} = A_3 A_2 E_1, \quad T_2^{\{1\}} = E_1 A_3 A_2, \quad T_3^{\{1\}} = A_2 E_1 A_3.$$

It follows that

$$\begin{aligned} A_1 T_1^{\{1\}} &= A_1 A_3 A_2 E_1 = \mu I_{H_2} E_1 = \mu E_1 \\ &= E_1 \mu I_{H_1} = E_1 (A_3 A_2) A_1 = T_2^{\{1\}} A_1, \end{aligned}$$

$$A_2 T_2^{\{1\}} = A_2 E_1 (A_3 A_2) = T_3^{\{1\}} A_2,$$

$$A_3 T_3^{\{1\}} = A_3 A_2 E_1 A_3 = T_1^{\{1\}} A_3.$$

Thus  $T^{\{1\}}$  is in  $End(H, f)$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{1\}} \in \mathbb{C}$  such that

$$A_3 A_2 E_1 = \mu^{\{1\}} I_{H_1}, \quad E_1 A_3 A_2 = \mu^{\{1\}} I_{H_2}, \quad A_2 E_1 A_3 = \mu^{\{1\}} I_{H_3}.$$

For  $S = \{2\}$ , we define  $T^S = T^{\{2\}} = (T_1^{\{2\}}, T_2^{\{2\}}, T_3^{\{2\}})$  by

$$T_1^{\{2\}} = A_3 E_2 A_1, \quad T_2^{\{2\}} = A_1 A_3 E_2, \quad T_3^{\{2\}} = E_2 A_1 A_3.$$

It follows that

$$\begin{aligned} A_1 T_1^{\{2\}} &= A_1 A_3 E_2 A_1 = T_2^{\{2\}} A_1, \\ A_2 T_2^{\{2\}} &= A_2 A_1 A_3 E_2 = \mu I_{H_3} E_2 = \mu E_2 \\ &= E_2 \mu I_{H_2} = E_2 A_1 A_3 A_2 = T_3^{\{2\}} A_2, \\ A_3 T_3^{\{2\}} &= A_3 E_2 A_1 A_3 = T_1^{\{2\}} A_3. \end{aligned}$$

Thus  $T^{\{2\}}$  is in  $End(H, f)$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{2\}} \in \mathbb{C}$  such that

$$A_3 E_2 A_1 = \mu^{\{2\}} I_{H_1}, \quad A_1 A_3 E_2 = \mu^{\{2\}} I_{H_2}, \quad (E_2 A_1) A_3 = \mu^{\{2\}} I_{H_3}.$$

For  $S = \{3\}$ , we define  $T^S = T^{\{3\}} = (T_1^{\{3\}}, T_2^{\{3\}}, T_3^{\{3\}})$  by

$$T_1^{\{3\}} = E_3 A_2 A_1, \quad T_2^{\{3\}} = A_1 E_3 A_2, \quad T_3^{\{3\}} = A_2 A_1 E_3.$$

It follows that

$$\begin{aligned} A_1 T_1^{\{3\}} &= A_1 E_3 A_2 A_1 = T_2^{\{3\}} A_1, \\ A_2 T_2^{\{3\}} &= A_2 A_1 E_3 A_2 = T_3^{\{3\}} A_2, \quad A_3 T_3^{\{3\}} = A_3 A_2 A_1 E_3 = \mu I_{H_1} E_3 = \mu E_3 \\ &= E_3 \mu I_{H_3} = E_3 A_2 A_1 A_3 = T_1^{\{3\}} A_3. \end{aligned}$$

Thus  $T^{\{3\}}$  is in  $End(H, f)$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{3\}} \in \mathbb{C}$  such that

$$E_3 A_2 A_1 = \mu^{\{3\}} I_{H_1}, \quad A_1 E_3 A_2 = \mu^{\{3\}} I_{H_2}, \quad A_2 A_1 E_3 = \mu^{\{3\}} I_{H_3}.$$

For  $S = \{1, 2\}$ , we have

$$T^{\{1,2\}} = (T_1^{\{1,2\}}, T_2^{\{1,2\}}, T_3^{\{1,2\}}) = (A_3 E_2 E_1, E_1 A_3 E_2, E_2 E_1 A_3).$$

It follows that

$$\begin{aligned} A_1 T_1^{\{1,2\}} &= A_1 A_3 E_2 E_1 = \mu^{\{2\}} I_{H_2} E_1 = \mu^{\{2\}} E_1 \\ &= E_1 \mu^{\{2\}} I_{H_1} = E_1 A_3 E_2 A_1 = T_2^{\{1,2\}} A_1, \\ A_2 T_2^{\{1,2\}} &= A_2 E_1 A_3 E_2 = \mu^{\{1\}} I_{H_3} E_2 = \mu^{\{1\}} E_2 \\ &= E_2 \mu^{\{1\}} I_{H_2} = E_2 E_1 A_3 A_2 = T_3^{\{1,2\}} A_2, \\ A_3 T_3^{\{1,2\}} &= A_3 E_2 E_1 A_3 = T_1^{\{1,2\}} A_3. \end{aligned}$$

Thus  $T^{\{1,2\}}$  is in  $End(H, f)$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{1,2\}} \in \mathbb{C}$  such that

$$A_3 E_2 E_1 = \mu^{\{1,2\}} I_{H_1}, \quad E_1 A_3 E_2 = \mu^{\{1,2\}} I_{H_2}, \quad E_2 E_1 A_3 = \mu^{\{1,2\}} I_{H_3}.$$

For  $S = \{1, 3\}$ , we have

$$T^{\{1,3\}} = (T_1^{\{1,3\}}, T_2^{\{1,3\}}, T_3^{\{1,3\}}) = (E_3 A_2 E_1, E_1 E_3 A_2, A_2 E_1 E_3).$$

It follows that

$$\begin{aligned} A_1 T_1^{\{1,3\}} &= A_1 E_3 A_2 E_1 = \mu^{\{3\}} I_{H_2} E_1 = \mu^{\{3\}} E_1 \\ &= E_1 \mu^{\{3\}} I_{H_1} = E_1 E_3 A_2 A_1 = T_2^{\{1,3\}} A_1, \\ A_2 T_2^{\{1,3\}} &= A_2 E_1 E_3 A_2 = T_3^{\{1,3\}} A_2, \\ A_3 T_3^{\{1,3\}} &= A_3 A_2 E_1 E_3 = \mu^{\{1\}} I_{H_1} E_3 = \mu^{\{1\}} E_3 \\ &= E_3 \mu^{\{1\}} I_{H_3} = E_3 A_2 E_1 A_3 = T_1^{\{1,3\}} A_3. \end{aligned}$$

Thus  $T^{\{1,3\}}$  is in  $End(H, f)$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{1,3\}} \in \mathbb{C}$  such that

$$E_3 A_2 E_1 = \mu^{\{1,3\}} I_{H_1}, \quad E_1 E_3 A_2 = \mu^{\{1,3\}} I_{H_2}, \quad A_2 E_1 E_3 = \mu^{\{1,3\}} I_{H_3}.$$

For  $S = \{2, 3\}$ , we have

$$T^{\{2,3\}} = (T_1^{\{2,3\}}, T_2^{\{2,3\}}, T_3^{\{2,3\}}) = (E_3 E_2 A_1, A_1 E_3 E_2, E_2 A_1 E_3).$$

It follows that

$$\begin{aligned} A_1 T_1^{\{2,3\}} &= A_1 E_3 E_2 A_1 = T_2^{\{2,3\}} A_1, \\ A_2 T_2^{\{2,3\}} &= A_2 A_1 E_3 E_2 = \mu^{\{3\}} I_{H_3} E_2 = \mu^{\{3\}} E_2 \\ &= E_2 \mu^{\{3\}} I_{H_2} = E_2 A_1 E_3 A_2 = T_3^{\{2,3\}} A_2, \\ A_3 T_3^{\{2,3\}} &= A_3 E_2 A_1 E_3 = \mu^{\{2\}} I_{H_1} E_3 = \mu^{\{2\}} E_3 \\ &= E_3 \mu^{\{2\}} I_{H_3} = E_3 E_2 A_1 A_3 = T_1^{\{2,3\}} A_3. \end{aligned}$$

Thus  $T^{\{2,3\}}$  is in  $End(H, f)$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{2,3\}} \in \mathbb{C}$  such that

$$E_3 E_2 A_1 = \mu^{\{2,3\}} I_{H_1}, \quad A_1 E_3 E_2 = \mu^{\{2,3\}} I_{H_2}, \quad E_2 A_1 E_3 = \mu^{\{2,3\}} I_{H_3}.$$

For  $S = \{1, 2, 3\}$ , we have

$$T^{\{1,2,3\}} = (T_1^{\{1,2,3\}}, T_2^{\{1,2,3\}}, T_3^{\{1,2,3\}}) = (E_3 E_2 E_1, E_1 E_3 E_2, E_2 E_1 E_3).$$

It follows that

$$\begin{aligned} A_1 T_1^{\{1,2,3\}} &= A_1 E_3 E_2 E_1 = \mu^{\{2,3\}} I_{H_1} E_1 = \mu^{\{2,3\}} E_1 \\ &= E_1 \mu^{\{2,3\}} I_{H_1} = E_1 E_3 E_2 A_1 = T_2^{\{1,2,3\}} A_1, \\ A_2 T_2^{\{1,2,3\}} &= A_2 E_1 E_3 E_2 = \mu^{\{1,3\}} I_{H_3} E_2 = \mu^{\{1,3\}} E_2 \\ &= E_2 \mu^{\{1,3\}} I_{H_2} = E_2 E_1 E_3 A_2 = T_3^{\{1,2,3\}} A_2, \\ A_3 T_3^{\{1,2,3\}} &= A_3 E_2 E_1 E_3 = \mu^{\{1,2\}} I_{H_1} E_3 = \mu^{\{1,2\}} E_3 \\ &= E_3 \mu^{\{1,2\}} I_{H_3} = E_3 E_2 E_1 A_3 = T_1^{\{1,2,3\}} A_3. \end{aligned}$$

Thus  $T^{\{1,2,3\}}$  is in  $End(H, f)$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu^{\{1,2,3\}} \in \mathbb{C}$  such that

$$E_3E_2E_1 = \mu^{\{1,2,3\}}I_{H_1}, \quad E_1E_3E_2 = \mu^{\{1,2,3\}}I_{H_2}, \quad E_2E_1E_3 = \mu^{\{1,2,3\}}I_{H_3}.$$

Take  $x (\neq 0) \in H_1$ . Since  $H_1 \subset H_i$  ( $1 \leq i \leq 3$ ),  $E_3E_2E_1x = x = \mu^{\{1,2,3\}}I_{H_1}x$ . Hence  $\mu^{\{1,2,3\}} = 1$ . By Lemma 4.4, we can represent  $H_i$  by  $H_i = K_1 \oplus K_2 \oplus \cdots \oplus K_{m(i)}$  ( $1 \leq i \leq 3$ ). We shall show that  $H_1 = H_2 = H_3$ .

Now,  $m(1) = 1$  and assume that  $m(2) \neq 1$ . We compare  $m(3)$  with  $m(2)$ . Assume that  $m(3) < m(2)$ . Take  $x (\neq 0) \in K_{m(2)} \subset H_2$ . Then  $E_2x = 0$ . This contradicts that  $E_1E_3E_2 = I_{H_2}$ . Assume that  $m(3) \geq m(2)$ . Take  $x (\neq 0) \in K_{m(2)} \subset H_2$ . Then  $E_2x = x$  and  $E_3x = 0$ . This contradicts that  $E_1E_3E_2 = I_{H_2}$ .

Hence  $m(1) = m(2)$  and  $H_1 = H_2$ . Next assume that  $H_3 \neq H_1$  (hence  $m(3) \neq 1$ ). Take  $x (\neq 0) \in K_{m(3)} \subset H_3$ . Then  $E_3x = 0$ . This contradicts that  $E_2E_1E_3 = I_{H_3}$ . Hence we have that  $H_1 = H_2 = H_3 := M$ . Therefore,  $E_1 = E_2 = E_3 = I_M$  and  $T_i^{\{1,2,3\}} = I_M \in \mathbb{C}$  Since

$$E_3E_2A_1 = \mu^{\{2,3\}}I_{H_1}, \quad E_1E_3A_2 = \mu^{\{1,3\}}I_{H_2}, \quad E_2E_1A_3 = \mu^{\{1,2\}}I_{H_3},$$

we have

$$A_1 = \mu^{\{2,3\}}I_M, \quad A_2 = \mu^{\{1,3\}}I_M, \quad A_3 = \mu^{\{1,2\}}I_M.$$

If  $\dim M > 1$ , there is a non-scalar operator  $B \in B(M)$ . Since  $A_1, A_2, A_3$  are scalar operators,  $(B, B, B) \in End(H, f)$ . This contradicts that  $(H, f)$  is transitive. Hence we have  $\dim M = 1$ . By Lemma 4.7,  $A_iA_j \neq 0$  for some  $i \neq j$  ( $i, j = 1, 2, 3$ ). Thus  $(H, f)$  is in the case (7).

Next we consider other cases. Assume that there exists  $H_i = 0$  for some  $i$ . Since  $(H, f)$  is transitive, the number  $|\{i; H_i \neq 0\}|$  is 1 or 2. If  $|\{i; H_i \neq 0\}| = 1 = |\{k\}|$ , then  $\dim H_k = 1$  because  $(H, f)$  is transitive. Hence these are in the cases (1), (2), (3). If  $|\{i; H_i \neq 0\}| = 2 = |\{k, \ell\}|$ , ( $k < \ell \pmod 3$ ), then we consider the reduction  $C_2$  of the quiver  $C_3$  as it is shown in Lemma 4.2. Let  $(K, g)$  be the reduced Hilbert representation of  $C_2$  from the Hilbert representation  $(H, f)$  of  $C_2$  by Lemma 4.2. We have  $End(H, f) \cong End(K, g)$ . Hence  $End(K, g)$  is transitive. By the same argument in the case (7), we have  $\dim H_k = \dim H_\ell = 1$ . Since  $(H, f)$  is transitive,  $A_k \neq 0$ . Thus these are in the cases (4), (5), (6). All these cases are summarized as the existence of unique  $(H, f)$ -connected component by Lemma 4.3.  $\square$

Let  $(H, f)$  be a Hilbert representation of  $C_n$ . In the below we denote  $f_{\alpha_1}, f_{\alpha_2}, \dots, f_{\alpha_n}$  by  $A_1, A_2, \dots, A_n$  for short.

**Lemma 4.9.** *Let  $(H, f)$  be a Hilbert representation of the oriented cyclic quiver  $C_n$ . Then  $(H, f)$  is transitive if and only if  $H_i = \mathbb{C}$  or 0 and there exists only one  $(H, f)$ -connected component in  $\{i \in V; H_i \neq 0\}$ .*

**Proof.** Assume that  $H_i = \mathbb{C}$  or 0 and there exists only one  $(H, f)$ -connected component in  $\{i \in V; H_i \neq 0\}$ . Then  $(H, f)$  is transitive by Lemma 4.3.

Conversely assume that  $(H, f)$  is transitive. At first we consider the case that  $H_i \neq 0$  for any  $i$ . By lemma 4.4, we may and do assume that the family  $(H_i)$  of Hilbert spaces are totally ordered under the inclusion order. We also assume that  $\dim H_1$  is the smallest dimension among  $\{\dim H_i; i = 1, \dots, n\}$ .

We define  $T = (T_1, T_2, \dots, T_n)$  by

$$T_1 = A_n \cdots A_3 A_2 A_1, T_2 = A_1 A_n \cdots A_4 A_3 A_2, \dots, T_n = A_{n-1} \cdots A_3 A_2 A_1 A_n.$$

Then  $T = (T_1, T_2, \dots, T_n)$  is clearly in  $End(H, f)$ .

We denote by  $E_i$  the following operator  $E_i : H_i \rightarrow H_{i+1}$ :

$$E_i = \begin{cases} \text{the inclusion map from } H_i \text{ into } H_{i+1}, & H_i \subset H_{i+1}, \\ \text{the projection map from } H_i \text{ onto } H_{i+1}, & H_{i+1} \subset H_i. \end{cases}$$

For  $S \subseteq \{1, 2, \dots, n\}$ , we define  $B_i \in B(H_i, H_{i+1})$ , which depends on  $S$ , by

$$B_i = \begin{cases} A_i, & \text{if } i \notin S \\ E_i, & \text{if } i \in S \end{cases}$$

We also define  $T_i^S \in B(H_i)$  and  $T^S \in B(H_1 \oplus \dots \oplus H_n)$  by

$$T_i^S = B_{i-1} B_{i-2} \cdots B_2 B_1 B_n B_{n-1} \cdots B_{i+1} B_i \quad \text{for } 1 \leq i \leq n,$$

and  $T^S = (T_1^S, T_2^S, \dots, T_n^S)$ . That is,  $T^S = (T_1^S, T_2^S, \dots, T_n^S)$  is obtained by replacing each word  $A_i$  in  $T = (T_1, T_2, \dots, T_n)$  with  $E_i$  for all  $i \in S$ .

For example,  $T^{\{1\}} = (T_1^{\{1\}}, T_2^{\{1\}}, \dots, T_n^{\{1\}})$  is given by

$$\begin{aligned} T_1^{\{1\}} &= A_n A_{n-1} \cdots A_2 E_1, \\ T_2^{\{1\}} &= E_1 A_n A_{n-1} \cdots A_3 A_2, \\ T_3^{\{1\}} &= A_2 E_1 A_n A_{n-1} \cdots A_4 A_3, \\ &\dots \\ T_n^{\{1\}} &= A_{n-1} A_{n-2} \cdots A_2 E_1 A_n. \end{aligned}$$

We regard  $T$  as  $T^\emptyset$ .

In the following we shall show that  $T^S = (T_1^S, T_2^S, \dots, T_n^S)$  is in  $End(H, f)$  for any  $S \subseteq \{1, 2, \dots, n\}$ . We shall prove it by the induction on the number  $k = |S|$ . First consider the case  $k = |S| = 0$ , that is,  $S = \emptyset$ . Then  $T^\emptyset = T = (T_1, T_2, \dots, T_n)$  is clearly in  $End(H, f)$ .

Next, we assume that  $T^S$  is in  $End(H, f)$  for  $|S| = k$ . Since  $(H, f)$  is transitive, there exists a constant  $\mu^S \in \mathbb{C}$  such that  $T_i^S = \mu^S I_{H_i}$  for any  $i = 1, \dots, n$ . Take  $S$  such that  $|S| = k + 1$ . We shall show that  $T^S$  is in  $End(H, f)$ . It is enough to show that, for any  $i = 1, \dots, n$ , we have  $A_i T_i^S = T_{i+1}^S A_i$ . First we consider the case that  $i = 1$ . We need to show the validity of the relation  $A_1 T_1^S = T_2^S A_1$ , that is,  $A_1 B_n \cdots B_2 B_1 = B_1 B_n \cdots B_2 A_1$ . Assume that 1 is in  $S$ . Then  $B_1 = E_1$  and  $T_i^{S \setminus \{1\}}$  is in  $End(H, f)$  by the

assumption of the induction. Since  $A_1B_nB_{n-1}\cdots B_2$  and  $B_nB_{n-1}\cdots B_2A_1$  have  $k$  changed letters, we have

$$T_2^{S\setminus\{1\}} = A_1B_nB_{n-1}\cdots B_2 = \mu^{S\setminus\{1\}}I_{H_2}$$

and

$$T_1^{S\setminus\{1\}} = B_nB_{n-1}\cdots B_2A_1 = \mu^{S\setminus\{1\}}I_{H_1}.$$

Therefore, we have

$$A_1T_1^S = A_1B_n\cdots B_1 = \mu^{S\setminus\{1\}}I_{H_2}B_1 = \mu^{S\setminus\{1\}}E_1$$

and

$$T_2^S A_1 = B_1B_nB_{n-1}\cdots B_2A_1 = B_1\mu^{S\setminus\{1\}}I_{H_1} = \mu^{S\setminus\{1\}}E_1.$$

Thus  $A_1T_1^S = T_2^S A_1$ .

Assume that 1 is not in  $S$ . Then  $B_1 = A_1$ . Hence

$$A_1T_1^S = A_1B_n\cdots B_1 = A_1B_n\cdots B_2A_1$$

and

$$T_2^S A_1 = B_1B_n\cdots B_2A_1 = A_1B_n\cdots B_2A_1.$$

Thus  $A_1T_1^S = T_2^S A_1$ .

For other cases that  $i = 2, 3, \dots, n$ , we also have that  $A_iT_i^S = T_{i+1}^S A_i$ . Hence, by induction, we have that  $T^S$  is in  $End(H, f)$  for any  $S \subset \{1, 2, \dots, n\}$ .

In particular, put  $S = \{1, 2, \dots, n\}$ . Since

$$T^{\{1,2,\dots,n\}} = (T_1^{\{1,2,\dots,n\}}, T_2^{\{1,2,\dots,n\}}, \dots, T_n^{\{1,2,\dots,n\}})$$

is in  $End(H, f)$  and  $(H, f)$  is transitive, there exists a constant  $\mu^{\{1,2,\dots,n\}} \in \mathbb{C}$  such that

$$T_i^{\{1,2,\dots,n\}} = E_{i-1}\cdots E_1E_nE_{n-1}\cdots E_{i+1}E_i = \mu^{\{1,2,\dots,n\}}I_{H_i}.$$

Take  $x (\neq 0) \in H_1$ . Since  $H_1 \subset H_j$  for any  $1 \leq j \leq n$  and  $E_n\cdots E_1 = \mu^{\{1,2,\dots,n\}}I_{H_1}$ , we have that  $x = \mu^{\{1,2,\dots,n\}}x$ . Hence  $\mu^{\{1,2,\dots,n\}} = 1$ .

We shall show that  $H_1 = H_2 = \cdots = H_n$ . On the contrary we assume that  $H_k \neq H_\ell$  for some  $k \neq \ell$ . Using Lemma 4.4, we can represent  $H_i$  as  $H_i = K_1 \oplus K_2 \oplus \cdots \oplus K_{m(i)}$  and  $m(1) = 1$ . Then there exists the smallest  $i$  such that  $m(i) > 1$ . We compare  $m(j)$  and  $m(i)$ . If there exists  $m(j)$  such that  $m(j) < m(i)$  ( $i \leq j \leq n$ ). Take  $x (\neq 0) \in K_{m(i)} \subset H_i$ . Then  $E_{j-1}E_{j-2}\cdots E_{i+1}E_ix = 0$ . This contradicts that  $E_{i-1}\cdots E_1E_nE_{n-1}\cdots E_{i+1}E_i = I_{H_i}$ .

If there exists no  $m(j)$  such that  $m(j) < m(i)$  ( $i \leq j \leq n$ ). Take  $x (\neq 0) \in K_{m(i)} \subset H_i$ . Then  $E_{n-1}E_{n-2}\cdots E_{i+1}E_ix = x$ , and  $E_nx = 0$ . This also contradicts that

$$E_{i-1}\cdots E_1E_nE_{n-1}\cdots E_{i+1}E_i = I_{H_i}.$$



Therefore we have that  $H_1 = H_2 = \dots = H_n =: M$ . Moreover we also have that  $E_1 = E_2 = \dots = E_n = I_M$ . In particular,  $T_i^{\{1,2,\dots,n\}} = I_M$  for any  $i$  and

$$A_i = E_{i-1} \cdots E_1 E_n E_{n-1} \cdots E_{i+1} A_i = T_i^{\{1,2,\dots,n\} \setminus k} = \mu^{\{1,2,\dots,n\} \setminus k} I_{H_k}.$$

We shall show that  $\dim M = 1$ . On the contrary, assume that  $\dim M \geq 2$ . Then there exists a non-scalar operator  $B \in B(M)$ . Since each  $A_k$  is a scalar operator for any  $k$ ,  $(B, \dots, B)$  is in  $End(H, f)$ . This contradicts to that  $(H, f)$  is transitive. Therefore  $\dim M = 1$ . Hence we may assume that  $H_i = \mathbb{C}$  for any  $i$ . Since  $(H, f)$  is transitive, there exists only one  $(H, f)$ -connected component on  $V = \{1, 2, \dots, n\}$  by Lemma 4.3.

Next we consider the case that there exists  $H_i = 0$  for some  $i$ . If there exists only one vertex  $i$  such that  $H_i \neq 0$ , then  $\dim H_i = 1$  because  $(H, f)$  is transitive. Therefore we may assume that there exists more than two vertices  $i$  such that  $H_i \neq 0$ .

We consider the reduction of the quiver  $C_n$  to the set of vertices  $i$  with  $H_i \neq 0$  to get another quiver  $C_m$  ( $2 \leq m \leq n$ ).

Let  $(K, g)$  be the reduced Hilbert representation of  $C_m$  from the Hilbert representation  $(H, f)$  of  $C_n$  by Lemma 4.2. Then  $End(H, f)$  is isomorphic to  $End(K, g)$ . Since  $(H, f)$  is transitive,  $(K, g)$  is also transitive.

Since we can adapt the above consideration to  $(K, g)$ , we have that  $H_i = \mathbb{C}$  for all  $i$  such that  $H_i \neq 0$ . Therefore in  $(H, f)$ , we may and do have that  $H_i = \mathbb{C}$  or  $0$  for  $1 \leq i \leq n$ . Since  $(H, f)$  is transitive, by Lemma 4.3, there exists only one  $(H, f)$ -connected component  $\{i \in V; H_i \neq 0\}$ .  $\square$


**Theorem 4.10.** *Let  $\Gamma$  be a quiver whose underlying undirected graph is an extended Dynkin diagram  $\widetilde{A}_n$ ,  $n \geq 0$ . Then there exists an infinite-dimensional transitive Hilbert representation of  $\Gamma$  if and only if  $\Gamma$  is not an oriented cyclic quiver.*

**Proof.** Assume that  $\Gamma$  is not an oriented cyclic quiver. By Theorem 4.1, there exists an infinite-dimensional transitive Hilbert representation of  $\Gamma$ . Conversely, assume that  $\Gamma$  is an oriented cyclic quiver. Then transitive Hilbert representations of  $\Gamma$  are finite-dimensional by Lemma 4.9. Hence there exist no infinite-dimensional transitive Hilbert representations of  $\Gamma$ .  $\square$

Gabriel’s theorem states that a finite, connected quiver has only finitely many indecomposable representations if and only if the underlying undirected graph is one of Dynkin diagrams  $A_n, D_n, E_6, E_7, E_8$ . In [EW3], we constructed some examples of indecomposable, infinite-dimensional representations of quivers with the underlying undirected graphs being extended Dynkin diagrams  $\widetilde{D}_n$  ( $n \geq 4$ ),  $\widetilde{E}_6, \widetilde{E}_7$  and  $\widetilde{E}_8$ . We used the quivers whose vertices are represented by a family of subspaces and whose arrows are represented by natural inclusion maps. Replacing the unilateral shift  $S$

with a transitive operator in the construction of examples of indecomposable, infinite-dimensional representations of quivers in [EW3], we shall give some examples of infinite-dimensional transitive representations of quivers with the underlying undirected graphs being extended Dynkin diagrams  $\tilde{D}_n$  ( $n \geq 4$ ),  $\tilde{E}_6$ ,  $\tilde{E}_7$  and  $\tilde{E}_8$ . Our construction of examples is considered as a modification of an unbounded operator used by Harrison, Radjavi and Rosenthal [HRR] to provide a transitive lattice.

**Lemma 4.11.** *Let  $\Gamma = (V, E, s, r)$  be the following quiver with the underlying undirected graph an extended Dynkin diagram  $\tilde{D}_n$  for  $n \geq 4$  :*




Then there exists an infinite-dimensional, transitive Hilbert representation  $(H, f)$  of  $\Gamma$ .

**Proof.** Let  $K = \ell^2(\mathbb{N})$  and  $S$  a transitive operator on  $K$  with the domain  $D(S)$ . We define a Hilbert representation  $(H, f) := ((H_v)_{v \in V}, (f_\alpha)_{\alpha \in E})$  of  $\Gamma$  as follows:  $H_1 = K \oplus 0$ ,  $H_2 = 0 \oplus K$ ,  $H_3 = \{(x, Sx) \in K \oplus K; x \in D(S)\}$ ,  $H_4 = \{(x, x) \in K \oplus K; x \in K\}$ , and  $H_5 = H_6 = \cdots = H_{n+1} = K \oplus K$ .

Let  $f_{\alpha_k} : H_{s(\alpha_k)} \rightarrow H_{r(\alpha_k)}$  be the inclusion map for any  $\alpha_k \in E$  for  $k = 1, 2, 3, 4$ , and  $f_\beta = id$  for other arrows  $\beta \in E$ . Take  $T = (T_v)_{v \in V} \in \text{End}(H, f)$ . Since  $T \in \text{End}(H, f)$  and any arrow is represented by the inclusion map, we have  $T_i = T_j$  ( $i = 5, \dots, n+1$ ),  $T_5x = T_vx$  for any  $v \in \{1, 2, 4\}$ , any  $x \in H_v$ . In particular,  $T_5H_v \subset H_v$  ( $v \in \{1, 2, 4\}$ ). Hence  $T_5$  is written as  $T_5 = A \oplus A$  as in [EW3, Lemma 6.1, Example 3]. Moreover  $H_3$  is also invariant under  $T_5$ . Since  $S$  is transitive, we have that  $A$  is a scalar by Lemma 3.3. Thus  $T$  is a scalar, that is,  $\text{End}(H, f) = \mathbb{C}$ . Therefore  $(H, f)$  is transitive.  $\square$

**Lemma 4.12.** *Let  $\Gamma = (V, E, s, r)$  be the following quiver with the underlying undirected graph an extended Dynkin diagram  $\tilde{E}_6$  :*




Then there exists an infinite-dimensional, transitive Hilbert representation  $(H, f)$  of  $\Gamma$ .

**Proof.** Let  $(H, f) = ((H_v)_{v \in V}, (f_\alpha)_{\alpha \in E})$  be the following Hilbert representation of  $\Gamma$ : Let  $K = \ell^2(\mathbb{N})$  and  $S$  a transitive operator on  $K$  with the domain  $D(S)$ . Define  $H_0 = K \oplus K \oplus K$ ,  $H_1 = 0 \oplus K \oplus K$ ,  $H_2 = 0 \oplus \{(y, Sy) \in K^2; y \in D(S)\}$ ,  $H_{1'} = K \oplus K \oplus 0$ ,  $H_{2'} = \{(x, x) \in K^2; x \in K\} \oplus 0$ ,

$H_{1''} = K \oplus 0 \oplus K$ , and  $H_{2''} = \{(x, 0, x) \in K^3; x \in K\}$ . For any arrow  $\alpha \in E$ , let  $f_\alpha : H_{s(\alpha)} \rightarrow H_{r(\alpha)}$  be the canonical inclusion map. Take  $T = (T_v)_{v \in V} \in \text{End}(H, f)$ . Since any arrow is represented by the inclusion map, we have  $T_0x = T_vx$  for any  $v \in \{1, 1', 2', 1'', 2''\}$  and any  $x \in H_v$ . In particular,  $T_0H_v \subset H_v$ . Hence  $T_0$  is written as  $T_0 = A \oplus A \oplus A$ . Moreover  $H_2 = \{(0, x, Sx) \in K^3; x \in D(S)\}$  is also invariant under  $T_0$ . Hence for any  $x \in D(S)$ , there exists  $y \in D(S)$  such that  $(0, Ax, ASx) = (0, y, Sy)$  as in [EW3, Example 4]. Since  $S$  is transitive, we have that  $A$  is a scalar by Lemma 3.3. Thus  $T$  is a scalar, that is,  $\text{End}(H, f) = \mathbb{C}$ . Therefore  $(H, f)$  is transitive.  $\square$


**Lemma 4.13.** *Let  $\Gamma = (V, E, s, r)$  be the following quiver with the underlying undirected graph an extended Dynkin diagram  $\tilde{E}_7$  :*



Then there exists an infinite-dimensional, transitive Hilbert representation  $(H, f)$  of  $\Gamma$ .

**Proof.** Let  $K = \ell^2(\mathbb{N})$  and  $S$  a transitive operator on  $K$  with the domain  $D(S)$ . Define a Hilbert representation  $(H, f) := ((H_v)_{v \in V}, (f_\alpha)_{\alpha \in E})$  of  $\Gamma$  as follows: Let  $H_0 = K \oplus K \oplus K \oplus K$ ,  $H_1 = K \oplus 0 \oplus K \oplus K$ ,  $H_2 = K \oplus 0 \oplus \{(x, x); x \in K\}$ ,  $H_3 = K \oplus 0 \oplus 0 \oplus 0$ ,  $H_{1'} = 0 \oplus K \oplus K \oplus K$ ,  $H_{2'} = 0 \oplus K \oplus \{(y, Sy) \in K^2; y \in D(S)\}$ ,  $H_{3'} = 0 \oplus K \oplus 0 \oplus 0$ , and  $H_{1''} = \{(x, y, x, y) \in K^4; x, y \in K\}$ . For any arrow  $\alpha \in E$ , let  $f_\alpha : H_{s(\alpha)} \rightarrow H_{r(\alpha)}$  be the canonical inclusion map. Take  $T = (T_v)_{v \in V} \in \text{End}(H, f)$ . Since any arrow is represented by the inclusion map, we have  $T_0x = T_vx$  for any  $v \in \{1, 2, 3, 1', 2', 3', 1''\}$  and any  $x \in H_v$ . In particular,  $T_0H_v \subset H_v$ . Hence  $T_0$  is written as  $T_0 = A \oplus A \oplus A \oplus A$ . Moreover  $H_1 \cap H_{2'} = \{(0, 0, x, Sx) \in K^4; x \in D(S)\}$  is also invariant under  $T_0$ . Hence for any  $x \in D(S)$ , there exists  $y \in D(S)$  such that  $(0, 0, Ax, ASx) = (0, 0, y, Sy)$  as in [EW3, Lemma 6.2]. Since  $S$  is transitive, we have that  $A$  is a scalar by Lemma 3.3. Thus  $T$  is a scalar, that is,  $\text{End}(H, f) = \mathbb{C}$ . Therefore  $(H, f)$  is transitive.  $\square$

**Lemma 4.14.** *Let  $\Gamma = (V, E, s, r)$  be the following quiver with the underlying undirected graph an extended Dynkin diagram  $\tilde{E}_8$  :*



Then there exists an infinite-dimensional, transitive Hilbert representation  $(H, f)$  of  $\Gamma$ .

**Proof.** Let  $K = \ell^2(\mathbb{N})$  and  $S$  a transitive operator on  $K$  with the domain  $D(S)$ . We define a Hilbert representation  $(H, f) := ((H_v)_{v \in V}, (f_\alpha)_{\alpha \in E})$  of  $\Gamma$  as follows: Let  $H_0 = K \oplus K \oplus K \oplus K \oplus K \oplus K$ ,  $H_1 = \{(x, x) \in K^2; x \in$

$K\} \oplus K \oplus K \oplus K \oplus K, H_2 = 0 \oplus 0 \oplus K \oplus K \oplus K \oplus K, H_3 = 0 \oplus 0 \oplus 0 \oplus K \oplus K \oplus K,$   
 $H_4 = 0 \oplus 0 \oplus 0 \oplus K \oplus \{(y, Sy) \in K^2; y \in D(S)\}, H_5 = 0 \oplus 0 \oplus 0 \oplus K \oplus 0 \oplus 0,$   
 $H_{1'} = K \oplus K \oplus \{(x, y, x, y) \in K^4; x, y \in K\}, H_{2'} = K \oplus K \oplus 0 \oplus 0 \oplus 0 \oplus 0,$   
 and  $H_{1''} = \{(y, z, x, 0, y, z) \in K^6; x, y, z \in K\}$ . For any arrow  $\alpha \in E$ , let  $f_\alpha : H_{s(\alpha)} \rightarrow H_{r(\alpha)}$  be the canonical inclusion map.

Take  $T = (T_v)_{v \in V} \in \text{End}(H, f)$ . Since any arrow is represented by the inclusion map, we have  $T_0x = T_vx$  for any  $v \in V$  and any  $x \in H_v$ . In particular,  $T_0H_v \subset H_v$ . Since  $T_0$  preserves subspaces  $H_v, v = 1, 1', 1'', 2, 2', 3, 5,$   $T_0$  is written as

$$T_0 = A \oplus A \oplus A \oplus A \oplus A \oplus A.$$

Finally,  $H_4 = 0 \oplus 0 \oplus 0 \oplus K \oplus \{(y, Sy) \in K^2; y \in D(S)\}$  is invariant under  $T_0$ . Then for any  $x \in K$  and  $y \in D(S)$ , there exist  $x' \in K$  and  $y' \in D(S)$  such that

$$T_0(0, 0, 0, x, y, Sy) = (0, 0, 0, Ax, Ay, ASy) = (0, 0, 0, x', y', Sy').$$

Hence  $ASy = Sy' = SAy$  as in [EW3, Lemma 6.3]. Since  $S$  is transitive, we have that  $A$  is a scalar by Lemma 3.3.

Thus  $T = (T_v)_{v \in V}$  is a scalar, that is,  $\text{End}(H, f) = \mathbb{C}$ . Therefore,  $(H, f)$  is transitive. □

Next, we shall investigate the endomorphism algebras of Hilbert representations. At first we recall some facts about reflection functors from [EW3].

Reflection functors are crucially used in the proof of the classification of finite-dimensional, indecomposable representations of tame quivers (cf. [As], [BGP], [DIR],[DoF], [GaR], [GeP]). As a matter of fact, many indecomposable representations of tame quivers can be reconstructed by iterating reflection functors on simple indecomposable representations. We can not expect such a best position in infinite-dimensional Hilbert representations. But reflection functors are still valuable to show that some property of representations of quivers on infinite-dimensional Hilbert spaces does not depend on the choice of orientations and does depend on the fact underlying undirected graphs are (extended) Dynkin diagrams or not.

Let  $\Gamma = (V, E, s, r)$  be a finite quiver. We say that a vertex  $v \in V$  is a *sink* if  $v \neq s(\alpha)$  for any  $\alpha \in E$ . Put  $E^v = \{\alpha \in E; r(\alpha) = v\}$ . We denote by  $\overline{E}$  the set of all formally reversed new arrows  $\overline{\alpha}$  for  $\alpha \in E$ . In this way if  $\alpha : x \rightarrow y$  is an arrow, then  $\overline{\alpha} : x \leftarrow y$ .

**Definition.**[EW3] Let  $\Gamma = (V, E, s, r)$  be a finite quiver. For a sink  $v \in V$ , we construct a new quiver  $\sigma_v^+(\Gamma) = (\sigma_v^+(V), \sigma_v^+(E), s, r)$  as follows: All the arrows of  $\Gamma$  having  $v$  as range are reversed and all the other arrows remain unchanged. That is,

$$\sigma_v^+(V) = V \quad \sigma_v^+(E) = (E \setminus E^v) \cup \overline{E^v},$$

where  $\overline{E^v} = \{\overline{\alpha}; \alpha \in E^v\}$ .

**Definition.** [EW3] (reflection functor  $\Phi_v^+$ .) Let  $\Gamma = (V, E, s, r)$  be a finite quiver. For a sink  $v \in V$ , we define a *reflection functor* at  $v$

$$\Phi_v^+ : HRep(\Gamma) \rightarrow HRep(\sigma_v^+(\Gamma))$$

between the categories of Hilbert representations of  $\Gamma$  and  $\sigma_v^+(\Gamma)$  as follows: For a Hilbert representation  $(H, f)$  of  $\Gamma$ , we define a Hilbert representation  $(K, g) = \Phi_v^+(H, f)$  of  $\sigma_v^+(\Gamma)$ . Let

$$h_v : \bigoplus_{\alpha \in E^v} H_{s(\alpha)} \rightarrow H_v$$

be a bounded linear operator defined by

$$h_v((x_\alpha)_{\alpha \in E^v}) = \sum_{\alpha \in E^v} f_\alpha(x_\alpha).$$

We shall define

$$K_v := \text{Ker } h_v = \{(x_\alpha)_{\alpha \in E^v} \in \bigoplus_{\alpha \in E^v} H_{s(\alpha)}; \sum_{\alpha \in E^v} f_\alpha(x_\alpha) = 0\}.$$

We also consider the canonical inclusion map  $i_v : K_v \rightarrow \bigoplus_{\alpha \in E^v} H_{s(\alpha)}$ . For  $u \in V$  with  $u \neq v$ , put  $K_u = H_u$ .

For  $\beta \in E^v$ , let

$$P_\beta : \bigoplus_{\alpha \in E^v} H_{s(\alpha)} \rightarrow H_{s(\beta)}$$

be the canonical projection. Then we shall define

$$g_{\bar{\beta}} : K_{s(\bar{\beta})} = K_v \rightarrow K_{r(\bar{\beta})} = H_{s(\beta)} \quad \text{by } g_{\bar{\beta}} = P_\beta \circ i_v$$

that is,  $g_{\bar{\beta}}((x_\alpha)_{\alpha \in E^v}) = x_\beta$ .

For  $\beta \notin E^v$ , let  $g_\beta = f_\beta$ . For a homomorphism  $T : (H, f) \rightarrow (H', f')$ , we define a homomorphism

$$S = (S_u)_{u \in V} = \Phi_v^+(T) : (K, g) = \Phi_v^+(H, f) \rightarrow (K', g') = \Phi_v^+(H', f')$$

If  $u = v$ , a bounded operator  $S_v : K_v \rightarrow K'_v$  is given by

$$S_v((x_\alpha)_{\alpha \in E^v}) = (T_{s(\alpha)}(x_\alpha))_{\alpha \in E^v}.$$

It is easily seen that  $S_v$  is well-defined and we have the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_v & \xrightarrow{i_v} & \bigoplus_{\alpha \in E^v} H_{s(\alpha)} & \xrightarrow{h_v} & H_v \\ & & S_v \downarrow & & (T_{s(\alpha)})_{\alpha \in E^v} \downarrow & & T_v \downarrow \\ 0 & \longrightarrow & K'_v & \xrightarrow{i'_v} & \bigoplus_{\alpha \in E^v} H'_{s(\alpha)} & \xrightarrow{h'_v} & H'_v \end{array}$$

For other  $u \in V$  with  $u \neq v$ , put

$$S_u = T_u : K_u = H_u \rightarrow K'_u = H'_u.$$

We also consider a dual of the above construction. We say that a vertex  $v \in V$  is a *source* if  $v \neq r(\alpha)$  for any  $\alpha \in E$ . Put  $E_v = \{\alpha \in E; s(\alpha) = v\}$ .

**Definition.**[EW3] Let  $\Gamma = (V, E, s, r)$  be a finite quiver. For a source  $v \in V$ , we shall construct a new quiver  $\sigma_v^-(\Gamma) = (\sigma_v^-(V), \sigma_v^-(E), s, r)$  as

follows: All the arrows of  $\Gamma$  having  $v$  as source are reversed and all the other arrows remain unchanged. That is,

$$\sigma_v^-(V) = V \quad \sigma_v^-(E) = (E \setminus E_v) \cup \overline{E_v},$$

where  $\overline{E_v} = \{\overline{\alpha}; \alpha \in E_v\}$ .

**Definition.**[EW3] (reflection functor  $\Phi_v^-$ .) Let  $\Gamma = (V, E, s, r)$  be a finite quiver. For a source  $v \in V$ , we shall define a *reflection functor* at  $v$

$$\Phi_v^- : HRep(\Gamma) \rightarrow HRep(\sigma_v^-(\Gamma))$$

between the categories of Hilbert representations of  $\Gamma$  and  $\sigma_v^-(\Gamma)$  as follows: For a Hilbert representation  $(H, f)$  of  $\Gamma$ , we define a Hilbert representation  $(K, g) = \Phi_v^-(H, f)$  of  $\sigma_v^-(\Gamma)$ . Let

$$\hat{h}_v : H_v \rightarrow \bigoplus_{\alpha \in E_v} H_{r(\alpha)}$$

be a bounded linear operator defined by

$$\hat{h}_v(x) = (f_\alpha(x))_{\alpha \in E_v} \quad \text{for } x \in H_v.$$

We shall define

$$K_v := (\text{Im } \hat{h}_v)^\perp = \text{Ker } \hat{h}_v^* \subset \bigoplus_{\alpha \in E_v} H_{r(\alpha)},$$

where  $\hat{h}_v^* : \bigoplus_{\alpha \in E_v} H_{r(\alpha)} \rightarrow H_v$  is given  $\hat{h}_v^*((x_\alpha)_{\alpha \in E_v}) = \sum f_\alpha^*(x_\alpha)$ . For  $u \in V$  with  $u \neq v$ , put  $K_u = H_u$ .

Let  $Q_v : \bigoplus_{\alpha \in E_v} H_{r(\alpha)} \rightarrow K_v$  be the canonical projection. For  $\beta \in E_v$ , let

$$j_\beta : H_{r(\beta)} \rightarrow \bigoplus_{\alpha \in E_v} H_{r(\alpha)}$$

be the canonical inclusion. We shall define

$$g_{\overline{\beta}} : K_{s(\overline{\beta})} = H_{r(\beta)} \rightarrow K_{r(\overline{\beta})} = K_v \quad \text{by } g_{\overline{\beta}} = Q_v \circ j_\beta.$$

For  $\beta \notin E_v$ , let  $g_\beta = f_\beta$ .

For a homomorphism  $T : (H, f) \rightarrow (H', f')$ , we shall define a homomorphism

$$S = (S_u)_{u \in V} = \Phi_v^-(T) : (K, g) = \Phi_v^-(H, f) \rightarrow (K', g') = \Phi_v^-(H', f').$$

For  $u = v$ , a bounded operator  $S_v : K_v \rightarrow K'_v$  is given by

$$S_v((x_\alpha)_{\alpha \in E_v}) = Q'_v((T_{r(\alpha)}(x_\alpha))_{\alpha \in E_v}),$$

where  $Q'_v : \bigoplus_{\alpha \in E_v} H'_{r(\alpha)} \rightarrow K'_v$  be the canonical projection.

We have the following commutative diagram:

$$\begin{array}{ccccccc} H_v & \xrightarrow{\hat{h}_v} & \bigoplus_{\alpha \in E_v} H_{r(\alpha)} & \xrightarrow{Q_v} & K_v & \longrightarrow & 0 \\ T_v \downarrow & & \bigoplus_{\alpha \in E_v} T_{r(\alpha)} \downarrow & & S_v \downarrow & & \\ H'_v & \xrightarrow{\hat{h}'_v} & \bigoplus_{\alpha \in E_v} H'_{r(\alpha)} & \xrightarrow{Q'_v} & K'_v & \longrightarrow & 0 \end{array}$$

For other  $u \in V$  with  $u \neq v$ , put

$$S_u = T_u : K_u = H_u \rightarrow K'_u = H'_u.$$

We shall describe a relation between two (covariant) functors  $\Phi_v^+$  and  $\Phi_v^-$ . We shall define another (contravariant) functor  $\Phi^*$  at the beginning.

Let  $\Gamma = (V, E, s, r)$  be a finite quiver. We shall define the opposite quiver  $\bar{\Gamma} = (\bar{V}, \bar{E}, s, r)$  by reversing all the arrows, more precisely,  $\bar{V} = V$  and  $\bar{E} = \{\bar{\alpha}; \alpha \in E\}$ .

**Definition.**[EW3] Let  $\Gamma = (V, E, s, r)$  be a finite quiver and  $\bar{\Gamma} = (\bar{V}, \bar{E}, s, r)$  its opposite quiver. We shall define a contravariant functor

$$\Phi^* : HRep(\Gamma) \rightarrow HRep(\bar{\Gamma})$$

between the categories of Hilbert representations of  $\Gamma$  and  $\bar{\Gamma}$  as follows: For a Hilbert representation  $(H, f)$  of  $\Gamma$ , we define a Hilbert representation  $(K, g) = \Phi^*(H, f)$  of  $\bar{\Gamma}$  by  $K_u = H_u$  for  $u \in V$  and  $g_{\bar{\alpha}} = f_{\alpha}^*$  for  $\alpha \in E$ . For a homomorphism  $T : (H, f) \rightarrow (H', f')$ , we define a homomorphism

$$S = (S_u)_{u \in V} = \Phi^*(T) : (K', g') = \Phi^*(H', f') \rightarrow (K, g) = \Phi^*(H, f),$$

by bounded operators  $S_u : K'_u = H'_u \rightarrow K_u = H_u$  given by  $S_u = T_u^*$ .

**Proposition 4.15.** [EW3, Proposition 4.2.] *Let  $\Gamma = (V, E, s, r)$  be a finite quiver. If  $v \in V$  is a source of  $\Gamma$ , then  $v$  is a sink of  $\bar{\Gamma}$ ,  $\sigma_v^-(\Gamma) = \sigma_v^+(\bar{\Gamma})$  and the following assertions hold:*

- (1) *For a Hilbert representation  $(H, f)$  of  $\Gamma$ ,*

$$\Phi_v^-(H, f) = \Phi^*(\Phi_v^+(\Phi^*(H, f))).$$

- (2) *For a homomorphism  $T : (H, f) \rightarrow (H', f')$ ,*

$$\Phi_v^-(T) = \Phi^*(\Phi_v^+(\Phi^*(T))).$$

**Proposition 4.16.** [EW3, Proposition 4.3.] *Let  $\Gamma = (V, E, s, r)$  be a finite quiver. If  $v \in V$  is a sink of  $\Gamma$ , then  $v$  is a source of  $\bar{\Gamma}$ ,  $\sigma_v^+(\Gamma) = \sigma_v^-(\bar{\Gamma})$  and the following assertions hold:*

- (1) *For a Hilbert representation  $(H, f)$  of  $\Gamma$ ,*

$$\Phi_v^+(H, f) = \Phi^*(\Phi_v^-(\Phi^*(H, f))).$$

- (2) *For a homomorphism  $T : (H, f) \rightarrow (H', f')$ ,*

$$\Phi_v^+(T) = \Phi^*(\Phi_v^-(\Phi^*(T))).$$

We shall investigate endomorphisms of Hilbert representations and its images of reflection functors. In the case of infinite-dimensional Hilbert representations, we need to assume a certain closedness condition at a sink or a source.

**Definition.**[EW3] Let  $\Gamma = (V, E, s, r)$  be a finite quiver and  $v \in V$  a sink. We recall that  $E^v = \{\alpha; r(\alpha) = v\}$ . It is said that a Hilbert representation  $(H, f)$  of  $\Gamma$  is *closed* at  $v$  if  $\sum_{\alpha \in E^v} \text{Im} f_{\alpha} \subset H_v$  is a closed subspace. It is said that  $(H, f)$  is *full* at  $v$  if  $\sum_{\alpha \in E^v} \text{Im} f_{\alpha} = H_v$ .

**Definition.**([EW3]) Let  $\Gamma = (V, E, s, r)$  be a finite quiver and  $v \in V$  a source. We recall that  $E_v = \{\alpha | s(\alpha) = v\}$ . It is said that a Hilbert

representation  $(H, f)$  of  $\Gamma$  is *co-closed* at  $v$  if  $\sum_{\alpha \in E^v} \text{Im} f_\alpha^* \subset H_v$  is a closed subspace. It is said that  $(H, f)$  is *co-full* at  $v$  if  $\sum_{\alpha \in E^v} \text{Im} f_\alpha^* = H_v$ .

We note that the properties of fullness, co-fullness, closedness and co-closedness are preserved under isomorphism of Hilbert representations.

**Lemma 4.17.** *Let  $\Gamma$  be a finite quiver and  $v \in \Gamma$  a sink. Let  $(H, f)$  and  $(K, g)$  be isomorphic Hilbert representations of  $\Gamma$ . If  $(H, f)$  is full (resp. closed) at  $v$ , then  $(K, g)$  is full (resp. closed) at  $v$ .*

**Proof.** Assume that  $(H, f)$  is full at  $v$ . Since  $(H, f)$  and  $(K, g)$  are isomorphic, there exists a family  $S = (S_u)_{u \in V}$  of bounded invertible operators such that  $S_{r(\alpha)} f_\alpha = g_\alpha S_{s(\alpha)}$  for  $\alpha \in E$ . Take an element  $y \in K_v$ . By the invertibility of  $S_v$ , there exists an element  $x \in H_v$  such that  $S_v(x) = y$ . Since  $(H, f)$  is full at  $v$ , there exist  $x_{s(\alpha)} \in H_{s(\alpha)}$  such that  $\sum_{\alpha \in E^v} f_\alpha(x_{s(\alpha)}) = x$ . We put  $y_{s(\alpha)} := S_{s(\alpha)}(x_{s(\alpha)})$ . Then

$$\begin{aligned} \sum_{\alpha \in E^v} g_\alpha(y_{s(\alpha)}) &= \sum_{\alpha \in E^v} g_\alpha S_{s(\alpha)}(x_{s(\alpha)}) = \sum_{\alpha \in E^v} S_v f_\alpha(x_{s(\alpha)}) \\ &= S_v \sum_{\alpha \in E^v} f_\alpha(x_{s(\alpha)}) = S_v(x) = y \end{aligned}$$

Hence  $(K, g)$  is full at  $v$ .

We can similarly prove that closedness property is preserved under isomorphism of Hilbert representations. □

**Lemma 4.18.** *Let  $\Gamma$  be a finite quiver and  $v \in V$  a source. Let  $(H, f)$  and  $(K, g)$  be isomorphic Hilbert representations of  $\Gamma$ . If  $(H, f)$  is co-full (resp. co-closed) at  $v$ , then  $(K, g)$  is co-full (resp. co-closed) at  $v$ .*

**Proof.** Since  $(H, f)$  and  $(K, g)$  are isomorphic,  $\Phi^*(H, f)$  and  $\Phi^*(K, g)$  are isomorphic. Hence the case of co-fullness is reduced to the case of fullness. We can similarly prove that co-closedness property is preserved under isomorphism of Hilbert representations. □

The following theorem is well known for finite-dimensional Hilbert spaces ([As, page289, 5.7. Corollary] and [DIR, page16, Proposition 2.1]).

**Theorem 4.19.** *Let  $\Gamma = (V, E, s, r)$  be a finite quiver and  $v \in V$  a sink. If a Hilbert representation  $(H, f)$  of  $\Gamma$  is full at  $v$ , then the map  $\Phi_v^+ : \text{End}(H, f) \rightarrow \text{End}(\Phi_v^+(H, f))$  is an isomorphism as  $\mathbb{C}$ -algebras.*

**Proof.** We put  $(K, g) := \Phi_v^+(H, f)$ . The mapping  $\Phi_v^+$  gives a mapping of  $\text{End}(H, f)$  to  $\text{End}(K, g)$ . At first we shall show that  $\Phi_v^+$  is one to one. Assume that  $S := \Phi_v^+(T) = 0$  for  $T \in \text{End}(H, f)$ . We have  $S_u = T_u = 0$  ( $u \neq v$ ). From this we shall show that  $T_v = 0$ . Since  $T \in \text{End}(H, f)$ ,  $T_v f_\alpha = f_\alpha T_{s(\alpha)}$  for  $\alpha \in E^v = \{\alpha \in E; r(\alpha) = v\}$ . Hence, for  $x_\alpha \in H_{s(\alpha)}$ ,

$$T_v \left( \sum_{\alpha \in E^v} f_\alpha(x_\alpha) \right) = \sum_{\alpha \in E^v} f_\alpha T_{s(\alpha)}(x_\alpha) = 0.$$



Since  $(H, f)$  is full at  $v$ ,  $T_v = 0$ . Thus  $\Phi_v^+$  is one to one. Next we shall show that  $\Phi_v^+$  is onto. Take  $S = (S_u)_{u \in V} \in \text{End}(K, g)$ . We put  $T_u = S_u$  for  $u \neq v$ . We shall define an operator  $T_v : H_v \rightarrow H_v$  such that  $T_v(\sum_{\alpha \in E^v} f_\alpha(x_\alpha)) = \sum_{\alpha \in E^v} f_\alpha(T_{s(\alpha)}(x_\alpha))$  for  $x_\alpha \in H_{s(\alpha)}$ . We need to show that  $T_v$  is well defined. If there exists an element  $(x'_\alpha)_{\alpha \in E^v} \in \bigoplus_{\alpha \in E^v} H_{s(\alpha)}$  such that

$$\sum_{\alpha \in E^v} f_\alpha(x_\alpha) = \sum_{\alpha \in E^v} f_\alpha(x'_\alpha),$$

then we must show that

$$\sum_{\alpha \in E^v} f_\alpha T_{s(\alpha)}(x_\alpha) = \sum_{\alpha \in E^v} f_\alpha T_{s(\alpha)}(x'_\alpha).$$

Since  $\sum_{\alpha \in E^v} f_\alpha(x_\alpha) = \sum_{\alpha \in E^v} f_\alpha(x'_\alpha)$ , we have

$$h_v((x_\alpha - x'_\alpha)_{\alpha \in E^v}) = \sum_{\alpha \in E^v} f_\alpha(x_\alpha - x'_\alpha) = 0.$$

Hence  $(x_\alpha - x'_\alpha)_{\alpha \in E^v} \in \ker h_v = K_v$ . Since  $S_v : K_v \rightarrow K_v$ , we have  $S_v((x_\alpha - x'_\alpha)_{\alpha \in E^v}) \in \ker h_v = K_v$ . Hence  $h_v(S_v((x_\alpha - x'_\alpha)_{\alpha \in E^v})) = 0$ . Since  $S \in \text{End}(K, g)$ , we have  $S_{s(\alpha)}g_{\bar{\alpha}} = g_{\bar{\alpha}}S_v$  for  $\alpha \in E^v$ ,

$$S_{s(\alpha)}g_{\bar{\alpha}}((x_\beta - x'_\beta)_{\beta \in E^v}) = S_{s(\alpha)}(x_\alpha - x'_\alpha) = T_{s(\alpha)}(x_\alpha - x'_\alpha),$$

and

$$g_{\bar{\alpha}}S_v((x_\beta - x'_\beta)_{\beta \in E^v}) = P_\alpha S_v((x_\beta - x'_\beta)_{\beta \in E^v}).$$

Hence

$$T_{s(\alpha)}(x_\alpha - x'_\alpha) = P_\alpha S_v((x_\beta - x'_\beta)_{\beta \in E^v}).$$

Then

$$\sum_{\alpha \in E^v} f_\alpha T_{s(\alpha)}(x_\alpha - x'_\alpha) = \sum_{\alpha \in E^v} f_\alpha P_\alpha S_v((x_\beta - x'_\beta)_{\beta \in E^v})$$

and

$$\sum_{\alpha \in E^v} f_\alpha P_\alpha S_v((x_\beta - x'_\beta)_{\beta \in E^v}) = h_v(S_v((x_\beta - x'_\beta)_{\beta \in E^v})) = 0.$$

This gives

$$\sum_{\alpha \in E^v} f_\alpha T_{s(\alpha)}(x_\alpha) = \sum_{\alpha \in E^v} f_\alpha T_{s(\alpha)}(x'_\alpha).$$

Thus  $T_v$  is well defined.

Next we shall show that  $T_v f_\alpha(x) = f_\alpha T_{s(\alpha)}(x)$  for  $x \in H_{s(\alpha)}$ . Take and fix  $x \in H_{s(\alpha)}$  for  $\alpha \in E^v$ . For  $\beta \in E^v$ , we put

$$x_\beta = \begin{cases} x & (\beta = \alpha), \\ 0 & (\beta \neq \alpha). \end{cases}$$

Since  $T_v(\sum_{\beta \in E^v} f_\beta(x_\beta)) = \sum_{\beta \in E^v} f_\beta(T_{s(\beta)}(x_\beta))$ , we have

$$T_v f_\alpha(x) = \sum_{\beta} f_\beta T_{s(\beta)}(x_\beta) = f_\alpha T_{s(\alpha)}(x).$$

Next we shall show that  $T_v : H_v \rightarrow H_v$  is bounded. We decompose

$$\bigoplus_{\alpha \in E^v} H_{s(\alpha)} = \ker h_v \oplus (\ker h_v)^\perp = K_v \oplus K_v^\perp.$$

By the Banach invertibility theorem,  $h_v|_{(K_v)^\perp} : (K_v)^\perp \rightarrow H_v$  is a bounded invertible operator. We shall show that there exists a positive constant  $c$  such that  $\| T_v x \| \leq c \| x \|$  for any  $x \in H_v$ . Take  $x = h((x_\alpha)_{\alpha \in E^v}) = \sum_{x \in E^v} f_\alpha(x_\alpha)$ . We get

$$\begin{aligned} \| T_v(x) \| &= \left\| \sum_{\alpha \in E^v} T_v(f_\alpha(x_\alpha)) \right\| = \left\| \sum_{x \in E^v} f_\alpha(T_{s(\alpha)}(x_\alpha)) \right\| \\ &= \left\| ((f_\alpha T_{s(\alpha)})_{\alpha \in E^v})((x_\alpha)_{\alpha \in E^v}) \right\| \leq \left\| ((f_\alpha T_{s(\alpha)})_{\alpha \in E^v}) \right\| \left\| ((x_\alpha)_{\alpha \in E^v}) \right\| \\ &= \left\| ((f_\alpha T_{s(\alpha)})_{\alpha \in E^v}) \right\| \| (h|_{K_v^\perp})^{-1} \| \| x \| \leq c \| x \| \end{aligned}$$

where  $((f_\alpha T_{s(\alpha)})_{\alpha \in E^v})$  is a row matrix and

$$c := \left\| ((f_\alpha T_{s(\alpha)})_{\alpha \in E^v}) \right\| \| (h|_{K_v^\perp})^{-1} \|.$$

Hence  $T_v$  is bounded. Next we shall show that  $\Phi_v^+(T) = S$ . Since  $S \in \text{End}(K, g)$ ,  $S_{s(\alpha)} P_\alpha i_v = S_{s(\alpha)} g_{\bar{\alpha}} = g_{\bar{\alpha}} S_v = P_\alpha i_v S_v$  for  $\alpha \in E^v$ . For  $((x_\alpha)_{\alpha \in E^v}) \in K_v$ , we have

$$S_v((x_\alpha)) = (P_\alpha i_v S_v((x_\alpha)))_{\alpha \in E^v} = (S_{s(\alpha)} P_\alpha i_v((x_\alpha)))_{\alpha \in E^v} = (S_{s(\alpha)}(x_\alpha)).$$

By the definition of  $\Phi_v^+(T)$ ,  $(\Phi_v^+(T))_u = S_u = T_u$  for  $u \neq v$ . For  $u = v$  and  $((x_\alpha)_{\alpha \in E^v}) \in K_v$ ,

$$\begin{aligned} (\Phi_v^+(T))_v((x_\alpha)_{\alpha \in E^v}) &= (T_{s(\alpha)}(x_\alpha))_{\alpha \in E^v} \\ &= ((S_{s(\alpha)}(x_\alpha))_{\alpha \in E^v}) = S_v((x_\alpha)_{\alpha \in E^v}). \end{aligned}$$

Thus  $(\Phi_v^+(T))_v = S_v$ . Hence  $\Phi_v^+(T) = S$ . Hence  $\Phi_v^+$  is onto. We conclude that  $\text{End}(H, f) \cong \text{End}(\Phi_v^+(H, f))$  as  $\mathbb{C}$ -algebras.  $\square$

**Corollary 4.20.** *Let  $\Gamma = (V, E, s, r)$  be a finite quiver and  $v \in V$  a sink. Assume that a Hilbert representation  $(H, f)$  of  $\Gamma$  is full at  $v$ . If  $(H, f)$  is transitive (resp. indecomposable), then  $\Phi_v^+(H, f)$  is transitive (resp. indecomposable).*

The following theorem is well known for finite-dimensional Hilbert spaces ([As, page289, 5.7. Corollary] and [DIR, page16, Proposition 2.1]).

**Theorem 4.21.** *Let  $\Gamma = (V, E, s, r)$  be a finite quiver and  $v \in V$  a source. If a Hilbert representation  $(H, f)$  of  $\Gamma$  is co-full at  $v$ . Then  $\Phi_v^- : \text{End}(H, f) \rightarrow \text{End}(\Phi_v^-(H, f))$  is an isomorphism as  $\mathbb{C}$ -algebras.*

**Proof.** We put  $(K, g) := \Phi_v^-(H, f)$ . The mapping  $\Phi_v^-$  gives a mapping of  $\text{End}(H, f)$  to  $\text{End}(K, g)$ . At first we shall show that  $\Phi_v^-$  is one to one. Assume that  $S := \Phi_v^-(T) = 0$  for  $T \in \text{End}(H, f)$ . We shall show that  $T_v = 0$ . Since  $T \in \text{End}(H, f)$ ,  $f_\alpha T_v = T_{r(\alpha)} f_\alpha$  for  $\alpha \in E_v$ . For  $(x_\alpha)_{\alpha \in E_v} \in \bigoplus_{\alpha \in E_v} H_{r(\alpha)}$ , we have

$$T_v^* \left( \sum f_\alpha^*(x_\alpha) \right) = \sum f_\alpha^*(T_{r(\alpha)}^*(x_\alpha)) = \sum f_\alpha^*(S_{r(\alpha)}^*(x_\alpha)) = 0.$$

Since  $(H, f)$  is co-full at  $v$ ,  $T_v^* = 0$ . Hence  $T_v = 0$ . Thus  $\Phi_v^-$  is one to one. Next we shall show that  $\Phi_v^-$  is onto. We put  $T_u = S_u$  for  $u \neq v$ . And we shall define an operator  $W_v : H_v \rightarrow H_v$  such that for  $(x_\alpha)_{\alpha \in E_v} \in \bigoplus_{\alpha \in E_v} H_{r(\alpha)}$ ,

$$W_v\left(\sum_{\alpha \in E_v} f_\alpha^*(x_\alpha)\right) = \sum_{\alpha \in E_v} f_\alpha^*(T_{r(\alpha)}^*(x_\alpha)).$$

We need to show that  $W_v$  is well defined. Assume that there exists an element  $(x'_\alpha)_{\alpha \in E_v} \in \bigoplus_{\alpha \in E_v} H_{r(\alpha)}$  such that

$$\sum_{\alpha \in E_v} f_\alpha^*(x_\alpha) = \sum_{\alpha \in E_v} f_\alpha^*(x'_\alpha).$$

We have

$$\hat{h}_v^*((x_\alpha - x'_\alpha)) = \sum_{\alpha \in E_v} f_\alpha^*(x_\alpha - x'_\alpha) = 0.$$

Hence  $(x_\alpha - x'_\alpha)_{\alpha \in E_v} \in \ker \hat{h}_v^* = K_v$ . Since  $S_v^* : K_v \rightarrow K_v$ , we have  $S_v^*((x_\alpha - x'_\alpha)_{\alpha \in E_v}) \in K_v$ . Hence  $\hat{h}_v^*(S_v^*((x_\alpha - x'_\alpha)_{\alpha \in E_v})) = 0$ . Since  $S \in \text{End}(K, g)$ , we have  $S_v g_\beta = g_\beta S_{r(\beta)}$  and  $g_\beta^* S_v^* = S_{r(\beta)}^* g_\beta^*$ . Hence

$$g_\beta^* S_v^*((x_\alpha - x'_\alpha)_{\alpha \in E_v}) = P_{r(\beta)} i_v S_v^*((x_\alpha - x'_\alpha)_{\alpha \in E_v})$$

and

$$S_{r(\beta)}^* g_\beta^*((x_\alpha - x'_\alpha)_{\alpha \in E_v}) = S_{r(\beta)}^*(x_\beta - x'_\beta).$$

Thus we have

$$P_{r(\beta)} i_v S_v^*((x_\alpha - x'_\alpha)_{\alpha \in E_v}) = S_{r(\beta)}^*(x_\beta - x'_\beta)$$

and

$$\sum f_\beta^* P_{r(\beta)} i_v (S_v^*((x_\alpha - x'_\alpha)_{\alpha \in E_v})) = \sum f_\beta^* S_{r(\beta)}^*(x_\beta - x'_\beta).$$

Since

$$\sum f_\beta^* P_{r(\beta)} i_v (S_v^*((x_\alpha - x'_\alpha)_{\alpha \in E_v})) = \hat{h}_v^*(S_v^*((x_\beta - x'_\beta)_{\beta \in E_v})) = 0,$$

we have

$$\sum f_\beta^* S_{r(\beta)}^*(x_\beta - x'_\beta) = \sum_{\beta \in E_v} f_\beta^* T_{r(\beta)}^*(x_\beta - x'_\beta) = 0.$$

Hence

$$\sum f_\beta^* T_{r(\beta)}^*(x_\beta) = \sum f_\beta^* T_{r(\beta)}^*(x'_\beta).$$

Thus  $W_v$  is well defined. Put  $T_v = W_v^*$ . Next we shall show that  $f_\alpha T_v = T_{s(\alpha)} f_\alpha$  and  $T_v^* f_\alpha^* = f_\alpha^* T_{s(\alpha)}^*$ . Take and fix  $x \in H_{r(\alpha)}$ . For  $\beta \in E_v$ , we put

$$x_\beta = \begin{cases} x & (\beta = \alpha), \\ 0 & (\beta \neq \alpha). \end{cases}$$

By the definition of  $W_v = T_v^*$ ,

$$W_v\left(\sum_{\alpha \in E^v} f_\alpha^*(x_\alpha)\right) = \sum_{\alpha \in E^v} f_\alpha^*(T_{r(\alpha)}^*(x_\alpha)).$$

Hence

$$T_v^* f_\alpha^*(x) = \sum_{\beta} f_\beta^* T_{r(\beta)}^*(x_\beta) = f_\alpha^* T_{r(\alpha)}^*(x) \text{ for } x \in H_{r(\alpha)}.$$

Thus we proved it.

Next we shall show that  $W_v = T_v^* : H_v \rightarrow H_v$  is bounded. By the Banach invertibility theorem,  $\hat{h}_v^*|_{(K_v)^\perp} : (K_v)^\perp \rightarrow H_v$  is a bounded invertible operator. We shall show that there exists a positive constant  $c$  such that  $\|T_v^* x\| \leq c \|x\|$  for any  $x \in H_v$ . For  $x \in H_v$ , there exists  $(x_\alpha)_{\alpha \in E_v} \in (K_v)^\perp$  such that  $x = \hat{h}_v^*((x_\alpha)_{\alpha \in E_v}) = \sum_{\alpha \in E_v} f_\alpha^*(x_\alpha)$ . We have

$$\begin{aligned} \|T_v^*(x)\| &= \left\| \sum_{\alpha \in E_v} T_v^*(f_\alpha^*(x_\alpha)) \right\| = \left\| \sum_{\alpha \in E_v} f_\alpha^*(T_{r(\alpha)}^*(x_\alpha)) \right\| \\ &= \left\| (f_\alpha^* T_{r(\alpha)}^*)_{\alpha \in E_v} (x_\alpha)_{\alpha \in E_v} \right\| \leq \left\| (f_\alpha^* T_{r(\alpha)}^*)_{\alpha \in E_v} \right\| \left\| (x_\alpha)_{\alpha \in E_v} \right\| \\ &= \left\| (f_\alpha^* T_{r(\alpha)}^*)_{\alpha \in E_v} \right\| \left\| (\hat{h}_v^*|_{K_v^\perp})^{-1} \right\| \|x\| \leq c \|x\|, \end{aligned}$$

where  $(f_\alpha^* T_{r(\alpha)}^*)_{\alpha \in E_v}$  is a row matrix and

$$c := \left\| ((f_\alpha^* T_{r(\alpha)}^*)_{\alpha \in E_v}) \right\| \left\| (\hat{h}_v^*|_{K_v^\perp})^{-1} \right\|.$$

Hence  $T_v$  is bounded. Next we shall show that  $\Phi_v^-(T) = S$ . By the definition of  $\Phi_v^-(T)$ ,  $(\Phi_v^-(T))_u = S_u = T_u$  for  $u (\neq v) \in V$ . Since  $S \in \text{End}(K, g)$ , we have

$$S_v Q_v j_\beta = S_v g_{\bar{\beta}} = g_{\bar{\beta}} S_{r(\beta)} = Q_v j_\beta S_{r(\beta)}$$

for  $\beta \in E_v$ . For  $(x_\beta)_{\beta \in E_v} \in K_v$ , we have

$$\begin{aligned} S_v((x_\beta)_{\beta \in E_v}) &= S_v Q_v \left( \sum_{\beta \in E_v} j_\beta(x_\beta) \right) = \sum_{\beta \in E_v} S_v Q_v j_\beta(x_\beta) \\ &= \sum_{\beta \in E_v} Q_v j_\beta(S_{r(\beta)} x_\beta) = Q_v \sum_{\beta \in E_v} j_\beta(S_{r(\beta)} x_\beta) \\ &= Q_v((S_{r(\beta)} x_\beta)_{\beta \in E_v}). \end{aligned}$$

Thus

$$S_v((x_\beta)_{\beta \in E_v}) = Q_v((S_{r(\beta)} x_\beta)_{\beta \in E_v}).$$

For  $u = v$  and  $((x_\alpha)_{\alpha \in E_v}) \in K_v$ ,

$$\begin{aligned} (\Phi_v^-(T))_v((x_\alpha)_{\alpha \in E_v}) &= Q_v((T_{r(\alpha)} x_\alpha)_{\alpha \in E_v}) = Q_v((S_{r(\alpha)} x_\alpha)_{\alpha \in E_v}) \\ &= S_v((x_\alpha)_{\alpha \in E_v}). \end{aligned}$$

Thus  $(\Phi_v^-(T))_v = S_v$ . Hence  $\Phi_v^-(T) = S$  and

$$\Phi_v^- : \text{End}(H, f) \rightarrow \text{End}(\Phi_v^-(H, f))$$

is onto. Thus we have  $\text{End}(H, f) \cong \text{End}(\Phi_v^-(H, f))$  as  $\mathbb{C}$ -algebras.  $\square$

**Corollary 4.22.** *Let  $\Gamma = (V, E, s, r)$  be a finite quiver and  $v \in V$  a source. Assume that a Hilbert representation  $(H, f)$  of  $\Gamma$  is co-full at  $v$ . If  $(H, f)$  is transitive, then  $\Phi_v^-(H, f)$  is transitive. Similarly, if  $(H, f)$  is indecomposable, then  $\Phi_v^{-1}(H, f)$  is indecomposable.*

Next, we shall show the existence of infinite-dimensional transitive Hilbert representations of quivers with any orientation whose underlying undirected graphs are extended Dynkin diagrams  $\tilde{D}_n$  ( $n \geq 4$ ),  $\tilde{E}_6$ ,  $\tilde{E}_7$  and  $\tilde{E}_8$ .

We recall some definitions and lemmas in [EW3].

**Definition.**[EW3] Let  $\Gamma$  be a quiver whose underlying undirected graph is Dynkin diagram  $A_n$ . We count the arrows from the left as  $\alpha_k : s(\alpha_k) \rightarrow r(\alpha_k)$ , ( $k = 1, \dots, n - 1$ ). Let  $(H, f)$  be a Hilbert representation of  $\Gamma$ . We denote  $f_{\alpha_k}$  briefly by  $f_k$ . For example,

$$\circ_{H_1} \xrightarrow{f_1} \circ_{H_2} \xrightarrow{f_2} \circ_{H_3} \xleftarrow{f_3} \circ_{H_4} \xrightarrow{f_4} \circ_{H_5} \xleftarrow{f_5} \circ_{H_6}$$

It is said that  $(H, f)$  is *positive-unitary diagonal* if there exist  $m \in \mathbb{N}$  and orthogonal decompositions (admitting zero components) of Hilbert spaces

$$H_k = \oplus_{i=1}^m H_{k,i} \quad (k = 1, \dots, n)$$

and decompositions of operators

$$f_k = \oplus_{i=1}^m f_{k,i} : \oplus_{i=1}^m H_{s(\alpha_k),i} \rightarrow \oplus_{i=1}^m H_{r(\alpha_k),i} \quad (k = 1, \dots, n)$$

such that each  $f_{k,i} : H_{s(\alpha_k),i} \rightarrow H_{r(\alpha_k),i}$  is written as  $f_{k,i} = 0$  or  $f_{k,i} = \lambda_{k,i} u_{k,i}$  for some positive scalar  $\lambda_{k,i}$  and onto unitary  $u_{k,i} \in B(H_{s(\alpha_k),i}, H_{r(\alpha_k),i})$ .

It is easily seen that if  $(H, f)$  is positive-unitary diagonal, then  $\Phi^*(H, f)$  is also positive-unitary diagonal.

**Lemma 4.23.** [EW3, Lemma 6.4.] *Let  $\Gamma$  be a quiver whose underlying undirected graph is Dynkin diagram  $A_n$  and  $(H, f)$  be a Hilbert representation of  $\Gamma$ . Suppose that  $(H, f)$  is positive-unitary diagonal. Then  $(H, f)$  is closed at any sink of  $\Gamma$  and co-closed at any source of  $\Gamma$ .*

**Proposition 4.24.** [EW3, Proposition 6.5.] *Let  $\Gamma$  be a quiver whose underlying undirected graph is Dynkin diagram  $A_n$  and  $(H, f)$  be a Hilbert representation of  $\Gamma$ . Let  $v$  be a source of  $\Gamma$ . Suppose that  $(H, f)$  is positive-unitary diagonal. Then  $\Phi_v^-(H, f)$  is also positive-unitary diagonal.*

It is known that every orientation of Dynkin diagram  $A_n$  is obtained by an iteration of  $\sigma_v^-$  at sources  $v$  except the right end from a particular orientation as follows:

**Lemma 4.25.** [EW3, Lemma 6.6.] *Let  $\Gamma_0$  and  $\Gamma$  be quivers whose underlying undirected graphs are the same Dynkin diagram  $A_n$  for  $n \geq 2$ . Assume that  $\Gamma_0$  is the following:*

$$\circ_1 \longrightarrow \circ_2 \longrightarrow \circ_3 \cdots \circ_{n-1} \longrightarrow \circ_n$$

*Then there exists a sequence  $v_1, \dots, v_m$  of vertices in  $\Gamma_0$  such that*

- (1) *for each  $k = 1, \dots, m$ ,  $v_k$  is a source in  $\sigma_{v_{k-1}}^- \cdots \sigma_{v_2}^- \sigma_{v_1}^- (\Gamma_0)$ ,*
- (2)  *$\sigma_{v_m}^- \cdots \sigma_{v_2}^- \sigma_{v_1}^- (\Gamma_0) = \Gamma$ ,*
- (3) *for each  $k = 1, \dots, m$ ,  $v_k \neq n$ .*

**Lemma 4.26.** [EW3, Lemma 5.6.] *Let  $\Gamma = (V, E, s, r)$  be a finite quiver and  $v \in V$  a sink. Then for any Hilbert representation  $(H, f)$  of  $\Gamma$ ,  $\Phi_v^+(H, f)$  is co-full at  $v$ .*

**Theorem 4.27.** [EW3, Theorem 5.13.] *Let  $\Gamma = (V, E, s, r)$  be a finite quiver and  $v \in V$  a source. Assume that a Hilbert representation  $(H, f)$  of  $\Gamma$  is indecomposable and co-closed at  $v$ . Then the following assertions hold:*

- (1) *If  $\Phi_v^-(H, f) = 0$ , then  $H_v = \mathbb{C}$ ,  $H_u = 0$  for any  $u \in V$  with  $u \neq v$  and  $f_\alpha = 0$  for any  $\alpha \in E$ .*
- (2) *If  $\Phi_v^-(H, f) \neq 0$ , then  $\Phi_v^-(H, f)$  is also indecomposable and  $(H, f) \cong \Phi_v^+ \Phi_v^-(H, f)$ .*

The following is one of the main theorems in this paper.

**Theorem 4.28.** *Let  $\Gamma$  be a quiver whose underlying undirected graph is an extended Dynkin diagram. Then there exists an infinite-dimensional transitive Hilbert representation of  $\Gamma$  if and only if  $\Gamma$  is not an oriented cyclic quiver.*

**Proof.** Suppose that  $\Gamma$  is an oriented cyclic quiver. Theorem 4.10 proves the nonexistence of infinite-dimensional transitive Hilbert representation of  $\Gamma$ . Suppose that  $\Gamma$  is not an oriented cyclic quiver. We shall prove the existence of infinite-dimensional transitive Hilbert representations of  $\Gamma$ . When  $\widetilde{A}_n$  case, Theorem 4.10 proves the existence of infinite-dimensional transitive Hilbert representations of  $\Gamma$ . Next we consider the case that the  $|\Gamma|$  is  $\widetilde{D}_n$ . Let  $\Gamma_0$  be the quiver of Lemma 4.11 and  $(H^{(0)}, f^{(0)})$  the Hilbert representation constructed there. Then  $|\Gamma_0| = |\Gamma| = \widetilde{D}_n$ , but their orientations are different in general. Let  $\Gamma_1$  be a quiver such that  $|\Gamma_1| = \widetilde{D}_n$  and the orientation is as same as  $\Gamma$  on the path between 5 and  $n+1$  and as same as  $\Gamma_0$  on the rest four “wings”. We shall define a Hilbert representation  $(H^{(1)}, f^{(1)})$  of  $\Gamma_1$  modifying  $(H^{(0)}, f^{(0)})$ . We put  $f_\beta^{(1)} = I$  for any arrow  $\beta$  in the path between 5 and  $n+1$ . and  $f_\beta^{(0)} = f_\beta^{(1)}$  for other arrow  $\beta$ . The same proof for  $(H^{(0)}, f^{(0)})$  shows that  $(H^{(1)}, f^{(1)})$  is transitive. Since  $f_{\alpha_i}^{(1)}$  ( $i = 1, \dots, 4$ ) is an inclusion map,  $(H^{(1)}, f^{(1)})$  is co-full at sources 1,2,3 and 4. By Theorem 4.21, a certain iteration of reflection functors at a source 1,2,3 or 4 on  $(H^{(1)}, f^{(1)})$  gives an infinite-dimensional, transitive, Hilbert representation of  $\Gamma$ . We have proved this case.

Next we consider the case that the  $|\Gamma|$  is  $\widetilde{E}_6$ . Let  $\Gamma_0$  be the quiver of Lemma 4.12, and we denote here by  $(H^{(0)}, f^{(0)})$  the Hilbert representation constructed there. Then  $|\Gamma_0| = |\Gamma| = \widetilde{E}_6$ , but their orientations are different in general. Three “wings” of  $|\Gamma_0|$   $2-1-0$ ,  $2'-1'-0$ ,  $2''-1''-0$  can be regarded as Dynkin diagrams  $A_3$ . Applying Lemma 4.25 for these wings locally, we can find a sequence  $v_1, \dots, v_m$  of vertices in  $\Gamma_0$  such that

- (1) for each  $k = 1, \dots, m$ ,  $v_k$  is a source in  $\sigma_{v_{k-1}}^- \dots \sigma_{v_2}^- \sigma_{v_1}^- (\Gamma_0)$ ,
- (2)  $\sigma_{v_m}^- \dots \sigma_{v_2}^- \sigma_{v_1}^- (\Gamma_0) = \Gamma$ ,

(3) for each  $k = 1, \dots, m, v_k \neq 0$ .

We note that co-closedness of Hilbert representations at a source can be checked locally around the source. Since the restriction of the representation  $(H^{(0)}, f^{(0)})$  to each “wing” is positive-unitary diagonal and the iteration of reflection functors does not move the vertex 0, we can apply Lemma 4.23 and Proposition 4.24 locally that  $\Phi_{v_{k-1}}^- \dots \Phi_{v_2}^- \Phi_{v_1}^-(H^{(0)}, f^{(0)})$  is co-closed at  $v_k$  for  $k = 1, \dots, m$ . Since the particular Hilbert space  $H_0^{(0)}$  associated with the vertex 0 is infinite-dimensional and remains unchanged under the iteration of the reflection functors above,  $\Phi_{v_i}^- \dots \Phi_{v_1}^-(H^{(0)}, f^{(0)})$  ( $1 \leq i \leq m$ ) is infinite-dimensional. Therefore Theorem 4.27 implies that

$$\Phi_{v_i}^- \dots \Phi_{v_1}^-(H^{(0)}, f^{(0)}) \quad (1 \leq i \leq m)$$

is an infinite-dimensional indecomposable Hilbert representation of

$$\sigma_{v_i}^- \dots \sigma_{v_2}^- \sigma_{v_1}^-(\Gamma).$$

By Theorem 4.27, for

$$(K, g) := \Phi_{v_i}^- \dots \Phi_{v_1}^-(H^{(0)}, f^{(0)}) \quad (1 \leq i \leq m),$$

we have

$$(K, g) \cong \Phi_{v_{i+1}}^+ \Phi_{v_{i+1}}^-(K, g).$$

On the other hand, by Lemma 4.26,  $\Phi_{v_{i+1}}^+ \Phi_{v_{i+1}}^-(K, g)$  is co-full at  $v_{i+1}$ . Since  $(K, g) \cong \Phi_{v_{i+1}}^+ \Phi_{v_{i+1}}^-(K, g)$ , by Lemma 4.18, we have that  $(K, g)$  is co-full at  $v_{i+1}$ . Hence Theorem 4.21 implies that  $End(K, g) \cong End(\Phi_{v_{i+1}}^-(K, g))$ . By induction, we have

$$End(H^{(0)}, f^{(0)}) \cong End(\Phi_{v_m}^- \dots \Phi_{v_1}^-(H^{(0)}, f^{(0)})).$$

Since  $(H^{(0)}, f^{(0)})$  is transitive,  $(\Phi_{v_m}^- \dots \Phi_{v_1}^-(H^{(0)}, f^{(0)}))$  is also transitive. Thus there exist infinite-dimensional transitive Hilbert representations for quivers with any orientation whose underlying undirected graphs is extended Dynkin diagram  $\widetilde{E}_6$ . The other cases  $\widetilde{E}_7$  and  $\widetilde{E}_8$  are proved similarly.  $\square$

### References

- [Ak] AKHIEZER, NAUM I.; GLAZMAN, IZRAIL' M. Theory of linear operators in Hilbert space. I. *Frederick Ungar Publishing Co., New York*, 1961. xi+147 pp. MR0264420, Zbl 0467.47001. 982
- [Ar] ARONSZAJN, NACHMAN; FIXMAN, URI. Algebraic spectral problems. *Studia Math.* **30** (1968), 273-338. MR0240114, Zbl 0165.15202, doi:10.4064/sm-30-3-273-338. 976
- [As] ASSEM, IBRAHIM; SIMSON, DANIEL; SKOWROŃSKI, ANDRZEJ. Elements of the representation theory of associative algebras. 1. Techniques of representation theory. London Mathematical Society Student Texts, 65 *Cambridge University Press, Cambridge*, 2006. x+458 pp. ISBN: 978-0-521-58423-4; 978-0-521-58631-3. MR2197389, Zbl 1092.16001, doi:10.1017/CBO9780511614309. 977, 980, 1002, 1006, 1008

- [Au] AUSLANDER, MAURICE. Large modules over Artin algebras. *Algebra, topology and category theory (a collection of papers in honor of Samuel Eilenberg)*, 1–17. Academic Press New York, 1976. MR0424874, Zbl 0442.16025, doi:10.1016/B978-0-12-339050-9.50006-7. 976
- [BGP] BERNŠTEĪN, JOESPH N.; GEL'FAND, ISRAIL' M.; PONOMAREV, V.A. Coxeter functors and Gabriel's theorem, *Russian Math. Surveys* **28** (1973), no. 2, 17–32. MR0393065, Zbl 0279.08001, doi:10.1070/RM1973v028n02ABEH001526. 976, 1002
- [DeZ1] DEAN, ANDREW P.; ZORZITTO, FRANK A. A criterion for pure simplicity. *J. Algebra* **132** (1990), no. 1, 50–71. MR1060831, Zbl 0703.16005, doi:10.1016/0021-8693(90)90251-I. 976
- [DeZ2] DEAN, ANDREW P.; ZORZITTO, FRANK A. Infinite dimensional representations of  $\tilde{D}_4$ . *Glasgow Math. J.* **32** (1990), no. 1, 25–33. MR1045083, Zbl 0689.16015, doi:10.1017/S0017089500009034. 976
- [DIR] DLAB, VLASTIMIL; RINGEL, CLAUS MICHAEL. Indecomposable representations of graphs and algebras. *Mem. Amer. Math. Soc.* **6** (1976), no. 173, v+57 pp. MR0447344, Zbl 0332.16015, doi:10.1090/memo/0173. 976, 1002, 1006, 1008
- [DoF] DONOVAN PETER; FREISLICH MARY RUTH. The representation theory of finite graphs and associated algebras. Carleton Mathematical Lecture Notes, 5. *Carleton University, Ottawa, Ont.*, 1973. iii+83 pp. MR0357233, Zbl 0304.08006. 976, 1002
- [E] ENOMOTO, MASATOSHI. A construction on Hilbert representations of quivers. *RIMS Kokyuroku* **1893** (2014), 102–114. 976
- [EW1] ENOMOTO, MASATOSHI; WATATANI, YASUO. Relative position of four subspaces in a Hilbert space. *Adv. Math.* **201** (2006), no. 2, 263–317. MR2211531, Zbl 1095.46020, arXiv:math/0404545, doi:10.1016/j.aim.2005.02.004. 976, 980
- [EW2] ENOMOTO, MASATOSHI; WATATANI, YASUO. Exotic indecomposable systems of four subspaces in a Hilbert space. *Integral Equations Operator Theory* **59** (2007), no. 2, 149–164. MR2345993, Zbl 1136.46016, arXiv:math/0607085, doi:10.1007/s00020-007-1512-2. 976
- [EW3] ENOMOTO, MASATOSHI; WATATANI, YASUO. Indecomposable representations of quivers on infinite-dimensional Hilbert spaces. *J. Funct. Anal.* **256** (2009), no. 4, 959–991. MR2488332, Zbl 1168.46010, arXiv:0707.0966, doi:10.1016/j.jfa.2008.12.011. 976, 979, 999, 1000, 1001, 1002, 1003, 1004, 1005, 1011, 1012
- [EW4] ENOMOTO, MASATOSHI; WATATANI, YASUO. Strongly irreducible operators and indecomposable representations of quivers on infinite-dimensional Hilbert spaces. *Integral Equations Operator Theory* **83** (2015), no. 4, 563–587. MR3430212, Zbl 1372.47028, arXiv:1303.2485, doi:10.1007/s00020-015-2228-3. 977, 979, 981, 986, 988, 989
- [Fi] FIXMAN, URI. On algebraic equivalence between pairs of linear transformations. *Trans. Amer. Math. Soc.* **113** (1964), 424–453. MR0169855, Zbl 0192.13105, doi:10.2307/1994142. 976
- [FiO] FIXMAN, URI; OKOH, FRANK. Extensions of pairs of linear transformations between infinite dimensional vector spaces. *Linear Algebra and Appl.* **19** (1978), no. 3, 275–291. MR0498624, Zbl 0381.15003, doi:10.1016/0024-3795(78)90015-0. 976
- [FiZ] FIXMAN, URI; ZORZITTO, FRANK A. A purity criterion for pairs of linear transformations. *Canadian J. Math.* **26** (1974), 734–745. MR0353054, Zbl 0255.15003, doi:10.4153/CJM-1974-068-8. 976
- [FiW] FILLMORE, PETER A.; WILLIAMS, JAMES P. On operator ranges. *Adv. Math.* **7** (1971), 254–281. MR0293441, Zbl 0224.47009, doi:10.1016/S0001-8708(71)80006-3. 977



- [Ga] GABRIEL, PETER. Unzerlegbare Darstellungen. I. *Manuscripta Math.* **6** (1972), 71–103; correction, *ibid.* **6** (1972) 309. MR0332887, Zbl 0232.08001, doi:10.1007/BF01298413. 976
- [GaR] GABRIEL, PETER; ROITER, ANDREI V. Representations of finite-dimensional algebras. *Springer-Verlag, Berlin*, 1997. iv+177 pp. ISBN: 3-540-62990-4. MR1475926. 976, 1002
- [GeP] GEL'FAND, ISRAIL' MOISEEVICH; PONOMAREV, V. A. Problems of linear algebra and classification of quadruples of subspaces in a finite-dimensional vector space. *Hilbert space operators and operator algebras* (Proc. Internat. Conf., Tihany, 1970), 163–237, Colloq. Math. Soc. János Bolyai, 5. *North-Holland, Amsterdam*, 1972. MR0357428, Zbl 0294.15002. 976, 1002
- [Gi] GILFEATHER, FRANK. Strong reducibility of operators. *Indiana Univ. Math. J.* **22** (1972/73), 393–397. MR0303322, Zbl 0277.47018, doi:10.1512/iumj.1972.22.22032. 976
- [GoHJ] GOODMAN, FREDERICK M.; DE LA HARPE, PIERRE; JONES, VAUGHAN F. R. Coxeter graphs and towers of algebras. Mathematical Sciences Research Institute Publications, 14, *Springer-Verlag, New York*, 1989. x+288 pp. ISBN: 0-387-96979-9. MR0999799, Zbl 0698.46050, doi:10.1007/978-1-4613-9641-3. 976
- [H] HALMOS, PAUL R. Ten problems in Hilbert space. *Bull. Amer. Math. Soc.* **76** (1970), 887–933. MR0270173, Zbl 0204.15001, doi:10.1090/S0002-9904-1970-12502-2. 977, 980, 981
- [HRR] HARRISON, KENNETH J.; RADJAVI, HEYDAR; ROSENTHAL, PETER. A transitive medial subspace lattice. *Proc. Amer. Math. Soc.* **28** (1971), 119–121. MR0283609, Zbl 0211.44402, doi:10.2307/2037768. 977, 978, 980, 981, 1000
- [JiW1] JIANG, CHUNLAN; WANG, ZONGYAO. Strongly irreducible operators on Hilbert space. Pitman Research Notes in Mathematics Series, 389. *Longman, Harlow*, 1998. x+243 pp. ISBN: 0-582-30594-2. MR1640067, Zbl 0946.47011. 976
- [JiW2] JIANG, CHUNLAN; WANG, ZONGYAO. Structure of Hilbert space operators. *World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ*, 2006. x+248 pp. ISBN: 981-256-616-3. MR2221863, Zbl 1102.47001, doi:10.1142/5993. 976
- [Jo] JONES, VAUGHAN F.R. Index for subfactors. *Invent. Math.* **72** (1983), no. 1, 1–25. MR0696688, Zbl 0508.46040, doi:10.1007/BF01389127. 976
- [Ka] KAC, VICTOR G. Infinite root systems, representations of graphs and invariant theory. *Invent. Math.* **56** (1980), no. 1, 57–92. MR0557581, Zbl 0427.17001, doi:10.1007/BF01403155. 976
- [Kau] KAUFMAN, WILLIAM E. Representing a closed operator as a quotient of continuous operators. *Proc. Amer. Math. Soc.* **72** (1978), no. 3, 531–534. MR0509249, Zbl 0404.47001, doi:10.2307/2042466. 977, 984
- [Ko] KOSAKI, HIDEKI. On intersections of domains of unbounded positive operators. *Kyushu J. Math.* **60**, (2006), no. 1, 3–25. MR2216946, Zbl 1102.47014, doi:10.2206/kyushujm.60.3. 977
- [KrR] KRAUSE, HENNING; RINGEL, CLAUS MICHAEL; DIRS. Infinite length modules. Invited lectures from the conference held in Bielefeld, September 7–11, 1998. Trends in Mathematics. *Birkhäuser Verlag, Basel*, 2000. x+439 pp. ISBN: 3-7643-6413-0. MR1789208, Zbl 0945.00021. 976
- [Kro] KRONECKER, L. Algebraische Reduktion der Scharen bilinearer Formen. *Sitzungsber. Akad. Berlin* (1890), 763–776. JFM 22.0169.01. 976
- [KrRS] KRUGLYAK, STANISLAV A.; RABANOVICH, VIACHESLAV I.; SAMOÏLENKO, YURIÏ S. On sums of projections. *Funct. Anal. Appl.* **36** (2002), no. 3, 182–195. MR1935900, Zbl 1038.47001, doi:10.1023/A:1020193804109. 977
- [KrRo] KRUGLYAK, STANISLAV A.; ROITER, ANDREI V. Locally scalar representations of graphs in the category of Hilbert spaces. *Funct. Anal. Appl.* **39** (2005), no. 2,

- 91–105. MR2161513, Zbl 1175.16011, arXiv:math/0307163, doi:10.1007/s10688-005-0022-8. 977
- [MS] MOSKALEVA, YULIYA P. ; SAMOÏLENKO, YURIÏ S. Systems of  $n$  subspaces and representations of  $*$ -algebras generated by projections. *Methods Funct. Anal. Topology* **12** (2006), no. 1, 57–73. MR2210905, Zbl 1092.46037, arXiv:math/0603503. 977
- [Na1] NAZAROVA, LIUDMILA A. Representation of a tetrad. *Izv. Akad. Nauk SSSR Ser. Mat.* **31** (1967), 1361–1378, translation in *Math. USSR Izv.* **1** (1967), 1305–1321. MR0223352, Zbl 0222.16028, doi:10.1070/IM1967v001n06ABEH000619. 976
- [Na2] NAZAROVA, LIUDMILA A. Representations of quivers of infinite type. *Izv. Akad. Nauk SSSR, Ser. Mat.* **37** (1973), 752–791. MR0338018, Zbl 0298.15012, doi:10.1070/IM1973v007n04ABEH001975. 976
- [Ok] OKOH, FRANK. Applications of linear functionals to Kronecker modules. I, II. *Linear Algebra Appl.* **76** (1986), 165–204. MR0830339, Zbl 0591.16011, doi:10.1016/0024-3795(86)90221-1, Zbl 0591.16012, doi:10.1016/0024-3795(86)90222-3. 976
- [RR] RADJAVI, HEYDAR; ROSENTHAL, PETER. Invariant subspaces. *Ergebnisse der Mathematik und ihrer Grenzgebiete, 77 Springer-Verlag, New York-Heidelberg*, 1973. xi+219 pp. ISBN: 978-3-642-65576-0. MR0367682, Zbl 0269.47003, doi:10.1007/978-3-642-65574-6. 978, 981
- [Ri1] RINGEL, CLAU M. Infinite dimensional representations of finite dimensional hereditary algebras. *Symposia Mathematica, XXIII* (Conf. Abelian Groups and their Relationship to the Theory of Modules, INDAM, Rome, 1977), 321–412. *Academic Press, London-New York, 1979*. MR0565613, Zbl 0429.16022. 976
- [Ri2] RINGEL, CLAU M. The rational invariants of the tame quivers. *Invent. Math.* **58** (1980), no. 3, 217–239. MR0571574, Zbl 0433.15009, doi:10.1007/BF01390253. 976
- [Ri3] RINGEL, CLAU M. Infinite length modules. Some examples as introduction. *Infinite Length Modules* (Bielefeld, 1998), 1–73, Trends Math. *Birkhäuser, Basel*, 2000. MR1789209, Zbl 0988.16002, doi:10.1007/978-3-0348-8426-6\_1. 976
- [Sh] SHIELDS, ALLEN L. Weighted shift operators and analytic function theory. *Topics in operator theory*, 49–128. Math. Surveys, 13. *Amer. Math. Soc., Providence, R.I.*, 1974. MR0361899, Zbl 0303.47021, doi:10.1090/surv/013/02. 987

(Masatoshi Enomoto) INSTITUTE OF EDUCATION AND RESEARCH, KOSHIEI UNIVERSITY,  
TAKARAZUKA, HYOGO 665-0006, JAPAN  
enomotoma@hotmail.co.jp

(Yasuo Watatani) DEPARTMENT OF MATHEMATICAL SCIENCES, KYUSHU UNIVERSITY,  
MOTOOKA, FUKUOKA, 819-0395, JAPAN  
watatani@math.kyushu-u.ac.jp

This paper is available via <http://nyjm.albany.edu/j/2019/25-43.html>.