

Sharp bound for embedded eigenvalues of Dirac operators with decaying potentials

Vishwam Khapre, Kang Lyu and Andrew Yu

ABSTRACT. We study eigenvalues of the Dirac operator with canonical form

$$L_{p,q} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} -p & q \\ q & p \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix},$$

where p and q are real functions. Under the assumption that

$$\limsup_{x \rightarrow \infty} x \sqrt{p^2(x) + q^2(x)} < \infty,$$

the essential spectrum of $L_{p,q}$ is $(-\infty, \infty)$. We prove that $L_{p,q}$ has no eigenvalues if

$$\limsup_{x \rightarrow \infty} x \sqrt{p^2(x) + q^2(x)} < \frac{1}{2}.$$

Given any $A \geq \frac{1}{2}$ and any $\lambda \in \mathbb{R}$, we construct functions p and q such that $\limsup_{x \rightarrow \infty} x \sqrt{p^2(x) + q^2(x)} = A$ and λ is an eigenvalue of the corresponding Dirac operator $L_{p,q}$. We also construct functions p and q so that the corresponding Dirac operator $L_{p,q}$ has any prescribed set (finitely or countably many) of eigenvalues.

CONTENTS

1. Introduction and main results	1317
2. Proof of Theorems 1.1 and 1.2	1320
3. Proof of Theorems 1.3 and 1.4	1324
Acknowledgments	1327
References	1327

1. Introduction and main results

The Schrödinger operator given by

$$Hu = -u'' + Vu \tag{1}$$

Received June 8, 2022.

Key words and phrases. Dirac operators, canonical form, embedded eigenvalues, essential spectrum.

2020 Mathematics Subject Classification. Primary: 34L15. Secondary: 34A30.

and the Dirac operator given by

$$L \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \frac{d}{dx} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} \quad (2)$$

are two basic models in mathematics and physics. We are interested in the embedded eigenvalue (eigenvalue embeds into the essential spectrum) problem of Schrödinger operators and Dirac operators. For Schrödinger operators, the problem is well understood. Kato's classical results [9] show that if

$$\limsup_{x \rightarrow \infty} |xV(x)| = A,$$

then the Schrödinger operator has no eigenvalues larger than A^2 . Wigner and von Neumann's examples [25] imply that there exist potentials with $A = 8$, such that $\lambda = 1$ is an eigenvalue of the associated Schrödinger operator. Finally, (see the survey [23] for the history), Atkinson and Everitt [1] obtained the sharp bound $\frac{4A^2}{\pi^2}$. They proved that there are no eigenvalues larger than $\frac{4A^2}{\pi^2}$, and for any $0 < \lambda < \frac{4A^2}{\pi^2}$, there are potentials with $\limsup_{x \rightarrow \infty} |xV(x)| = A$ so that λ is an eigenvalue of the associated Schrödinger operator.

Equations (1) and (2) are closely related. For example, by letting $p_{11} = V$, and $p_{12} = p_{21} = p_{22} = 0$, one can obtain

$$-u'' + \lambda V u = \lambda^2 u$$

from

$$L \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix}.$$

In this article, we study embedded eigenvalue problems of a particular type of Dirac operators on $L^2[0, \infty) \oplus L^2[0, \infty)$, namely Dirac operators with canonical form,

$$L_{p,q} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \frac{d}{dx} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} -p & q \\ q & p \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}, \quad (3)$$

where $p \in L^2[0, \infty)$ and $q \in L^2[0, \infty)$ are real functions (referred to as potentials). The canonical form of Dirac operators plays an important role in spectral theory [19, Theorem 5.1]. In the study of asymptotics of eigenvalues and the inverse problems of Dirac operators, it is crucial to use the canonical form [10, pp. 185-187], [27, 28]. We refer readers to [3, 4, 5, 6, 7, 20] for more recent development about various types of Dirac operators.

For any $\phi_0 \in [0, \pi)$, under the boundary condition

$$u(0) \sin \phi_0 - v(0) \cos \phi_0 = 0, \quad (4)$$

the Dirac operator $L_{p,q}$ defined by (3) is self-adjoint.

Denote by $\sigma_{\text{ess}}(L_{p,q})$ the essential spectrum of $L_{p,q}$. Recall that $\lambda \in \sigma_{\text{ess}}(L_{p,q})$ if and only if there is an orthonormal sequence $\{\varphi_n\}_{n=1}^{\infty}$ such that

$$\|L_{p,q}\varphi_n - \lambda\varphi_n\| \rightarrow 0, \quad n \rightarrow \infty.$$

It is well known that

$$\sigma_{\text{ess}}(L_{0,0}) = (-\infty, \infty),$$

and $L_{0,0}$ has no eigenvalues.

By [26, Theorem 6.4], if

$$\sqrt{p^2(x) + q^2(x)} = o(1),$$

as $x \rightarrow \infty$, then

$$\sigma_{\text{ess}}(L_{p,q}) = (-\infty, \infty).$$

In the first part of our paper, under the assumption that p and q are Coulomb type potentials (but without singularity at $x = 0$), we study the question when $L_{p,q}$ has embedded eigenvalues.

Theorem 1.1. *If*

$$\limsup_{x \rightarrow \infty} x \sqrt{p(x)^2 + q(x)^2} = A < \frac{1}{2},$$

then under any boundary condition (4), $L_{p,q}$ has no eigenvalues in $(-\infty, \infty)$.

Theorem 1.2. *For any $\phi_0 \in [0, \pi)$, $\lambda \in (-\infty, \infty)$, and $A \geq \frac{1}{2}$, there exist potentials p and q such that*

$$\limsup_{x \rightarrow \infty} x \sqrt{p(x)^2 + q(x)^2} = A,$$

and the Dirac operator $L_{p,q}$ has an eigenvalue λ under the boundary condition (4).

We say that the potential is C^∞ if p, q are C^∞ . In the second part of the paper, we will construct C^∞ potentials with which $L_{p,q}$ has many embedded eigenvalues.

Theorem 1.3. *Let $S = \{\lambda_j\}_{j=1}^N$ be a set of distinct real numbers. Let $\{\theta_j\}_{j=1}^N \subset [0, \pi)$ be a set of angles. There exist C^∞ potentials satisfying*

$$\sqrt{p(x)^2 + q(x)^2} = \frac{O(1)}{1+x},$$

where $O(1)$ depends on S , such that the associated $L_{p,q}$ has $L^2[0, \infty) \oplus L^2[0, \infty)$ solutions $(u_j, v_j)^T$ satisfying

$$L_{p,q} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda_j \begin{pmatrix} u \\ v \end{pmatrix}$$

with the boundary condition

$$\frac{u(0)}{v(0)} = \cot \theta_j,$$

for $j = 1, \dots, N$.

Theorem 1.4. *Let $S = \{\lambda_j\}_{j=1}^\infty$ be a set of distinct real numbers. Let $\{\theta_j\}_{j=1}^\infty \subset [0, \pi)$ be a set of angles. If $h(x)$ is a positive function with $\lim_{x \rightarrow \infty} h(x) = \infty$, then there exist C^∞ potentials satisfying*

$$\sqrt{p(x)^2 + q(x)^2} \leq \frac{h(x)}{1+x},$$

such that the associated Dirac operator $L_{p,q}$ has $L^2[0, \infty) \oplus L^2[0, \infty)$ solutions $(u_j, v_j)^T$ satisfying

$$L_{p,q} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda_j \begin{pmatrix} u \\ v \end{pmatrix}$$

with the boundary condition

$$\frac{u(0)}{v(0)} = \cot \theta_j,$$

for $j = 1, 2, \dots$.

For Dirac operators with single embedded eigenvalue, Evans and Harris [2] obtained the sharp bound for the separated Dirac equation with the form

$$\tilde{L} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} p+1 & q \\ q & p-1 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix},$$

where their results are under the assumption that q is locally absolutely continuous. For more results on embedded single eigenvalue, one can refer to [11, 15].

For many embedded eigenvalues of Schrödinger operators or Dirac operators, Naboko [18] constructed smooth potentials such that $L_{0,q}$ has dense (rationally independent) embedded eigenvalues. Naboko's constructions work for Schrödinger operators as well. Simon [22] constructed potentials such that the associated Schrödinger operator has dense embedded eigenvalues. More recently, Jitomirskaya and Liu [8] introduced a novel idea to construct embedded eigenvalues for Laplacian on manifolds, which is referred to as piecewise constructions. This approach turns out to be quite robust. Liu and his collaborators developed the approach of piecewise constructions to construct embedded eigenvalues for various models [13, 15, 16, 17]. For more results on embedded eigenvalue problems, one can refer to [12, 14, 21].

In this paper, we adapt the approach of piecewise construction to study embedded eigenvalue problems of Dirac operators. The main strategy of proofs for our main theorems follow from that of [8, 13, 17]. In the current case of Dirac operators, new difficulties and challenges arise from the Dirac operator being vector valued and its potential consisting of a pair of functions p and q (unlike the models in [8, 13, 15, 16, 17]).

2. Proof of Theorems 1.1 and 1.2

Let $(u(x), v(x))^T$ be a solution of

$$L_{p,q} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix}.$$

We define the Prüfer variables $R(x)$ and $\theta(x)$ of λ by

$$u(x) = R(x) \cos \theta(x),$$

and

$$v(x) = R(x) \sin \theta(x).$$

Clearly, we have

Proposition 2.1. *Let $R(x)$ and $\theta(x)$ be the Prüfer variables of λ . Then λ is an eigenvalue of the Dirac operator if and only if $R \in L^2(0, \infty)$.*

By the equation

$$L_{p,q} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix},$$

we obtain

$$\frac{R'}{R} = -q(x) \cos 2\theta(x) - p(x) \sin 2\theta(x), \quad (5)$$

and

$$\theta' = -\lambda + q(x) \sin 2\theta(x) - p(x) \cos 2\theta(x). \quad (6)$$

Set $q(x) = V(x) \cos \varphi(x)$, $p(x) = V(x) \sin \varphi(x)$. Note that p and q are completely determined by V and φ . By (5) and (6), one has

$$\frac{R'}{R} = -V(x) \cos(2\theta(x) - \varphi(x)), \quad (7)$$

and

$$\theta' = -\lambda + V(x) \sin(2\theta(x) - \varphi(x)). \quad (8)$$

It is obvious that equations (7) and (8) are equivalent to

$$L_{p,q} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix}.$$

By Proposition 2.1, we only need to study (7) and (8).

Proof of Theorem 1.1. Assume

$$\limsup_{x \rightarrow \infty} |xV(x)| = \limsup_{x \rightarrow \infty} x \sqrt{p(x)^2 + q(x)^2} = A < \frac{1}{2}. \quad (9)$$

For any $\epsilon > 0$ (small enough so that $A + \epsilon < \frac{1}{2}$), there exists x_0 so that for any $x > x_0$, one has

$$|V(x)| \leq \frac{A + \epsilon}{1 + x}.$$

By (7) and (9), we have

$$\begin{aligned}\ln R(x) &= \ln R(x_0) - \int_{x_0}^x V(t) \cos(2\theta(t) - \varphi(t)) dt \\ &\geq O(1) - (A + \epsilon) \int_{x_0}^x \frac{1}{1+t} dt \\ &= O(1) - (A + \epsilon) \ln x.\end{aligned}$$

By the assumption, there exists a positive constant k such that, for large x , we have

$$R(x) \geq kx^{-\frac{1}{2}}.$$

This implies that $R \notin L^2(0, \infty)$. Hence by Proposition 2.1, λ is not an eigenvalue of $L_{p,q}$. \square

Proof of Theorem 1.2 for $A > \frac{1}{2}$. We construct p and q as follows:

$$V(x) = \frac{A}{1+x}, \quad x \geq 0,$$

and

$$\varphi(x) = -2\lambda x + 2\theta(0), \quad x \geq 0.$$

By (8) and the uniqueness theorem (see for example [24, Theorem 2.2]), one has for any $x \geq 0$,

$$2\theta(x) - \varphi(x) \equiv 0.$$

Thus from (7) we obtain

$$\begin{aligned}\ln R(x) &= \ln R(0) - \int_0^x \frac{A}{1+t} dt \\ &= O(1) - A \ln x.\end{aligned}$$

We immediately obtain that for some small $\epsilon > 0$ and any large x ,

$$R(x) \leq x^{-\frac{1}{2}-\epsilon}.$$

Therefore, $R \in L^2(0, \infty)$ and by Proposition 2.1, λ is an eigenvalue of the corresponding Dirac operator $L_{p,q}$. \square

Proof of Theorem 1.2 for $A = \frac{1}{2}$. Let $\epsilon_n = \frac{1}{2n}$, $a_n = e^{n^3}$. Set

$$V(x) = \frac{A + \epsilon_n}{x}, \quad x \in [a_n, a_{n+1}),$$

and

$$\varphi(x) = -2\lambda x + 2\theta(0).$$

By (8) and the uniqueness theorem, one has for any $x \geq 0$,

$$2\theta(x) - \varphi(x) \equiv 0.$$

By (7), one has

$$\begin{aligned}\ln R(a_{n+1}) - \ln R(a_n) &= - \int_{a_n}^{a_{n+1}} \frac{A + \epsilon_n}{x} dx \\ &= -(A + \epsilon_n) \ln \frac{a_{n+1}}{a_n}.\end{aligned}\quad (10)$$

For $t \in [a_n, a_{n+1})$, we have

$$\begin{aligned}\ln R(t) - \ln R(a_n) &= - \int_{a_n}^t \frac{A + \epsilon_n}{x} dx \\ &= -(A + \epsilon_n) \ln \frac{t}{a_n}.\end{aligned}\quad (11)$$

From (10), we obtain

$$\ln R(a_n) = \ln R(a_0) - \sum_{j=0}^{n-1} (A + \epsilon_j) \ln \frac{a_{j+1}}{a_j}.$$

Therefore, one has

$$\begin{aligned}R(a_n) &= O(1)e^{-\sum_{j=0}^{n-1} (A + \epsilon_j) \ln \frac{a_{j+1}}{a_j}} \\ &= O(1) \prod_{j=0}^{n-1} a_{j+1}^{-(A + \epsilon_j)} a_j^{A + \epsilon_j} \\ &= O(1) \prod_{j=1}^n a_j^{-(A + \epsilon_{j-1})} \prod_{j=1}^{n-1} a_j^{A + \epsilon_j} \\ &= O(1) a_n^{-(A + \epsilon_{n-1})} \prod_{j=1}^{n-1} a_j^{\epsilon_j - \epsilon_{j-1}}.\end{aligned}\quad (12)$$

By (11) and (12), we conclude

$$\begin{aligned}R(t) &= O(1)R(a_n)e^{-(A + \epsilon_n) \ln \frac{t}{a_n}} \\ &= O(1)R(a_n)t^{-(A + \epsilon_n)} a_n^{A + \epsilon_n} \\ &= O(1) \prod_{j=1}^n a_j^{\epsilon_j - \epsilon_{j-1}} t^{-(A + \epsilon_n)}.\end{aligned}\quad (13)$$

It follows that

$$\begin{aligned} \int_{a_n}^{a_{n+1}} R(t)^2 dt &= O(1) \int_{a_n}^{a_{n+1}} \prod_{j=1}^n a_j^{\frac{1}{j}-\frac{1}{j-1}} t^{-1-\frac{1}{n}} dt \\ &\leq O(1) \prod_{j=1}^n e^{-j} \frac{n}{e^{n^2}} \\ &\leq O(1) \frac{n}{e^{n^2}}. \end{aligned} \tag{14}$$

This implies that $R \in L^2(0, \infty)$, by Proposition 2.1, λ is an eigenvalue of the corresponding Dirac operator $L_{p,q}$. \square

3. Proof of Theorems 1.3 and 1.4

We assume that λ and λ_j are different values. Denote the Prüfer variables of λ and λ_j by $R(x), \theta(x)$ and $R_j(x), \theta_j(x)$, respectively.

Recall that $V(x)$ and $\varphi(x)$ uniquely determine p and q . Define $V(x) = V(x, b)$ and $\varphi(x) = \varphi(x, \lambda, a, \varphi_0)$ on $[a, \infty)$ by

$$V(x, b) = \frac{C}{1+x-b}, \tag{15}$$

and

$$\varphi(x, \lambda, a, \varphi_0) = -2\lambda(x-a) + 2\varphi_0, \tag{16}$$

where C is a constant will be defined later, $a > b$ and $\varphi_0 = \theta(a)$.

Lemma 3.1. Fix $b > 0$. Let $V(x)$ be defined by (15). Let $\varphi(x)$ be defined by (16), and $\lambda \neq \lambda_j$. Let $\theta_j(x)$ be a solution of

$$\theta_j'(x) = -\lambda_j + V(x) \sin(2\theta_j(x) - \varphi(x)), \tag{17}$$

then we have

$$\int_{x_0}^x \frac{1}{1+t-b} \cos(2\theta_j(t) - \varphi(t)) dt = \frac{O(1)}{x_0-b}, \tag{18}$$

for any $x > x_0 > a$.

Proof. By (16) and (17) we have

$$2\theta_j'(t) - \varphi'(t) = 2(\lambda - \lambda_j) + \frac{O(1)}{1+t-b},$$

and

$$2\theta_j''(t) - \varphi''(t) = \frac{O(1)}{1+t-b}.$$

It follows that

$$\begin{aligned} & \int_{x_0}^x \frac{1}{1+t-b} \cos(2\theta_j(t) - \varphi(t)) dt \\ &= \frac{\sin(2\theta_j(t) - \varphi(t))}{2(\lambda - \lambda_j) + \frac{O(1)}{1+t-b}} \frac{1}{1+t-b} \Big|_{x_0}^x + O(1) \int_{x_0}^x \frac{1}{(1+t-b)^2} dt \\ &= \frac{O(1)}{x_0 - b}. \end{aligned}$$

□

Lemma 3.2. Fix $b > 0$. Let $V(x)$ be defined by (15) on $[a, \infty)$. Let $\varphi(x)$ be defined by (16) on $[a, \infty)$, and $\lambda \neq \lambda_j$. Let $R(x), \theta(x)$ and $R_j(x), \theta_j(x)$ be the Prüfer variables of λ and λ_j , respectively. For any $x > a$,

$$\ln R(x) - \ln R(a) \leq -100 \ln \frac{x-b}{a-b} + C, \tag{19}$$

$$\ln R(x) \leq \ln R(a), \tag{20}$$

where C is a large constant depending on λ and λ_j , and for any $x > x_0 \geq a$ with large enough $x_0 - b$, we have

$$R_j(x) \leq 1.5R_j(x_0). \tag{21}$$

Proof. By (8), (15) and (16), and the uniqueness theorem, one has

$$2\theta(x) - \varphi(x) = 0.$$

Therefore, by (7) and (15), we have

$$\begin{aligned} \ln R(x) &= \ln R(a) - \int_a^x \frac{C}{1+t-b} dt \\ &= \ln R(a) - C \ln \frac{1+x-b}{1+a-b}. \end{aligned}$$

Then we immediately obtain (19) and (20).

By (5) and (18), we have

$$\begin{aligned} \ln R_j(x) &= \ln R_j(x_0) - \int_{x_0}^x \frac{C \cos(2\theta_j(t) - \varphi(t))}{1+t-b} dt \\ &= \ln R_j(x_0) + \frac{O(1)}{x_0 - b}. \end{aligned}$$

Hence we obtain (21). □

Proposition 3.3. Let λ and $S = \{\lambda_j\}_{j=1}^k$ be distinct real numbers. Given $\varphi_0 \in [0, \pi)$, if $x_1 > x_0 > b$, then there exist constants $K(\lambda, S), C(\lambda, S)$ (independent of b, x_0 and x_1) and $\tilde{V}(x, \lambda, S, x_0, x_1, b) \in C^\infty$ and $\varphi(x, \lambda, S, x_0, x_1, b, \varphi_0)$ such that for $x_0 - b > K(\lambda, S)$ the following holds:

(1): for $x_0 \leq x \leq x_1$, $\text{supp}(\tilde{V}) \subset (x_0, x_1)$, and

$$|\tilde{V}(x, \lambda, S, x_0, x_1, b)| \leq \frac{C(\lambda, S)}{x - b}. \tag{22}$$

(2): the solution of Dirac equation

$$L_{p,q} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix},$$

with the boundary condition $\frac{u(x_0)}{v(x_0)} = \cot \varphi_0$ satisfies

$$R(x_1) \leq C(\lambda, S) \left(\frac{x_1 - b}{x_0 - b} \right)^{-100} R(x_0), \tag{23}$$

and for $x_0 < x < x_1$,

$$R(x) \leq 2R(x_0). \tag{24}$$

(3): the solution of Dirac equation

$$L_{p,q} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda_j \begin{pmatrix} u \\ v \end{pmatrix},$$

with any boundary condition satisfies for $x_0 < x \leq x_1$,

$$R_j(x) \leq 2R_j(x_0). \tag{25}$$

Proof. Let $V(x)$ be given by (15) and $\varphi(x)$ be given by (16), with $a = x_0$ and $C = C(\lambda, S)$. Let $x = x_1$ in (19), (20) and (21). We smooth $V(x)$ near x_0, x_1 to obtain $\tilde{V}(x)$. Notice that by (7), a small perturbation of $V(x)$ will only give a small change of $R(x)$ and $R_j(x)$. Hence Lemma 3.2 still holds with slightly larger constants. We complete the proof. \square

Proof of Theorems 1.3 and 1.4. With the help of Proposition 3.3, the proofs of Theorems 1.3 and 1.4 follow from the construction step by step as appearing in [8, 13, 17]¹.

We only give an outline of the proof here. Let $\{N_r\}_{r \in \mathbb{Z}^+}$ be a non-decreasing sequence which goes to infinity arbitrarily slowly depending on $h(x)$ ². We further assume $N_{r+1} = N_r + 1$ when $N_{r+1} > N_r$. At the r th step, we take N_r eigenvalues into consideration. Applying Proposition 3.3, we construct potentials with N_r pieces, where each piece comes from (22) with λ being an eigenvalue. The main difficulty is to control the size of each piece (denote by T_r). The construction in [8, 13, 17] only uses inequalities (22), (23) and (24) to obtain appropriate T_r and N_r . Hence Proposition 3.3 implies Theorems 1.3 and 1.4. \square

¹We should mention that although models in [8, 13, 17] are second-order differential equations, the first step is to write those equations in a system of two first order differential equations.

²For most $r \in \mathbb{N}$, we have $N_{r+1} = N_r$, and when $N_{r+1} > N_r$, we take $N_{r+1} = N_r + 1$. This will ensure N_r increases to infinity slowly.

Acknowledgments

This work was completed in an ongoing High School and Undergraduate Research Program “STODO” (Spectral Theory Of Differential Operators) at Texas A&M University. We would like to thank Wencai Liu for managing the program, introducing this project and many inspiring discussions. The authors are also grateful to the anonymous referee, whose comments led to an improvement of our manuscript. This work was partially supported by NSF DMS-2015683 and DMS-2000345.

References

- [1] ATKINSON, F. V.; EVERITT, W. N. Bounds for the point spectrum for a Sturm-Liouville equation. *Proc. Roy. Soc. Edinburgh Sect. A* **80** (1978), no. 1-2, 57–66. MR529569, Zbl 0426.34015. 1318
- [2] EVANS, W. D.; HARRIS, B. J. Bounds for the point spectra of separated Dirac operators. *Proc. Roy. Soc. Edinburgh Sect. A* **88** (1981), no. 1-2, 1–15. MR611296, Zbl 0447.34022. 1320
- [3] GESZTESY, F.; NICHOLS, R. On absence of threshold resonances for Schrödinger and Dirac operators. *Discrete Contin. Dyn. Syst. Ser. S* **13** (2020), no. 12, 3427–3460. MR4160130, Zbl 1459.35107. 1318
- [4] GESZTESY, F.; SAKHNOVICH, A. The inverse approach to Dirac-type systems based on the A -function concept. *J. Funct. Anal.* **279** (2020), no. 6, 108609, 40. MR4096724, Zbl 1470.34053. 1318
- [5] GESZTESY, F.; ZINCHENKO, M. Renormalized oscillation theory for Hamiltonian systems. *Adv. Math.* **311** (2017), 569–597. MR3628224, Zbl 1381.34050. 1318
- [6] HARRIS, B. J. Bounds for the eigenvalues of separated Dirac operators. *Proc. Roy. Soc. Edinburgh Sect. A* **95** (1983), no. 3-4, 341–366. MR726883, Zbl 0563.34025. 1318
- [7] HU, Y.; BONDARENKO, N. P.; SHIEH, C.; YANG, C. Traces and inverse nodal problems for Dirac-type integro-differential operators on a graph. *Appl. Math. Comput.* **363** (2019), 124606, 10. MR3984178, Zbl 1433.34027. 1318
- [8] JITOMIRSKAYA, S.; LIU, W. Noncompact complete Riemannian manifolds with dense eigenvalues embedded in the essential spectrum of the Laplacian. *Geom. Funct. Anal.* **29** (2019), no. 1, 238–257. MR3925109, Zbl 1430.58022. 1320, 1326
- [9] KATO, T. Growth properties of solutions of the reduced wave equation with a variable coefficient. *Comm. Pure Appl. Math.* **12** (1959), 403–425. MR108633, Zbl 0091.09502. 1318
- [10] LEVITAN, B. M.; SARGSIAN, I. S. Sturm-Liouville and Dirac operators. Mathematics and its Applications (Soviet Series), 59. *Kluwer Academic Publishers Group, Dordrecht*. Translated from the Russian, 1991. MR1136037, Zbl 0657.34002. 1318
- [11] LIU, W. Sharp bound on the largest positive eigenvalue for one-dimensional Schrödinger operators. Preprint, 2017. arXiv:1709.05611. 1320
- [12] LIU, W. The asymptotical behaviour of embedded eigenvalues for perturbed periodic operators. *Pure Appl. Funct. Anal.* **4** (2019), no. 3, 589–602. MR4008385, Zbl 1458.47014. 1320
- [13] LIU, W. Criteria for eigenvalues embedded into the absolutely continuous spectrum of perturbed Stark type operators. *J. Funct. Anal.* **276** (2019), no. 9, 2936–2967. MR3926137, Zbl 1412.34235. 1320, 1326
- [14] LIU, W. Sharp bounds for finitely many embedded eigenvalues of perturbed Stark type operators. *Math. Nachr.* **293** (2020), no. 9, 1776–1790. MR4148833, Zbl 07261817. 1320
- [15] LIU, W. Criteria for embedded eigenvalues for discrete Schrödinger operators. *Int. Math. Res. Not. IMRN* (2020), no. 20, 15803–15832. MR4329883, Zbl 1480.39016. 1320
- [16] LIU, W.; LYU, K. One dimensional discrete Schrödinger operators with resonant embedded eigenvalues. Preprint, 2022. arXiv:2207.00194. 1320

- [17] LIU, W.; ONG, D. C. Sharp spectral transition for eigenvalues embedded into the spectral bands of perturbed periodic operators. *J. Anal. Math.* **141** (2020), no. 2, 625–661. MR4179772, Zbl 1466.34074. 1320, 1326
- [18] NABOKO, S. N. On the dense point spectrum of Schrödinger and Dirac operators. *Teoret. Mat. Fiz.* **68** (1986), no. 1, 18–28. MR875178, Zbl 0607.34023. 1320
- [19] REMLING, C. Spectral theory of canonical systems. De Gruyter Studies in Mathematics, 70. *De Gruyter, Berlin*, 2018. MR3890099, Zbl 1401.37004. 1318
- [20] REMLING, C.; SCARBROUGH, K. The essential spectrum of canonical systems. *J. Approx. Theory.* **254** (2020), 105395, 11. MR4076316, Zbl 1457.34122. 1318
- [21] SCHMIDT, K. M. Dense point spectrum for the one-dimensional Dirac operator with an electrostatic potential. *Proc. Roy. Soc. Edinburgh Sect. A* **126** (1996), no. 5, 1087–1096. MR1415824, Zbl 0859.34074. 1320
- [22] SIMON, B. Some Schrödinger operators with dense point spectrum. *Proc. Amer. Math. Soc.* **125** (1997), no. 1, 203–208. MR1346989, Zbl 0888.34071. 1320
- [23] SIMON, B. Tosio Kato’s work on non-relativistic quantum mechanics: part 1. *Bull. Math. Sci.* **8** (2018), no. 1, 121–232. MR3775269, Zbl 1416.81063. 1318
- [24] TESCHL, G. Ordinary differential equations and dynamical systems. Graduate Studies in Mathematics, 140. *American Mathematical Society, Providence, RI.*, 2012. MR2961944, Zbl 1263.34002. 1322
- [25] VON NEUMANN, J.; WIGNER, E. Über merkwürdige diskrete Eigenwerte. *Phys. Zeit.* **30** (1929), 467–470. JFM 55.0520.04. 1318
- [26] WEIDMANN, J. Oszillationsmethoden für Systeme gewöhnlicher Differentialgleichungen. *Math. Z.* **119** (1971), 349–373. MR285758, Zbl 0206.10002. 1319
- [27] YANG, C.; HUANG, Z. Inverse spectral problems for $2m$ -dimensional canonical Dirac operators. *Inverse Problems* **23** (2007), no. 6, 2565–2574. MR2441020, Zbl 1153.34006. 1318
- [28] ZHANG, R.; YANG, C.; BONDARENKO, N. P. Inverse spectral problems for the Dirac operator with complex-valued weight and discontinuity. *J. Differential Equations* **278** (2021), 100–110. MR4199393, Zbl 1462.34042. 1318

(Vishwam Khapre) DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA
vishwam28@tamu.edu

(Kang Lyu) SCHOOL OF MATHEMATICS AND STATISTICS, NANJING UNIVERSITY OF SCIENCE AND TECHNOLOGY, NANJING 210094, JIANGSU, CHINA
lvkang201905@outlook.com

(Andrew Yu) PHILLIPS ACADEMY, 180 MAIN ST, ANDOVER, MA 01810, USA
andrewyu45@gmail.com

This paper is available via <http://nyjm.albany.edu/j/2022/28-55.html>.