

Bounds for the number of rational points on curves over function fields

Amílcar Pacheco and Fabien Pazuki

ABSTRACT. We provide an upper bound for the number of rational points on a nonisotrivial curve defined over a one variable function field over a finite field. The bound only depends on the curve and the field, and not on the Jacobian variety of the curve.

CONTENTS

1. Introduction	131
Acknowledgements	133
2. Proof of Theorem 1.1 part (a)	134
2.1. Tools from étale cohomology	135
3. Proof of Theorem 1.1 part (b): F -descent in characteristic p	136
3.1. Selmer groups	136
3.2. Group cohomology	137
3.3. p -rank and Lie algebras	137
3.4. Local computation	138
3.5. Global result	139
3.6. Using F -descent and finishing the proof	140
4. Further remarks	141
References	142

1. Introduction

Let k be a finite field of cardinality q and of positive characteristic p . Let \mathcal{C} a smooth, projective, geometrically connected curve defined over k of genus g . Denote by $K = k(\mathcal{C})$ its function field. Let K_s be a separable closure of K . Given a smooth, projective, geometrically connected curve X

Received February 16, 2013; revised April 12, 2013.

2010 *Mathematics Subject Classification*. 11G35, 11R58, 14G40, 14G25, 14H05.

Key words and phrases. Abelian varieties, rational points, curves, function fields.

The first author wishes to thank the Institut de Mathématiques de Jussieu, Paris, for its support during his sabbatical. This stay had also the support of CNPq (Brazil), CNRS (France) and Paris Science Foundation. Both authors would like to thank Université Bordeaux 1, ANR-10-BLAN-0115 Hamot and ANR-10-JCJC-0107 Arivaf (France).

defined over K of genus $d \geq 2$, the analogue of the Mordell's conjecture asks whether the set $X(K)$ is finite.

This does not come without a constraint, otherwise this question would have a trivial negative answer. One has to assume that X is nonisotrivial. This means that there does not exist a smooth projective geometrically connected curve X_0 defined over a finite extension l of k and a common extension L of both K and l such that $X \times_K L \cong X_0 \times_l L$ (cf. [Sa66]). Under the aforementioned condition the finiteness of $X(K)$ is a theorem due to Samuel [Sa66].

Our purpose is to give an effective upper bound for the cardinality of the set $X(K)$ in terms of a minimal number of invariants associated with our given geometric situation. Namely, our upper bound will depend on the following parameters:

- (i) The genus d of X/K .
- (ii) The genus g of \mathcal{C}/k .
- (iii) The inseparable degree p^e of the map $u : \mathcal{U} \rightarrow \mathcal{M}_g$ from the affine sub-curve \mathcal{U} of \mathcal{C} (where X has good reduction) to the fine moduli scheme of genus g curves. The map u is induced by a model $\mathcal{X} \rightarrow \mathcal{C}$ of X/K .¹
- (iv) The conductor $f_{X/K}$ of X/K (this will be defined later in the text).

Let us insist on the fact that this bound does not depend on the Jacobian variety J_X of the curve X . The rank r of the Mordell–Weil group $J_X(K)$ is not used in the bound. In the geometric case this rank is bounded in terms of d and g (cf. Ogg's bound, see Remark 2.3). We observe that the bound in terms of the conductor of J_X/K would be stronger (cf. Proposition 2.8), but the point is to show that the bound can be expressed in terms of only the curve itself. Our main result is the following theorem.

Theorem 1.1. *Let k be a finite field of cardinality q and characteristic p , \mathcal{C} a smooth, projective, geometrically connected curve defined over k of genus g and denote by $K = k(\mathcal{C})$ its function field. Let X/K be a smooth, projective, geometrically connected curve defined over K of genus $d \geq 2$. We suppose that X is nonisotrivial.*

- (a) *If X is defined over K , but not over K^p , then the following inequality holds:*

$$\#X(K) \leq p^{2d \cdot (2g+1) + f_{X/K}} \cdot 3^d \cdot (8d - 2) \cdot d!.$$

Denote the right hand side of the latter inequality by C_{BV} .

- (b) *More generally, suppose that $p > 2d + 1$. If X is defined over K^{p^e} , but not over $K^{p^{e+1}}$ for some natural integer e , then*

$$\#X(K) \leq C_{\text{BV}} \cdot C_{\text{desc}}^e,$$

¹Observe that p^e does not depend on the choice of the model $\mathcal{X} \rightarrow \mathcal{C}$, for a further discussion see Section 3.

where one can take

$$C_{\text{desc}} = q^{c_0} \text{ and } c_0 = g - 1 + f_{X/K} + \frac{1}{2} \cdot p^{e+1} \cdot d \cdot (2g - 2 + 2^{4d^2} \cdot f_{X/K}).$$

Remark 1.2. In a recent paper [CoUlVo12], Conceição, Ulmer and Voloch provide some explicit examples of curves X_a for which the number of rational points cannot be bounded by a quantity independent of X_a . Consider the curve $X_a/\mathbb{F}_p(t)$ defined by the affine equation $y^2 = x \cdot (x^r + 1) \cdot (x^r + a^r)$, where $p > 3$ and r is coprime to $2p$ and $a = t^{p^n+1}$. Say $n = 2^m$ with $m \in \mathbb{N}$ big enough. Then

$$\#X_a(\mathbb{F}_p(t)) \geq d(n) \gg \log n \gg \log \log h(X_a/\mathbb{F}_p(t)) \gg \log \log f_{J_{X_a/\mathbb{F}_p(t)}},$$

where the last step is obtained thanks to [HiPa13, Corollary 6.12], this inequality relates the conductor of the Jacobian to its differential height.

In fact, the height $h(X_a/\mathbb{F}_p(t))$ is the height of the equation defining the curve (for instance defined through its associated Chow form). One can give an upper bound for the theta height in terms of the height of the equation as in [Re10, Théorème 1.3 and Proposition 1.1]. Next the theta height of an abelian variety can be bounded from above by the differential height of the abelian variety, because the former can be realized as the height on an appropriate moduli space (cf. [HiPa13, Section 3], this also known in the case of number fields, cf. [Pa12, Theorem 1.1]).

The history of explicit upper bounds for $\#X(K)$ starts with the work of Szpiro [Sz81] which in fact gives an explicit upper bound for the height of points in $X(K)$. This depends, however, on the geometry of a semi-stable fibration on curves $\phi : \mathcal{X} \rightarrow \mathcal{C}$ which gives a minimal model of X/K over \mathcal{C} . One of the goals of the current paper is to obtain a bound which does not depend on the geometry of any model of X/K over \mathcal{C} .

We start with an upper bound for the number of elements of $X(K)$, when X is defined over K , but not over K^p . This follows from a result due to Buïum and Voloch [BuVo96]. In fact, their result gives an explicit proof of a conjecture of Lang, Mordell's conjecture is a particular case of the latter. We then extend the first result to curves which can be defined over K^{p^n} for some integer $n \geq 1$. The crucial step is the F -descent of abelian varieties in characteristic $p > 0$ (see Section 3).

Acknowledgements. The authors would like to thank José Felipe Voloch for inspiring conversations during our stay at Ennio di Giorgi Center in Pisa, Italy, in October 2012. This paper was completed during the sabbatical leave of the first author which was spent at the Institut de Mathématiques de Jussieu, Paris. He would like to thank this institution for its support. This stay had also the support of CNPq (Brazil), CNRS (France) and Paris Science Foundation. The first author would also like to thank UFRJ for giving him this leave. Both authors would like to thank Université Bordeaux 1, ANR-10-BLAN-0115 Hamot and ANR-10-JCJC-0107 Arivaf (France). Both

authors would like to thank the organizers of the conference in Pisa for their invitation.

2. Proof of Theorem 1.1 part (a)

We start by recalling:

Theorem 2.1 (Buium–Voloch, [BuVo96, Theorem]). *Let k be a finite field of characteristic p , K a one variable function field over k , X/K a smooth, projective, geometrically connected curve defined over K of genus $d \geq 2$. We suppose that X is not defined over K^p . Let Γ a subgroup of $J_X(K_s)$ such that $\Gamma/p\Gamma$ is finite. The following inequality holds:*

$$\#(X \cap \Gamma) \leq \#(\Gamma/p\Gamma) \cdot p^d \cdot 3^d \cdot (8d - 2) \cdot d!.$$

Remark 2.2. Let $\Gamma = J_X(K)$. Then $J_X(K)/pJ_X(K)$ is a finite group by the Mordell–Weil theorem. Writing $J_X(K) = \mathbb{Z}^r \times J_X(K)_{\text{tor}}$, where $r = \text{rk } J_X(K)$, one has $J_X(K)/pJ_X(K) = (\mathbb{Z}/p\mathbb{Z})^r \times J_X(K)_{\text{tor}}/pJ_X(K)_{\text{tor}}$. Its order is bounded from above by p^{d+r} . Next we discuss an upper bound for the rank.

Remark 2.3. Let k be any field and \mathcal{C} smooth projective geometrically connected curve over k . Denote by $K = k(\mathcal{C})$ its function field. Let A/K be a nonconstant abelian variety over K and denote by (τ, B) its K/k -trace (cf. [La83]). Let \bar{k} be an algebraic closure of k . A theorem due to Lang and Néron ([La83], [LaNe59]) states that the quotient group $A(\bar{k}(\mathcal{C}))/\tau B(\bar{k})$ is a finitely generated abelian group. A fortiori, the quotient group $A(K)/\tau B(k)$ is also finitely generated. Ogg in the 60’s (cf. [Ogg62]) produced the following upper bound for the rank of the geometric quotient $A(\bar{k}(\mathcal{C}))/\tau B(\bar{k})$ (hence of $A(K)/\tau B(k)$). Below we define the conductor $f_{A/K}$ of A/K . Let $d_0 = \dim B$. Then the upper bound is

$$2d \cdot (2g - 2) + f_{A/K} + 4d_0 \leq 4d \cdot g + f_{A/K}.$$

In particular, if K is a one variable function field over a finite field, then

$$\text{rk } A(K) \leq 4d \cdot g + f_{A/K}.$$

Definition 2.4. Let $\ell \neq p$ be a prime number. Denote by $T_\ell(A)$ the ℓ -adic Tate module of A and define $V_\ell(A) = T_\ell(A) \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell$. For each place v of K , denote by I_v an inertia group at v (well-defined up to conjugation). Let ϵ_v be the codimension of the subgroup of I_v -invariants $V_\ell(A)^{I_v}$ in $V_\ell(A)$. Let δ_v be the Swan conductor of $H_{\text{ét}}^1(A_{K_s}, \mathbb{Q}_\ell)$ (cf. [Se69]). Define the conductor divisor $\mathfrak{F}_{A/K} = \sum_v (\epsilon_v + \delta_v) \cdot [v]$, where v runs through the places of K . Denote $f_{A/K} = \deg \mathfrak{F}_{A/K}$.

Definition 2.5. A model of X/K over \mathcal{C} is a smooth, projective, geometrically connected surface \mathcal{X} defined over k and a proper flat morphism $\phi : \mathcal{X} \rightarrow \mathcal{C}$. Each place v of K is identified with a point of \mathcal{C} . Denote by κ_v the residue field at v (which is a finite field) and let $\bar{\kappa}_v$ be an algebraic

closure of κ_v . Denote by \mathcal{X}_v the fiber of ϕ at v . For an algebraic variety Z defined over a field l and for an extension L of l , denote $Z_L = Z \times_l L$.

2.1. Tools from étale cohomology.

Definition 2.6. Let Z be a smooth variety defined over a field l with algebraic closure \bar{l} . Denote by $n = \dim Z$, for each $0 \leq i \leq 2n$, let $H_{\text{ét}}^i(X_{\bar{l}}, \mathbb{Q}_\ell)$ be the i -th étale cohomology group of Z/\bar{l} . Define the Euler–Poincaré characteristic of Z/\bar{l} by $\chi(Z/\bar{l}) = \sum_{i=0}^{2n} (-1)^i \dim_{\mathbb{Q}_\ell} H_{\text{ét}}^i(Z_{\bar{l}}, \mathbb{Q}_\ell)$. This number is indeed independent from the choice of ℓ .

Definition 2.7. Fix a place v of K . The Artin conductor of the curve X over K at v is defined as $f_{X/K,v} = -\chi(X_{K_s}) + \chi(\mathcal{X}_{v,\bar{\kappa}_v}) + \delta_v$, where $\chi(X_{K_s})$, respectively $\chi(\mathcal{X}_{v,\bar{\kappa}_v})$ denotes the Euler–Poincaré characteristic of X_{K_s} , respectively $\mathcal{X}_{v,\bar{\kappa}_v}$. The term δ_v denotes the Swan conductor of $H^1(X_{K_s}, \mathbb{Q}_\ell)$ at v (cf. [LiSa00, end of p. 414] for the definition of the Artin conductor, [Se69] for the definition of the Swan conductor, as well as [Bl87, §1]). Define the global conductor of the curve X/K by $f_{X/K} = \sum_v f_{X/K,v} \cdot \deg v$, where v runs through the places of K .

The following proposition is a consequence of the subsequent lemma in [Bl87].

Proposition 2.8. *We have the inequality $f_{J_X/K} \leq f_{X/K}$.*

Lemma 2.9 ([Bl87, Lemma 1.2]). *Fix a place v of K and let I_v be an inertia subgroup of $\text{Gal}(K_s/K)$ at v . Then:*

- (I) $H_{\text{ét}}^i(X_{K_s}, \mathbb{Q}_\ell)^{I_v} \cong H_{\text{ét}}^i(\mathcal{X}_{v,\bar{\kappa}_v}, \mathbb{Q}_\ell)$ for $i = 0, 1$.
- (II) *Let M_v be the free abelian group generated by the irreducible components of $\mathcal{X}_{v,\bar{\kappa}_v}$. Since the individual components are not necessarily defined over κ_v , there is an action of $\hat{\mathbb{Z}} \cong \text{Gal}(\bar{\kappa}_v/\kappa_v)$ on M_v . Moreover, there is an exact sequence of $\hat{\mathbb{Z}}$ -modules:*

$$0 \rightarrow \mathbb{Q}_\ell(-1) \rightarrow M_v \otimes \mathbb{Q}_\ell(-1) \rightarrow H_{\text{ét}}^2(\mathcal{X}_{v,\bar{\kappa}_v}, \mathbb{Q}_\ell) \rightarrow H_{\text{ét}}^2(X_{K_s}, \mathbb{Q}_\ell)^{I_v} \rightarrow 0.$$

Remark 2.10. The definition of the conductor given in [LiSa00] agrees with that given in [Bl87] (up to sign).

Proof of Proposition 2.8. It follows from the definition of $f_{X/K,v}$, Lemma 2.9 and the fact that the action of the Galois group $\text{Gal}(K_s/K)$ on the étale cohomology groups $H_{\text{ét}}^i(X_{K_s}, \mathbb{Q}_\ell)$ (for $i = 0, 2$) is trivial that we have an equality:

$$f_{X/K,v} = \dim_{\mathbb{Q}_\ell} H_{\text{ét}}^1(X_{K_s}, \mathbb{Q}_\ell) - \dim_{\mathbb{Q}_\ell} H_{\text{ét}}^1(X_{K_s}, \mathbb{Q}_\ell)^{I_v} + m_v - 1 + \delta_v,$$

where m_v denotes the number of the irreducible components of $\mathcal{X}_{v,\bar{\kappa}_v}$. The proposition now follows from observing that $H_{\text{ét}}^1(X_{K_s}, \mathbb{Q}_\ell) \cong H_{\text{ét}}^1(J_{K_s}, \mathbb{Q}_\ell)$ (cf. [Mi85, Corollary 9.6]). □

Definition 2.11. Let l be a field of characteristic $p > 0$ and Z/l a smooth algebraic variety. Let $F_{\text{abs}} : l \rightarrow l$ be the absolute Frobenius map defined by $a \mapsto a^p$. We define the smooth variety $Z^{(p)}$ by the Cartesian diagram

$$\begin{array}{ccc} Z^{(p)} & \longrightarrow & Z \\ \downarrow & & \downarrow \\ \text{Spec } l & \xrightarrow{F_{\text{abs}}} & \text{Spec } l. \end{array}$$

The relative Frobenius morphism $F : Z \rightarrow Z^{(p)}$ is defined so that composed with the upper horizontal arrow of the diagram gives the absolute Frobenius morphism $F_{\text{abs}} : Z \rightarrow Z$. This situation can be iterated by taking for any integer $e \geq 1$ to get the e -th power $F^e : Z \rightarrow Z^{(p^e)}$ of F .

Proof of Theorem 1.1 part (a). Let $j : X \hookrightarrow J_X$ be the embedding of X into its Jacobian variety. Denote by $X(K) = \{x_1, \dots, x_m\}$ the finite set of K -rational points of X . Let Γ be the subgroup of $J_X(K)$ generated by the images $\{j(x_1), \dots, j(x_m)\}$ of these points under the embedding j . Observe that

$$\#(\Gamma/p\Gamma) \leq \#(J_X(K)/pJ_X(K)) \leq p^{r+d} \leq p^{d(4g+1)+f_{X/K}}$$

by Remarks 2.2 and 2.3 and Proposition 2.8. The result is now a consequence of Theorem 2.1. \square

3. Proof of Theorem 1.1 part (b): F -descent in characteristic p

Let K be a one variable function field over a finite field of characteristic $p > 0$.

3.1. Selmer groups. (See [Ul91, §1].) We start with the more general set-up of an isogeny $f : A \rightarrow B$ of nonconstant abelian varieties defined over K . We use the convention that all cohomology groups will be computed in terms of the flat site. As a consequence, on the flat site of K , we have a short exact sequence of group schemes given by $0 \rightarrow \ker f \rightarrow A \rightarrow B \rightarrow 0$.

For any place v of K , let K_v be the completion of K at v . Denote by $\text{Sel}(K_v, f)$ the image of the coboundary map $\delta_v : B(K_v) \rightarrow H^1(K_v, \ker f)$. The global Selmer group $\text{Sel}(K, f)$ is defined as the subset of those elements in $H^1(K, \ker f)$ whose restriction modulo v is trivial in $\text{Sel}(K_v, f)$ for every place v of K .

Recall that the Tate–Shafarevich group $\text{III}(A/K)$ is defined as

$$\ker(H^1(K, A) \rightarrow \prod_v H^1(K_v, A)).$$

The isogeny f induces a map $f : \text{III}(A/K) \rightarrow \text{III}(B/K)$ whose kernel is denoted by $\text{III}(A/K)_f$. Then $\text{Sel}(K, f)$ appears in the following exact sequence of groups: $0 \rightarrow B(K)/f(A(K)) \rightarrow \text{Sel}(K, f) \rightarrow \text{III}(A/K)_f \rightarrow 0$. In practice $\text{Sel}(K, f)$ is finite and effectively computable.

Denote by \mathcal{O}_v the valuation ring of K_v . If both A and B have good reduction over \mathcal{O}_v , then the restriction map $H^1(\mathcal{O}_v, \ker f) \rightarrow H^1(K_v, \ker f)$ induces an isomorphism $\text{Sel}(K_v, f) \cong H^1(\mathcal{O}_v, \ker f)$. If L/K_v is a Galois extension of degree prime to $\deg f$, then the inclusion map

$$H^1(K_v, \ker f) \rightarrow H^1(L, \ker f)$$

induces an isomorphism $\text{Sel}(K_v, f) \cong \text{Sel}(L, f)^G$. Similarly, if L/K is a finite Galois extension of degree prime to $\deg f$, then $\text{Sel}(K, f) = \text{Sel}(L, f)^G$.

3.2. Group cohomology. (See [Se79, Chapter VII, §2].) Let G be a group, A an abelian group with an action of G on the left, denoted by

$$(\sigma \in G, a \in A) \mapsto \sigma \cdot a.$$

A one cocycle is a map $a : G \rightarrow A$ such that $a_{\sigma\sigma'} = \sigma \cdot a_{\sigma'} + a_\sigma$. Note that if $A = B \oplus C$, where B and C are also abelian groups, then composing a with projections on B , respectively C , one gets two one cocycles $b : G \rightarrow B$ and $c : G \rightarrow C$ so that $a = (b, c)$. A one cocycle $a : G \rightarrow A$ is a coboundary if there exists $\alpha \in A$ such that $a_\sigma = \sigma \cdot \alpha - \alpha$, for every $\sigma \in G$. Again, if a is a one cocycle which is a coboundary and $A = B \oplus C$, then b and c are coboundaries as well. Denote by $H^1(A, G)$ the group of one cocycles with values in A modulo coboundaries. In the previous case, we have

$$H^1(A, G) = H^1(G, B) \oplus H^1(G, C).$$

3.3. p -rank and Lie algebras. (See [Mu70, Theorem, p. 139].) In the case of an abelian variety \mathfrak{A} defined over an algebraically closed field l , denote by $\mathfrak{A}^\vee = \text{Pic}^0 \mathfrak{A}$ its dual abelian variety. Then $\text{Lie } \mathfrak{A}^\vee \cong H^1(\mathfrak{A}, \mathcal{O}_{\mathfrak{A}})$, moreover under this isomorphism the p -th power map on $\text{Lie } \mathfrak{A}^\vee$ corresponds to the Frobenius map F on $H^1(\mathfrak{A}, \mathcal{O}_{\mathfrak{A}})$. In particular,

$$r = p\text{-rk } \mathfrak{A} = \dim_{\mathbb{F}_p} \mathfrak{A}[p] = \dim_{\mathbb{F}_p} H^1(\mathfrak{A}, \mathcal{O}_{\mathfrak{A}})^F = \dim_{\mathbb{F}_p} (\text{Lie } \mathfrak{A})_{\text{ss}}.$$

It is known from p -linear algebra that $\ker F \cong \mu_p^{\oplus r}$. Therefore, by group cohomology, $H^1(K_v, \ker F) \cong H^1(K_v, \mu_p)^{\oplus r} \cong (K_v^*/K_v^{*p})^{\oplus r}$.

We now return to our original abelian variety A/K , and denote by $\varphi : A \rightarrow \mathcal{C}$ its Néron model over \mathcal{C} . Let $e_{\mathcal{A}} : \mathcal{C} \rightarrow \mathcal{A}$ be its neutral section. Denote $\omega_{\mathcal{A}/\mathcal{C}} = e_{\mathcal{A}}^* \Omega_{\mathcal{A}/\mathcal{C}}^1$ and $\tilde{\omega}_{\mathcal{A}/\mathcal{C}} = \wedge^d \omega_{\mathcal{A}/\mathcal{C}}$, where $d = \dim A$. The degree of $\tilde{\omega}_{\mathcal{A}/\mathcal{C}}$ is defined as the differential height of A/K and denoted by $h_{\text{diff}}(A/K)$. Then $\tilde{\omega}_{\mathcal{A}/\mathcal{C}}$ corresponds to a unique Weil divisor $\mathcal{D}_{\mathcal{A}/\mathcal{C}}$ on \mathcal{C} .

The relative version of the first paragraph of this section states that if $\varphi^\vee : \mathcal{A}^\vee \rightarrow \mathcal{C}$ is the dual group scheme of $\varphi : \mathcal{A} \rightarrow \mathcal{C}$, then $\text{Lie } \mathcal{A}^\vee \cong R^1 \varphi_* \mathcal{O}_{\mathcal{A}}$. The latter is dual to $\Omega_{\mathcal{C}}^1(\mathcal{D}_{\mathcal{A}/\mathcal{C}})$.

Denote by \mathcal{C} the Cartier operator acting on $\Omega_{\mathcal{C}}^1$ (cf. [Se56]). By the previous isomorphism the p -th power map on $\text{Lie } \mathcal{A}^\vee$ corresponds to the map F on $R^1 \varphi_* \mathcal{O}_{\mathcal{A}}$. Next by Serre's duality theorem for curves, the latter map corresponds to the map \mathcal{C} on $\Omega_{\mathcal{C}}^1(\mathcal{D}_{\mathcal{A}/\mathcal{C}})$. In particular, $p\text{-Lie } \mathcal{A}^\vee$ is dual to $\Omega_{\mathcal{C}}^1(\mathcal{D}_{\mathcal{A}/\mathcal{C}})^{\mathcal{C}}$.

Let D_1, \dots, D_r be a basis of $H^0(\mathcal{C}, p\text{-Lie } \mathcal{A}^\vee)$, then there exists $a_i \in \bar{k}$ such that $D_i^p = a_i \cdot D_i$, where \bar{k} denotes the algebraic closure of k . Denote $\mathcal{L}_{\mathcal{A}} = \text{Lie } \mathcal{A}^\vee$. In this case the Oort–Tate classification of finite flat group schemes of order p in characteristic p implies that we may associate to a_i a group scheme $G_{\mathcal{L}_{\mathcal{A}}, 0, a_i} = G_{0, a_i}$ over \mathcal{C} (cf. [Mi86, Chapter III, 0.9], [OoTa70]).

3.4. Local computation.

3.4.1. Potential good reduction. We fix a place v of K . Let K_s be a separable closure of K and denote by I_v an inertia subgroup of $\text{Gal}(K_s/K)$ at v (this is well-defined up to conjugation). By definition $A_{K_v} = A \times_K K_v$ has potential good reduction at v , if there exists a finite extension K' of K_v such that $A_{K'} = A_{K_v} \times_{K_v} K'$ has good reduction at v . By [SeTa68, Theorem 2], if $\ell \neq p$ is a prime number and $\rho_\ell : \text{Gal}(K_s/K) \rightarrow \text{Aut}(T_\ell(A))$ is the Galois representation on the Tate module, then A has potential good reduction at v if and only if $\rho_\ell(I_v)$ is finite.

3.4.2. Description of Selmer groups. (See [U191, §3].) Suppose that we are in this case and let K' be as above. Denote by v' the valuation of K' over v . Let $n = -v'(\mathcal{D}_{\mathcal{A}/\mathcal{C}})$. Define $U^{[i]} = \{\bar{f} \in K_v^*/K_v^{*p} \mid \text{ord}_v(f) \geq 1 - i\}$. Apply [Mi86, Chapter III, §7.5] to get $H^1(\mathcal{O}_{K'}, G_{0, a_i}) \cong U_{K'}^{[pm]}$. The previous properties of Selmer groups give

$$\text{Sel}(K', F) \cong H^1(\mathcal{O}_{K'}, \ker F) \cong H^1(\mathcal{O}_{K'}, \mu_p)^{\oplus r} \cong (U_{K'}^{[pm]})^{\oplus r}.$$

Then taking Galois invariants as in Subsection 3.1, we get

$$\text{Sel}(K, F) \cong \text{Sel}(K', F)^{\text{Gal}(K'/K)} \cong (U_{K_v}^{[i]})^{\oplus r},$$

where $i = -p \cdot v(\mathcal{D}_{\mathcal{A}/\mathcal{C}})$.

3.4.3. Potential semi-abelian reduction. We suppose that $p > 2d + 1$, where $d = \dim A$. In this case, A acquires everywhere semi-stable reduction over $L = K(A[\ell])$ for any prime $\ell \neq p$ (cf. [Gr72]). In particular, for the places where the reduction is already good, we are reduced to the latter subsection. So we suppose that we are in the case where A has bad semi-abelian reduction at a place w of L . In this case by [BoLuRa90] there exists a semi-abelian variety $G \in \text{Ext}^1(B, \mathbb{G}_m^t)$ defined over L_w , where B is an abelian variety with good reduction at w , and a lattice $\Lambda \subset G(L_w)$ such that $A(L_w) \cong G(L_w)/\Lambda$.

The action of the absolute Frobenius map F of G engenders the semi-abelian variety $G^{(p)} \in \text{Ext}^1(B^{(p)}, \mathbb{G}_m^t)$, where $B^{(p)}$ is the image of B under F . One checks that $A^{(p)}(L_w) \cong G^{(p)}(L_w)/\Lambda^{(p)}$, where the lattice $\Lambda^{(p)}$ is generated by the vectors obtained from the generators of Λ by raising each component to p . Recall that there exists an isogeny $V : A^{(p)} \rightarrow A$ (called the Verschiebung) such that $V \circ F = [p]_A$ and $F \circ V = [p]_{A^{(p)}}$.

The coboundary map is given by $A(L_w) \rightarrow H^1(L_w, \ker F)$. We have already shown that the latter is isomorphic to $(L_w^*/L_w^{*p})^{\oplus r}$. The previous parametrization composed with V then gives a surjective map

$$G^{(p)}(L_w)/\Lambda^{(p)} \rightarrow (L_w^*/L_w^{*p})^{\oplus r}.$$

In particular, this implies that the coboundary map is surjective, i.e.,

$$\mathrm{Sel}(L_w, F) \cong (L_w^*/L_w^{*p})^{\oplus r}.$$

Finally once more taking Galois invariants we get

$$\mathrm{Sel}(K_v, F) \cong \mathrm{Sel}(L_w, F)^{\mathrm{Gal}(L_w/K_v)} \cong (K_v^*/K_v^{*p})^{\oplus r},$$

(cf. [UI91, §3]).

3.5. Global result. We denote by v, good the set of places v of K where A has good reduction. Similarly v, bad denotes the set of places v of K where A has bad reduction. Let

$$D = \sum_{v, \text{bad}} [v] - \sum_{v, \text{good}} i_v \cdot [v] \in \mathrm{Div} \mathcal{C}, \text{ where } i_v = -p \cdot v(\mathcal{D}_{A/\mathcal{C}}).$$

We observe that

$$0 < \deg D \leq f_{A/K} + p \cdot h_{\mathrm{diff}}(A/K).$$

Note there exists an injective map $K^*/K^{*p} \hookrightarrow \Omega_K^1$ given by $\bar{f} \mapsto df/f$. Observe that the image is exactly $(\Omega_{\mathcal{C}}^1)^{\mathcal{C}}$. The local results imply $\bar{f} \in \mathrm{Sel}(K, F)$ if and only if $df/f \in H^0(\mathcal{C}, \Omega_{\mathcal{C}}^1(-D))^{\mathcal{C}}$. By the Riemann–Roch theorem, one gets

$$\begin{aligned} \dim_{\mathbb{F}_q} H^0(\mathcal{C}, \Omega_{\mathcal{C}}^1(-D))^{\mathcal{C}} &\leq \dim_{\mathbb{F}_q} H^0(\mathcal{C}, \Omega_{\mathcal{C}}^1(-D)) \\ &= g - 1 + \deg D \leq g - 1 + f_{A/K} + p \cdot h_{\mathrm{diff}}(A/K). \end{aligned}$$

Remark 3.1. As we have mentioned before p^e is the inseparable degree of the map $u : \mathcal{U} \rightarrow \mathcal{M}_g$ from the open sub-curve \mathcal{U} of \mathcal{C} to the fine moduli space \mathcal{M}_g of genus g curves induced from a model $\mathcal{X} \rightarrow \mathcal{C}$ of X/K . This invariant is indeed birational. It may be interpreted as follows: p^e is the largest power of p such that X is defined over K^{p^e} , but not over $K^{p^{e+1}}$.

We need to assume from now on that $p > 2d + 1$. In this case, if $\ell \neq p$ is a prime number and $L = K(A[\ell])$, then A has semi-abelian reduction over L . Furthermore, since L/K is tamely ramified of degree prime to p , then the Swan conductor makes no contribution to $f_{A/K}$, hence $\mathfrak{F}_{A/K} = \sum_v \epsilon_v \cdot [v]$. We now recall the abc-theorem for semi-abelian schemes in characteristic $p > 0$.

Theorem 3.2 ([HiPa13, Theorem 5.3]). *Let A/K be a nonconstant abelian variety with everywhere semi-abelian reduction. Denote by $\phi : \mathcal{A} \rightarrow \mathcal{C}$ a Néron model of A/K . Let $\mathfrak{P}_{A/K}$ be the set of places of K where A has bad reduction. Denote $\bar{s} = \sum_{v \in \mathfrak{P}_{A/K}} \deg v$. Let $e_{\mathcal{A}} : \mathcal{C} \rightarrow \mathcal{A}$ be a section of ϕ and*

$\omega_{A/C} = e_A^* \wedge^d \Omega_{A/C}^1$. Suppose that $p > 2d + 1$. Then the following inequality holds

$$(3.1) \quad h_{\text{diff}}(A/K) = \deg \omega_{A/C} \leq \frac{1}{2} \cdot p^e \cdot d \cdot (2g - 2 + \bar{s}).$$

Applying Theorem 3.2 to A_L/L we get the following upper bound:

$$h_{\text{diff}}(A_L/L) \leq \frac{1}{2} \cdot p^e \cdot d \cdot (2g - 2 + f_{A_L/L}).$$

By [Pa05, Proposition 3.7] page 371, one has

$$f_{A_L/L} \leq [L : K] \cdot f_{A/K} \leq \ell^{4d^2} \cdot f_{A/K}.$$

Choosing $\ell = 2$ (remember that $p > 2d + 1 \geq 3$, so $p \neq 2$) provides:

$$(3.2) \quad h_{\text{diff}}(A_L/L) \leq \frac{1}{2} \cdot p^e \cdot d \cdot (2g - 2 + 2^{4d^2} \cdot f_{A/K}).$$

Let $c_0 = g - 1 + f_{A/K} + \frac{1}{2} \cdot p^e \cdot d \cdot (2g - 2 + 2^{4d^2} \cdot f_{A/K})$. Then $\#\text{Sel}_{A_L}(L, F) \leq q^{c_0}$. Recall that since L/K is Galois of order prime to p ,

$$\text{Sel}_{A_L}(L, F)^G = \text{Sel}_A(K, F).$$

We conclude that $\#\text{Sel}_A(K, F) \leq q^{c_0}$. Denote $C_{\text{desc}} = q^{c_0}$.

3.6. Using F -descent and finishing the proof. The following lemma allows us to conclude the proof of item (b) of Theorem 1.1.

Lemma 3.3. *Let $X \hookrightarrow J_X$ be a curve over a field K as before embedded into its Jacobian variety J_X . Suppose X is defined over K^{p^e} , but not over $K^{p^{e+1}}$. Suppose that one has the estimate*

$$\#(J_X(K)/F(J_X(K))) \leq C_{\text{desc}}.$$

Then one obtains the upper bound

$$\#X(K) \leq C_{\text{BV}} \cdot C_{\text{desc}}^e,$$

where $C_{\text{BV}} = p^{2d \cdot (2g+1) + f_{X/K}} \cdot 3^d \cdot (8d - 2) \cdot d!$.

Proof. Suppose that X is defined over K , but not over K^p . Then without any further hypothesis the theorem is proven in part (a). Suppose now that X is defined over K^p , but not over K^{p^2} . Then there exists a smooth, geometrically connected, projective curve X_1 defined over K , but not over K^p such that

$$F : X_1 \rightarrow X_1^{(p)} = X$$

is the relative Frobenius morphism of X_1 . Consider the following decomposition into right cosets

$$X(K) = \bigcup_i F(X_1(K)) + P_i.$$

Under the embedding $j : X \hookrightarrow J_X$ this decomposition is included in the decomposition

$$\bigcup_i F(J_{X_1}(K)) + j(P_i).$$

Note that these classes are not necessarily distinct, however this decomposition is contained in the decomposition

$$\bigcup_l F(J_{X_1}(K)) + \alpha_l,$$

where we now consider all representatives of $J_X(K)$ modulo $F(J_{X_1}(K))$. As a consequence we get

$$(X(K) : F(X_1(K))) \leq (J_X(K) : F(J_{X_1}(K))) \leq \#\text{Sel}_{J_X}(K, \ker F) \leq C_{\text{desc}},$$

Recall that F is purely inseparable, therefore $\#F(X_1(K)) \leq C_{\text{BV}}$. Finally we get

$$\#X(K) \leq C_{\text{BV}} \cdot C_{\text{desc}}.$$

Suppose now that X is defined over K^{p^2} , but not over K^{p^3} . As before there exist curves X_1, X_2 (with the same description as in the last paragraph) such that

$$X_2 \xrightarrow{F} X_1 = X_2^{(p)} \xrightarrow{F} X = X_1^{(p)} = X_2^{(p^2)}.$$

In this case we have got inequalities

$$\#X_1(K) \leq \#X_2(K) \cdot \#(J_{X_1}(K)/F(J_{X_2}(K))),$$

$$\#X(K) \leq \#X_1(K) \cdot \#(J_X(K)/F(J_{X_1}(K))).$$

Observe that $\#(J_{X_1}(K)/F(J_{X_2}(K))) \leq \#\text{Sel}_{J_{X_1}}(K, \ker F)$. An upper bound for the latter term depends only on J_{X_1} through its conductor. Since J_X and J_{X_1} are F -isogeneous, their conductors coincide. Whence,

$$\#X(K) \leq C_{\text{BV}} \cdot C_{\text{desc}}^2.$$

An easy induction argument then finishes the proof. \square

4. Further remarks

Remark 4.1. We would now like to compare our result with a result similar in nature when we replace the one variable function field K defined over a finite field k by a number field K . In order to do this we refer to the work of Rémond (cf. [Re10]).

Theorem 4.2 (Rémond). *Let X be a smooth, projective, geometrically connected curve of genus $d \geq 2$ defined over a number field K , then one has*

$$\#X(K) \leq (2^{38+2d} \cdot [K : \mathbb{Q}] \cdot d \cdot \max(1, h_\Theta))^{(r+1) \cdot d^{20}},$$

where h_Θ is the theta height of J_X and $r = \text{rk } J_X(K)$.

Remark 4.3. Using Proposition 5.1 page 775 of [Re10], one has $r \ll \log f_{J_X/K}$, as in the function field case, but the bound on the number of points is still dependent on the height of the Jacobian variety. To be more precise, Rémond shows in loc. cit. how to produce a bound depending on the height of a model of the curve (and not of its Jacobian variety), but it seems difficult to get rid of this height. It would be a consequence of a conjecture of Lang and Silverman, as explained in the introduction of [Pa12]. Note that in the function field case, the height of the Jacobian variety J_X is comparable to the degree of its conductor $f_{J_X/K}$. More precisely it is proven in [HiPa13, Corollary 5.12] that we have the following inequalities

$$p^e \cdot f_{A/K} \ll h_{\text{diff}}(A/K) \ll p^e \cdot f_{A/K},$$

where the implied constants depend only on g and d . Note that the upper bound is a consequence of Theorem 3.2. The lower inequality has a simpler proof (cf. loc. cit.).

References

- [BoLuRa90] BOSCH, SIEGFRIED; LÜTKEBOHMERT, WERNER; RAYNAUD, MICHEL. Néron models. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*, 21. Springer-Verlag, Berlin, 1990. x+325 pp. ISBN: 3-540-50587-3. MR1045822 (91i:14034), Zbl 0705.14001, doi:10.1007/978-3-642-51438-8.
- [Bl87] BLOCH, SPENCER. De Rham cohomology and conductors of curves. *Duke Math. J.* **54** (1987), no.2, 295–308. MR0899399 (89h:11028), Zbl 0632.14018, doi:10.1215/S0012-7094-87-05417-2.
- [BuVo96] BUIUM, ALEXANDRU; VOLOCH, JOSÉ FELIPE. Lang’s conjecture in characteristic p : an explicit bound. *Compositio Math.* **103** (1996), no. 1, 1–6. MR1404995 (98a:14038), Zbl 0885.14010.
- [CoUIVo12] CONCEIÇÃO, RICARDO; ULMER, DOUGLAS; VOLOCH, JOSÉ FELIPE. Unboundedness of the number of rational points on curves over function fields. *New York J. Math.* **18** (2012), 291–293. MR2928577, Zbl 06032712, arXiv:1204.2001.
- [Gr72] GROTHENDIECK, A. Modèles de Néron et monodromie. *Sém. Géom. algébrique Bois-Marie 1967–1979. Exp. IX in SGA 7*, no. 9, Lect. Notes in Math., 288. Springer, 1972. 313–523. Zbl 0248.14006.
- [HiPa13] HINDRY, M.; PACHECO, A. An analogue of the Brauer–Siegel theorem for abelian varieties in positive characteristic. Preprint, <https://sites.google.com/site/amilcarpachecoresearch/publications>, 2013.
- [La83] LANG, SERGE. *Fundamentals of Diophantine geometry*. Springer-Verlag, New York, 1983. xviii+370 pp. ISBN: 0-387-90837-4. MR0715605 (85j:11005), Zbl 0644.14007.
- [LaNe59] LANG, SERGE; NÉRON, A. Rational points of abelian varieties over function fields. *Amer. J. Math.* **81** (1959), 95–118. MR0102520 (21 #1311), Zbl 0099.16103.
- [Liu94] LIU, QING. Conducteur et discriminant minimal de courbes de genre 2. *Compositio Math.* **94** (1994), no. 1, 51–79. MR1302311 (96b:14038), Zbl 0837.14023.
- [LiSa00] LIU, QING; SAITO, TAKESHI. Inequality for conductor and differential of a curve over a local field. *J. Algebraic Geom.* **9** (2000), no. 3, 409–424. MR1752009 (2001g:14043), Zbl 0992.14008.

- [Mi85] MILNE, J. S. Jacobian varieties. *Arithmetic Geometry*, 167–212. Springer, New York, 1986. MR0861976, Zbl 0604.14018, doi:10.1007/978-1-4613-8655-1_7.
- [Mi86] MILNE, J. S. Arithmetic duality theorems. Perspectives in Mathematics, 1. Academic Press, Inc., Boston, MA, (1986). x+421 pp. ISBN: 0-12-498040-6. MR0881804 (88e:14028), Zbl 0613.14019.
- [MB85] MORET-BAILLY, LAURENT. Pinceaux de variétés abéliennes. *Astérisque* **129** (1985), 266 p. MR0797982 (87j:14069), Zbl 0595.14032.
- [Mu70] MUMFORD, DAVID. Abelian varieties. Tata Institute of Fundamental Research Studies in Mathematics, no. 5. Oxford University Press, London 1970. viii+242 pp. MR0282985 (44 #219), Zbl 0223.14022.
- [Ogg62] OGG, A. P. Cohomology of abelian varieties over function fields. *Ann. of Math. (2)* **76** (1962) 185–212. MR0155824 (27 #5758), Zbl 0121.38002.
- [Pa05] PACHECO, AMÍLCAR. On the rank of abelian varieties over function fields. *Manuscripta Math.* **118** (2005), no. 3, 361–381. MR2183044 (2006g:11115), Zbl 1082.11037, arXiv:math/0404384, doi:10.1007/s00229-005-0597-7.
- [Pa12] PAZUKI, FABIEN. Theta height and Faltings height. *Bull. Soc. Math. France* **140** (2012), no.1, 19–49. MR2903770, Zbl 1245.14029, arXiv:0907.1458.
- [Re10] RÉMOND, GAËL. Nombre de points rationnels des courbes. *Proc. Lond. Math. Soc. (3)* **101** (2010), 759–794. MR2734960 (2011k:11088), Zbl 1210.11073, doi:10.1112/plms/pdq005.
- [Sa66] SAMUEL, PIERRE. Compléments à un article de Hans Grauert sur la conjecture de Mordell *Inst. Hautes Études Sci. Publ. Math.* **29** (1966), 55–62. MR0204430 (34 #4272), Zbl 0144.20102, doi:10.1007/BF02684805.
- [Se56] SERRE, JEAN-PIERRE. Sur la topologie des variétés algébriques en caractéristique p . *Symposium internacional de topología algebraica*, 24–53. Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958. MR0098097 (20 #4559), Zbl 0098.13103.
- [Se69] SERRE, JEAN-PIERRE. Facteurs locaux des fonctions zêta des variétés algébriques (Définitions et conjectures). *Sém. Delange-Pisot-Poitou*, **11** (1969/70), no. 19. Zbl 0214.48403.
- [Se79] SERRE, JEAN-PIERRE. Local fields. Graduate Texts in Mathematics, 67. Springer-Verlag, New York-Berlin, 1979. viii+241 pp. ISBN: 0-387-90424-7. MR0554237 (82e:12016), Zbl 0423.12016.
- [SeTa68] SERRE, JEAN-PIERRE; TATE, JOHN. Good reduction of abelian varieties. *Ann. of Math. (2)* **88** (1968), 492–517. MR0236190 (38 #4488), Zbl 0172.46101, doi:10.2307/1970722.
- [Sz81] SZPIRO, LUCIEN. Propriétés numériques du faisceau dualisant relatif. *Astérisque* **86** (1981), 44–78. Zbl 0517.14006.
- [OoTa70] TATE, JOHN; OORT, FRANS. Group schemes of prime order. *Ann. Sci. École Norm. Sup. (4)* **3** (1970), 1–21. MR0265368 (42 #278), Zbl 0195.50801.
- [Ul91] ULMER, DOUGLAS L. p -descent in characteristic p . *Duke Math. J.* **62** (1991), no. 2, 237–265. MR1104524 (92i:11068), Zbl 0742.14028, doi:10.1215/S0012-7094-91-06210-1.
- [Vo91] VOLOCH, J. F. On the conjectures of Mordell and Lang in positive characteristics. *Invent. Math.* **104** (1991), no. 3, 643–646. MR1106753 (92d:11067), Zbl 0735.14019, doi:10.1007/BF01245094.

UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, INSTITUTO DE MATEMÁTICA, AV. ATHOS DA SILVEIRA RAMOS 149, CT, BL. C, CIDADE UNIVERSITÁRIA, ILHA DO FUNDÃO, CAIXA POSTAL 68530, 21941-909 RIO DE JANEIRO, RJ, BRASIL

amilcar@acd.ufrj.br

INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UNIVERSITÉ BORDEAUX I, 351, COURS DE LA LIBÉRATION, 33405 TALENCE, FRANCE

fabien.pazuki@math.u-bordeaux1.fr

This paper is available via <http://nyjm.albany.edu/j/2013/19-9.html>.