

A note on multipliers between model spaces

Emmanuel Fricain and Rishika Rupam

ABSTRACT. In this note, we study the multipliers from one model space to another. In the case when the corresponding inner functions are meromorphic, we give both necessary and sufficient conditions ensuring this set of multipliers is not trivial. Our conditions involve the Beurling–Malliavin densities and are based on the deep work of Makarov–Poltoratski on injectivity of Toeplitz operators.

CONTENTS

| | |
|---|-----|
| 1. Introduction | 147 |
| 2. Preliminaries | 148 |
| 2.1. Basic notations | 148 |
| 2.2. Meromorphic inner functions and model spaces | 149 |
| 2.3. Toeplitz operators and a characterization of multipliers | 149 |
| 2.4. Beurling–Malliavin densities | 150 |
| 3. Main theorem and examples | 152 |
| References | 155 |

1. Introduction

For a pair of inner functions U and V on the upper half-plane

$$\mathbb{C}_+ = \{z \in \mathbb{C} : \Im(z) > 0\},$$

the multipliers set $\mathcal{M}(U, V)$ is the set of analytic functions Φ on \mathbb{C}_+ such that

$$\Phi K_U \subset K_V.$$

Here K_U (respectively K_V) is the model space associated to U (respectively to V). See Section 2.2 for the definition. A basic question here is whether or not

$$\mathcal{M}(U, V) \neq \{0\}?$$

Received June 21, 2017.

2010 *Mathematics Subject Classification.* 30J05, 30H10.

Key words and phrases. Multipliers, model spaces, Beurling–Malliavin densities.

The authors were supported by Labex CEMPI (ANR-11-LABX-0007-01).

A source of inspiration for this paper stems from [GMR16, Tim16] which examined various pre-orders on the set of partial isometries and contractions on Hilbert spaces and their relationship to their associated Livšic characteristic functions. It turns out, for example, that when the Livšic characteristic functions u and v for two partial isometries A and B are inner (on the unit disc), the issue of whether or not A is "less than" B can be rephrased as to whether or $\mathcal{M}(u, v) \neq \{0\}$. Another motivation comes from the work of Crofoot [Cro94] who studied the onto multipliers.

In [FHR18], the authors characterize the multipliers from one model space to another in terms of kernels of Toeplitz operators and Carleson measures for model spaces. However, it is widely understood that both the injectivity problem of Toeplitz operators and the Carleson measures question for model spaces are rather difficult. As a result, it is not easy to apply the characterization obtained in [FHR18] in concrete situations. In this paper, we pursue this line of research. We consider the case when U and V are both meromorphic on \mathbb{C} . Our aim is to simplify the characterization proved in [FHR18] and to apply it to several examples.

2. Preliminaries

2.1. Basic notations. We use the standard notation

$$\mathcal{H}^p = \mathcal{H}^p(\mathbb{C}_+),$$

$1 \leq p \leq \infty$, for the Hardy space of the upper half-plane and as usual we identify functions in \mathcal{H}^p with their boundary values on \mathbb{R} . We denote by Π the Poisson measure on \mathbb{R} ,

$$d\Pi(t) = \frac{dt}{1+t^2},$$

and by $L^1_{\Pi} = L^1(\mathbb{R}, \Pi)$. The Hilbert transform of a function $h \in L^1_{\Pi}$ is defined as the singular integral

$$\tilde{h}(x) = \lim_{\varepsilon \rightarrow 0} \frac{1}{\pi} \int_{|x-t|>\varepsilon} \left[\frac{1}{x-t} + \frac{t}{1+t^2} \right] h(t) dt.$$

Recall that outer functions H are of the form

$$H = e^{h+i\tilde{h}} \quad \text{on } \mathbb{R},$$

for some $h \in L^1_{\Pi}$. Recall also that if $h \in L^1_{\Pi}$, then $\tilde{h} \in L^{o(1,\infty)}_{\Pi}$ (the weak L^1 space), i.e.,

$$\Pi\{|\tilde{h}| > A\} = o\left(\frac{1}{A}\right), \quad A \rightarrow \infty.$$

See [Mas09, Corollary 14.6].

We shall need the elementary Blaschke factor on \mathbb{C}_+ with zero at i :

$$b_i(z) := \frac{z-i}{z+i},$$

and

$$k_i(z) = \frac{1}{\pi} \frac{1}{z+i},$$

the corresponding kernel (of \mathcal{H}^2) at i .

2.2. Meromorphic inner functions and model spaces. Recall that an inner function U on the upper half-plane is a bounded and analytic function on \mathbb{C}_+ with boundary values of modulus one almost everywhere on \mathbb{R} . In this paper, we are interested in the situation when the inner function U can be extended into a meromorphic function in \mathbb{C} . Such functions are called meromorphic inner functions (MIF) on the upper half-plane. They can be easily described via the standard Blaschke/singular factorization. All MIFs have the following form:

$$U(z) = Ce^{iaz} \prod_{n=0}^{\infty} e^{i\alpha_n} \frac{z-w_n}{z-\bar{w}_n}, \quad (z \in \mathbb{C}_+),$$

where a is a nonnegative constant, w_n is a sequence of points in \mathbb{C}_+ tending to infinity as $n \rightarrow \infty$ and satisfying the Blaschke condition

$$\sum_{n=0}^{\infty} \frac{\Im(w_n)}{1+|w_n|^2} < \infty,$$

C is a unimodular constant and α_n is a real number chosen so that

$$e^{i\alpha_n} = \frac{\left| \frac{i-w_n}{i-\bar{w}_n} \right|}{\frac{i-w_n}{i-\bar{w}_n}}.$$

Associated to an inner function U on \mathbb{C}_+ , the model space K_U is defined by

$$K_U := \mathcal{H}^2 \cap (U\mathcal{H}^2)^\perp.$$

We also have the following equivalent definition

$$(1) \quad K_U = \mathcal{H}^2 \cap U\overline{\mathcal{H}^2},$$

where $\overline{\mathcal{H}^2}$ is often regarded as the Hardy space of the lower half-plane.

2.3. Toeplitz operators and a characterization of multipliers. Recall that to every $\varphi \in L^\infty(\mathbb{R})$, there corresponds the Toeplitz operator $T_\varphi : \mathcal{H}^2 \rightarrow \mathcal{H}^2$ defined by

$$T_\varphi(f) = P_+(\varphi f), \quad f \in \mathcal{H}^2,$$

where P_+ is the orthogonal projection of $L^2(\mathbb{R})$ onto \mathcal{H}^2 . Using (1), it is immediate to see that, when the function U is inner, then

$$(2) \quad \ker T_{\overline{U}} = K_U.$$

In [FHR18], the following characterization of multipliers is proved.

Theorem 2.1 (Fricain–Hartmann–Ross). *Let U and V be inner functions with $|U'(x)| \asymp 1, x \in \mathbb{R}$, and let Φ be a function holomorphic on \mathbb{C}_+ . Then the following are equivalent:*

- (1) $\Phi \in \mathcal{M}(U, V)$.
- (2) $\Phi k_i \in \ker T_{\overline{b_i} \overline{V} U}$ and $\sup_{x \in \mathbb{R}} \int_x^{x+1} |\Phi(t)|^2 dt < \infty$.

Note that the second condition appearing in (2) says that the measure $|\Phi(t)|^2 dt$ is a Carleson measure for K_U (see [Bar00, Theorem 5.1]), ensuring that $\Phi K_U \subset \mathcal{H}^2$.

As one see from Theorem 2.1, the noninjectivity of a certain Toeplitz operator is necessary for the set of multipliers being nontrivial. The problem of injectivity of Toeplitz operators is a classical problem in analysis, being related to completeness of exponential systems on $L^2(0, 2\pi)$. In [MP05, MP10], Makarov–Poltoratski extended the theory of Beurling–Malliavin density to model spaces related to MIF. See next section for a brief discussion on their results. We just mention here an easy result which shall be used below.

Lemma 2.2. *Let B be a finite Blaschke product, Θ an inner function which is not a finite Blaschke product and let $1 \leq p \leq \infty$. Then*

$$\ker T_{B\overline{\Theta}} \cap \mathcal{H}^p \neq \{0\}.$$

Proof. Let us write

$$B(z) = \prod_{j=1}^k \left(\frac{z - w_j}{z - \overline{w_j}} \right)^{m_j}$$

and define the linear map

$$T : \begin{cases} K_{\Theta} \cap \mathcal{H}^p & \longrightarrow \mathbb{C}^N \\ f & \longmapsto (f^{(s)}(\lambda_j))_{\substack{1 \leq j \leq k \\ 1 \leq s \leq m_j}} \end{cases}$$

where $N = \sum_{1 \leq j \leq k} m_j$. Since Θ is not a finite Blaschke product, we know that $K_{\Theta} \cap \mathcal{H}^p$ is of infinite dimension and then T is not one-to-one. Hence there exists a function $f \in K_{\Theta} \cap \mathcal{H}^p$, $f \neq 0$, such that for every $1 \leq j \leq k$, $1 \leq s \leq m_j$, $f^{(s)}(\lambda_j) = 0$. We can write $f = Bg$ for some $g \in \mathcal{H}^p$. It remains to note that using (2), we have

$$T_{B\overline{\Theta}}(g) = P_+(\overline{\Theta}Bg) = T_{\overline{\Theta}}(f) = 0. \quad \square$$

2.4. Beurling–Malliavin densities. Let $\Lambda \subset \mathbb{C}_+ \cup \mathbb{R}$. In [MP05, MP10], Makarov and Poltoratski connected the Beurling–Malliavin density of Λ to the injectivity of the kernel of a related Toeplitz operator. We briefly recall some of these facts here. First, let $\Lambda \subset \mathbb{R}$ be a discrete sequence. We say that Λ is *strongly a -regular* if

$$(3) \quad \int_{\mathbb{R}} \frac{|n_{\Lambda}(x) - ax|}{1 + x^2} dx < \infty,$$

where n_Λ is the counting function of Λ defined by

$$n_\Lambda(x) = \begin{cases} \text{card}(\Lambda \cap [0, x]) & \text{if } x \geq 0 \\ -\text{card}(\Lambda \cap [x, 0]) & \text{if } x < 0. \end{cases}$$

It is known (see [Pol15, MiP10]) that the interior Beurling–Malliavin (BM) density of a discrete sequence Λ can be defined as

$$D_*(\Lambda) := \sup\{a : \exists \text{ strongly } a\text{-regular subsequence } \Lambda' \subset \Lambda\}.$$

Similarly, the exterior BM density is defined as

$$D^*(\Lambda) := \inf\{a : \exists \text{ strongly } a\text{-regular supsequence } \Lambda' \supset \Lambda\}.$$

These definitions extend to the upper half-plane as well [MP10] in the following way. Let $\Lambda \subset \mathbb{C}_+$ be a discrete sequence, then

$$D_*(\Lambda) := D_*(\Lambda^*),$$

where $\Lambda^* := \{\lambda^* : \lambda \in \Lambda, \Re \lambda \neq 0\}$, $\lambda^* := [\Re(\lambda^{-1})]^{-1}$.

Example 2.3. Let $\Lambda = \{n + i\}_{n \in \mathbb{Z}}$. Then $D_*(\Lambda) = D^*(\Lambda) = 1$.

Proof. For $n \in \mathbb{Z}^*$, we have $\lambda_n^* = [\Re(1/(n + i))]^{-1} = (n^2 + 1)/n$. The counting function of this sequence is odd and $n_{\Lambda^*}(x) = n$, for

$$x \in (n + 1/n, n + 1 + 1/(n + 1)),$$

$n > 0$. Then

$$\begin{aligned} \int_2^\infty \frac{|n_{\Lambda^*}(x) - x|}{1 + x^2} dx &= \sum_{n \geq 1} \int_{n+1/n}^{n+1+1/(n+1)} \frac{x - n}{1 + x^2} dx \\ &\leq \sum_n \frac{3}{2} \left(1 + \frac{1}{n+1} - \frac{1}{n}\right) \cdot \frac{1}{n^2 + 1} < \infty. \end{aligned}$$

Thus Λ^* is itself a 1– strongly regular sequence and so

$$D_*(\Lambda) = D^*(\Lambda) = 1. \quad \square$$

It turns out that when Λ is a discrete sequence on \mathbb{R} , then we can construct a MIF Θ with $\sigma(\Theta) := \{x \in \mathbb{R} : \Theta(x) = 1\} = \Lambda$. Then it is proved in [MiP10, MP05] that

$$D_*(\Lambda) = \frac{1}{2\pi} \inf\{a : \ker T_{S^a \bar{\Theta}} = \{0\}\},$$

and

$$D^*(\Lambda) = \frac{1}{2\pi} \sup\{a : \ker T_{\bar{S}^a \Theta} = \{0\}\},$$

where S is the singular inner function defined by $S(z) = e^{iz}$. In terms of Toeplitz kernels, when Λ is a Blaschke sequence in \mathbb{C}_+ , we can replace Θ by

the Blaschke product B_Λ with zeroes on Λ , and we have

$$D_*(\Lambda) = \frac{1}{2\pi} \inf\{a : \ker T_{S^a \bar{B}_\Lambda} = \{0\}\},$$

$$D^*(\Lambda) = \frac{1}{2\pi} \sup\{a : \ker T_{\bar{S}^a B_\Lambda} = \{0\}\}.$$

Note that if $a > b$, then

$$(4) \quad \ker T_{\bar{S}^b B_\Lambda} \neq \{0\} \implies \ker T_{\bar{S}^a B_\Lambda} \neq \{0\}.$$

3. Main theorem and examples

In this section, we give a class of MIFs U and V for which the triviality of $\mathcal{M}(U, V)$ can be reduced to the injectivity of the Toeplitz operator $T_{U\bar{V}}$. We end the section by showing examples of MIFs that fall into this category.

Theorem 3.1. *Let U and V be MIFs with $|U'| \asymp 1$ on \mathbb{R} and let*

$$m := \arg(U) - \arg(Vb_i)$$

on \mathbb{R} . Suppose that either $m \notin \tilde{L}_\Pi^1$ or if $m = \tilde{h}$ for some $h \in L_\Pi^1$, then $e^{-h} \notin L^1(\mathbb{R})$. Then the following three conditions are equivalent.

- (1) $\dim \ker T_{U\bar{V}b_i} \geq 2$.
- (2) $\ker T_{U\bar{V}} \neq \{0\}$.
- (3) $\mathcal{M}(U, V) \neq \{0\}$.

Proof. (1) \implies (2): Since $\dim \ker T_{U\bar{V}b_i} \geq 2$, we can find a function $\Psi \in \ker T_{U\bar{V}b_i}$, $\Psi \neq 0$, such that $\Psi(i) = 0$. Then we can write $\Psi = b_i \Psi_1$ with $\Psi_1 \in \mathcal{H}^2$. Since

$$0 = T_{U\bar{V}b_i}(\Psi) = T_{U\bar{V}}(\Psi_1),$$

we have $\Psi_1 \in \ker T_{U\bar{V}}$ and $\Psi_1 \neq 0$.

(2) \implies (3): Let $\Phi \in \ker T_{U\bar{V}}$ be nonzero. Then there is a function $g \in \mathcal{H}^2$ such that on \mathbb{R} , we have

$$\Phi.U\bar{V} = \bar{g}.$$

Since

$$\frac{\Phi}{z+i}.U\bar{V}.b_i = \frac{\bar{g}}{z+i} \cdot \frac{z+i}{z-i} = \overline{\left(\frac{g}{z+i}\right)} \in \overline{\mathcal{H}^2},$$

then $\Phi b_i \in \ker T_{U\bar{V}b_i}$. Moreover, using $\Phi \in \mathcal{H}^2$, we also have

$$\sup_{x \in \mathbb{R}} \int_x^{x+1} |\Phi(t)|^2 dt < \infty.$$

Thus by Theorem 2.1, we deduce that $\Phi \in \mathcal{M}(U, V)$, which gives (3).

(3) \implies (1): Now assume that $\mathcal{M}(U, V) \neq \{0\}$. Then, according to Theorem 2.1, we know that $\ker T_{U\bar{V}b_i} \neq \{0\}$. We argue by contradiction and suppose that $\dim \ker T_{U\bar{V}b_i} = 1$. First let us prove that $\ker T_{U\bar{V}b_i}$ is generated by an outer function. Indeed, let $f \in \mathcal{H}^2$ such that $\ker T_{U\bar{V}b_i} =$

$\mathbb{C}f$ and write $f = \Theta f_0$ where Θ and f_0 are respectively the inner and outer part of f . Notice that

$$T_{U\overline{Vb_i}}(f_0) = P_+(U\overline{Vb_i}\Theta f) = T_{\Theta}T_{U\overline{Vb_i}}(f) = 0,$$

whence $f_0 \in \ker T_{U\overline{Vb_i}}$ and there exists a $\lambda \in \mathbb{C}$ such that $f_0 = \lambda f$. Thus f is outer.

By definition, there is a function $g \in \mathcal{H}^2$ such that on \mathbb{R} ,

$$U\overline{Vb_i}f = \overline{g}.$$

Let $g = g_i g_0$ be the inner-outer factorization of g . Then

$$U\overline{Vb_i}f g_i = \overline{g_0}.$$

We deduce $g_i f \in \ker T_{U\overline{Vb_i}}$. Since $\ker T_{U\overline{Vb_i}}$ is generated by f , we necessarily get that g_i is a constant of modulus one which we may of course assume to be one. Using that f and g_0 are outer and satisfy $|f| = |g_0|$ on \mathbb{R} , we obtain that $g_0 = f$, and thus

$$(5) \quad Uf = Vb_i \overline{f}.$$

Since f is an outer function that is square integrable on \mathbb{R} , there must exist a function $h_1 \in L^1_{\mathbb{H}}(\mathbb{R})$ such that $f = e^{h_1 + ih_1}$ on \mathbb{R} and $|f| = e^{h_1} \in L^2(\mathbb{R})$. We compare the arguments in (5) which gives

$$m = \arg(U) - \arg(Vb_i) = -2\tilde{h}_1 = \tilde{h},$$

with $h = -2h_1$. But $h \in L^1_{\mathbb{H}}$ and $e^{-h} \in L^1(\mathbb{R})$ a contradiction to our hypothesis. Thus $\dim \ker T_{U\overline{Vb_i}} \geq 2$. \square

Remark 3.2. For the assertions (1) \implies (2) and (2) \implies (3), we only use that U and V are MIFs with $|U'| \asymp 1$ on \mathbb{R} . It is only in the assertion (3) \implies (1) that we use the full hypothesis of the theorem.

It is natural to wonder for which MIFs U and V are the hypotheses of the above theorem satisfied. We give examples here to illustrate that for many pairs of MIFs, this is indeed the case.

Let us denote the singular inner function e^{iz} by $S(z)$. We know that MIFs have the form $S^a B_{\Lambda}$, where $a \geq 0$ and B_{Λ} is a Blaschke product. So we assume that $U = S^a B_{\Lambda_1}$ and $V = S^b B_{\Lambda_2}$.

Example 3.3. Let $U = S^a$ and $V = S^b$. Then we have

$$\mathcal{M}(U, V) \neq \{0\} \iff b \geq a.$$

Indeed, if $b = a$ then $U = V$ and of course the constant functions are multipliers from K_U into K_V . We may assume now that $a \neq b$. Note that

$$m = \arg(U) - \arg(Vb_i) = (a - b)x + 2 \arctan(x)$$

on \mathbb{R} . Since $2 \arctan(x) \in L^{\infty}(\mathbb{R})$ and $(a - b)x \notin L^{\infty(1, \infty)}_{\mathbb{H}}$, the function m does not belong to the space $\tilde{L}^1_{\mathbb{H}}(\mathbb{R})$. Of course, we also have $|U'| \asymp 1$ on

\mathbb{R} . Therefore, we can apply Theorem 3.1 which gives that $\mathcal{M}(U, V) \neq \{0\}$ if and only if $\ker T_{U\bar{V}} \neq \{0\}$. Since $T_{U\bar{V}} = T_{S^{b-a}}$, we get from (2) that

$$b > a \implies \ker T_{U\bar{V}} = K_{S^{b-a}} \implies \mathcal{M}(U, V) \neq \{0\}.$$

On the other hand, if $b < a$, then $T_{U\bar{V}} = T_{S^{a-b}}$ and the operator $T_{U\bar{V}}$ is thus one-to-one, which gives $\mathcal{M}(U, V) = \{0\}$. Note that the result can also be obtained from Crofoot's paper [Cro94]. See also [FHR18, Proposition 2.2].

Example 3.4. Let $U = S^a B_{\Lambda_1}$ and $V = S^b B_{\Lambda_2}$ such that $a \neq b$ and B_{Λ_1} and B_{Λ_2} are finite Blaschke products. Then

$$\mathcal{M}(U, V) \neq \{0\} \iff b > a.$$

Indeed, note that

$$\begin{aligned} m &= \arg(U) - \arg(Vb_i) \\ &= (a - b)x + \arg(B_{\Lambda_1}) - \arg(B_{\Lambda_2}) + 2 \arctan(x). \end{aligned}$$

Since B_{Λ_1} and B_{Λ_2} are finite Blaschke products,

$$\arg(B_{\Lambda_1}) - \arg(B_{\Lambda_2}) + 2 \arctan(x) \in L^\infty(\mathbb{R}).$$

The function $(a - b)x \notin L_\Pi^{o(1, \infty)}$. Thus, the function $m \notin \tilde{L}_\Pi^1(\mathbb{R})$. We also have $|U'| \asymp 1$ on \mathbb{R} . Therefore, we can apply Theorem 3.1 which gives that $\mathcal{M}(U, V) \neq \{0\}$ if and only if $\ker T_{U\bar{V}} \neq \{0\}$. Now if $b > a$, then $T_{U\bar{V}} = T_{\Theta B_{\Lambda_1}}$ where Θ is the inner function defined by $\Theta = S^{b-a} B_{\Lambda_2}$. Hence, by Lemma 2.2, $\ker T_{U\bar{V}} \neq \{0\}$ and thus $\mathcal{M}(U, V) \neq \{0\}$. Note that Coburn's Lemma (see [Nik86, Page 318]) implies that if $b > a$, then $\ker T_{V\bar{U}} = \{0\}$. By symmetry, we thus get that if $b < a$, then $\mathcal{M}(U, V) = \{0\}$.

Example 3.5. Let $U = B_{\Lambda_1} S^a$ and $V = S^b$ where $a \geq 0$, $b > 0$, B_{Λ_1} is an infinite Blaschke product, and let $D := D^*(\Lambda_1)$. Assume that $|U'| \asymp 1$ on \mathbb{R} and $b - a \neq 2\pi D$. Then

$$\mathcal{M}(U, V) \neq \{0\} \iff b - a > 2\pi D.$$

Indeed, if $b - a > 2\pi D$, then by definition of D ,

$$\ker T_{U\bar{V}} = \ker T_{B_{\Lambda_1} S^{b-a}} \neq \{0\}.$$

By Theorem 3.1 and Remark 3.2, we deduce that $\mathcal{M}(U, V) \neq \{0\}$.

Let us now assume that $b - a < 2\pi D$. Using once more the definition of D , there exists $\beta > b - a$ such that $\ker T_{S^\beta B_{\Lambda_1}} = \{0\}$. Since

$$T_{U\bar{V}}(f) = T_{S^{a-b} B_{\Lambda_1}}(f) = T_{S^\beta S^{\beta+a-b} B_{\Lambda_1}}(f) = T_{S^\beta B_{\Lambda_1}}(f S^{\beta+a-b}), \quad f \in \mathcal{H}^2,$$

we get that $\ker T_{U\bar{V}} = \{0\}$. It thus remain to prove that that U and V satisfy the hypothesis of Theorem 3.1 to get that $\mathcal{M}(U, V) = \{0\}$. So let $m = \arg(U) - \arg(Vb_i) = \arg(B_{\Lambda_1} S^a) - \arg(S^b b_i)$. We argue by contradiction and assume that $m = \tilde{h}$ for some $h \in L_\Pi^1(\mathbb{R})$ and $e^{-h} \in L^1(\mathbb{R})$. Let us choose an $\varepsilon > 0$ such that $b - a + \varepsilon < 2\pi D$. By Lemma 2.2, we know that

$\ker T_{b_i \overline{S^\epsilon}} \cap \mathcal{H}^\infty \neq \{0\}$. Therefore, we use [MP05, Proposition 3.14] to see that $\arg(b_i \overline{S^\epsilon})$ is of the form $-\alpha + \tilde{h}_2$, where α is the argument of a MIF, $h_2 \in L^1_{\Pi}(\mathbb{R})$ and $e^{-h_2} \in L^\infty(\mathbb{R})$. Thus,

$$\begin{aligned} \arg(B_{\Lambda_1} \overline{S^{b-a+\epsilon}}) &= \arg(B_{\Lambda_1} \overline{S^{b-a} b_i \overline{S^\epsilon}}) \\ &= \arg(B_{\Lambda_1} \overline{S^{b-a} b_i}) + \arg(b_i \overline{S^\epsilon}) \\ &= -\alpha + \widetilde{h + h_2}, \end{aligned}$$

where $h+h_2 \in L^1_{\Pi}(\mathbb{R})$ and $e^{-(h+h_2)} \in L^1(\mathbb{R})$. Using [MP05, Proposition 3.14] once more, we have that $\ker T_{B_{\Lambda_1} \overline{S^{b-a+\epsilon}}} \neq \{0\}$, and we get a contradiction between (4) and the fact that $b - a + \epsilon < 2\pi D$.

Example 3.6. Let $U = S^a$ and $V = B_{\Lambda_2} S^b$ with $a > 0$, $b \geq 0$. Let $D := D_*(\Lambda_2)$ and assume that $a - b \neq 2\pi D$. By similar computations as above, we can say that

$$\mathcal{M}(U, V) \neq \{0\} \iff a - b < 2\pi D.$$

Corollary 3.7. Let $\Lambda = \{n + i\}_{n \in \mathbb{Z}}$, $U = S^a$, $V = B_\Lambda$ and assume that $a \neq 2\pi$. Then

$$\mathcal{M}(U, V) \neq \{0\} \iff a < 2\pi.$$

Proof. By Example 2.3, we know that $D_*(\Lambda) = 1$. Thus the conclusion follows from Example 3.6. \square

References

- [Bar00] BARANOV, ANTON D. Differentiation in the Branges spaces and embedding theorems. Nonlinear equations and mathematical analysis. *J. Math. Sci. (New York)* **101** (2000), no. 2, 2881–2913. MR1784683 (2002g:46039), Zbl 1060.46508, doi: 10.1007/BF02672176.
- [Cro94] CROFOOT, R. BRUCE. Multipliers between invariant subspaces of the backward shift. *Pacific J. Math.* **166** (1994), no. 2, 225–246. MR1313454 (95k:47009), Zbl 0819.47042, doi: 10.2140/pjm.1994.166.225.
- [FHR18] FRICAIN, EMMANUEL; HARTMANN, ANDREAS; ROSS, WILLIAM T. Multipliers between model spaces. *Studia Math.* **240** (2018), no. 2, 177–191. MR3720929, arXiv:1605.07418, doi: 10.4064/sm8782-4-2017.
- [GMR16] GARCIA, STEPHAN RAMON; MARTIN, ROBERT T. W.; ROSS, WILLIAM T. Partial orders on partial isometries. *J. Operator Theory* **75** (2016), no. 2, 409–442. MR3509136, Zbl 06658786, arXiv:1501.04888, doi: 10.7900/jot.2015may20.2062.
- [MP05] MAKAROV, NIKOLAI G.; POLTORATSKI, ALEXEI G. Meromorphic inner functions, Toeplitz kernels and the uncertainty principle. *Perspectives in analysis*, 185–252. Math. Phys. Stud., 27. Springer, Berlin, 2005. MR2215727 (2007b:47079), Zbl 1118.47020, doi: 10.1007/3-540-30434-7_10.
- [MP10] MAKAROV, NIKOLAI G.; POLTORATSKI, ALEXEI G. Beurling–Malliavin theory for Toeplitz kernels. *Invent. Math.* **180** (2010), no. 3, 443–480. MR2609247 (2011h:47058), Zbl 1186.47025, arXiv:math/0702497, doi: 10.1007/s00222-010-0234-2.

- [Mas09] MASHREGHI, JAVAD. Representation theorems in Hardy spaces. London Mathematical Society Student Texts, 74. *Cambridge University Press, Cambridge*, 2009. xii+372 pp. ISBN: 978-0-521-73201-7. [MR2500010](#) (2011e:30001), [Zbl 1169.22001](#), doi: [10.1017/CBO9780511814525](#).
- [MiP10] MITKOVSKI, MISHKO; POLTORATSKI, ALEXEI. Pólya sequences, Toeplitz kernels and gap theorems. *Adv. Math.* **224** (2010), no. 3, 1057–1070. [MR2628803](#) (2011c:30076), [Zbl 1204.30018](#), [arXiv:0903.4499](#), doi: [10.1016/j.aim.2009.12.014](#).
- [Nik86] NIKOL'SKIĬ, N.K. Treatise on the shift operator. Spectral function theory. With an appendix by S.V. Khrushchëv and V.V. Peller. Translated from Russian by Jaak Peetre. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 273. *Springer-Verlag, Berlin*, 1986. xii+491 pp. ISBN: 3-540-15021-8. [MR0827223](#) (87i:47042), [Zbl 0587.47036](#), doi: [10.1007/978-3-642-70151-1](#).
- [Pol15] POLTORATSKI, ALEXEI. Toeplitz approach to problems of the uncertainty principle. CBMS Regional Conference Series in Mathematics, 121. *Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI*, 2015. viii+216 pp. ISBN: 978-1-4704-2017-8. [MR3309830](#), [Zbl 1317.47003](#), doi: [/10.1090/cbms/121](#).
- [Tim16] TIMOTIN, DAN. On a preorder relation for contractions. *Acta Sci. Math. (Szeged)* **82** (2016), no. 3–4, 629–640. [MR3616198](#), [Zbl 06707925](#), [arXiv:1507.07128](#), doi: [10.14232/actasm-015-068-5](#).

(Emmanuel Fricain) LABORATOIRE PAUL PAINLEVÉ, UNIVERSITÉ LILLE 1, 59 655 VILLENEUVE D'ASCQ CÉDEX

emmanuel.fricain@math.univ-lille1.fr

(Rishika Rupam) LABORATOIRE PAUL PAINLEVÉ, UNIVERSITÉ LILLE 1, 59 655 VILLENEUVE D'ASCQ CÉDEX

rishika.rupam@math.univ-lille1.fr

This paper is available via <http://nyjm.albany.edu/j/2018/24-6.html>.