

# The eigensheaf of an operator

Sameer Chavan and Archana Morje

ABSTRACT. The eigensheaf  $\mathcal{F}_T$  of an operator  $T$  on a Hilbert space  $H$  is the subsheaf of  $\mathcal{O}_H$  defined by the prescription:

$$\mathcal{F}_T(U) = \{f \in \mathcal{O}_H(U) \mid T(f(w)) = wf(w) \text{ for all } w \in U\},$$

where  $U$  is open in  $\mathbb{C}$  and  $\mathcal{O}_H$  is the sheaf of  $H$ -valued holomorphic functions defined on  $\mathbb{C}$ . If  $T$  lies in the Cowen-Douglas class, then its eigensheaf is locally free, but not conversely. We obtain a model for operators whose eigensheaves are locally free. We describe the eigensheaves for certain coanalytic Toeplitz operators, we show that the map from an operator to its eigensheaf is a functor from the category of bounded linear operators on Hilbert space to the category of Hilbert space-valued analytic sheaves, and we discuss relation between the eigensheaf of an operator and the sheaf that Putinar associates to an operator.

## CONTENTS

1. The Cowen-Douglas correspondence	868
2. The eigensheaf of an operator	870
3. Toeplitz operators with coanalytic symbols	876
4. Relation with the Putinar's sheaf	879
Acknowledgment	881
References	881

## 1. The Cowen-Douglas correspondence

This article is partly motivated by the celebrated work [9] of Cowen and Douglas. In that paper, they introduced and studied a class of bounded linear Hilbert space operators with point spectrum containing an open set (now well-known as the Cowen-Douglas class  $\mathbf{B}_n(\Omega)$ ). One of the main results in [9] proves that the classification of operators in  $\mathbf{B}_n(\Omega)$  is equivalent to the classification of a class of holomorphic vector bundles associated to the operators in  $\mathbf{B}_n(\Omega)$ . Moreover, the present paper is very much in the spirit of the Putinar's

Received September 8, 2021.

2020 *Mathematics Subject Classification*. Primary 47B13, 47B35; Secondary 55Rxx.

*Key words and phrases*. Cowen-Douglas class, Toeplitz operator, holomorphic vector bundle, locally free, cokernel sheaf.

The work of the second author was supported through UGC SAP (DSA 1).

work [26, 27, 28] on a sheaf associated to a bounded linear operator. In particular, we associate the class of analytic eigensheaves to the class of bounded linear Hilbert space operators (see Definition 2.1 below). Needless to say, it is different from the sheaf constructions appearing in [4, 12, 13, 26, 27, 28]. Indeed, the sheaf of this paper is a kernel sheaf while the one introduced and studied in [26] is a cokernel sheaf (see Section 4).

All the vector spaces here are over the field  $\mathbb{C}$  of complex numbers. If  $E$  is a complex vector space, then  $\mathbf{1}_E$  denotes the identity operator on  $E$ , and for every complex number  $w$ , we denote the linear operator  $w\mathbf{1}_E : E \rightarrow E$  also by  $w$ . Throughout this paper,  $\Omega$  stands for a bounded connected open subset of  $\mathbb{C}$  and  $\Omega^*$  denotes the set of complex conjugates of elements in  $\Omega$ . For a complex Hilbert space, the space of bounded linear operators on  $H$  is denoted by  $\mathcal{L}(H)$ . For  $T \in \mathcal{L}(H)$ , the kernel, the point spectrum, the approximate point spectrum and the essential spectrum of  $T$  are denoted by  $\text{Ker}(T)$ ,  $\sigma_p(T)$ ,  $\sigma_{ap}(T)$  and  $\sigma_e(T)$ , respectively.

We now recall the definition of the Cowen-Douglas class (the reader is referred to [23] for an up-to-date exposition on this class). For a connected open subset  $\Omega$  of  $\mathbb{C}$  and  $n$  a positive integer, let  $\mathbf{B}_n(\Omega)$  denote the collection of all operators  $T \in \mathcal{L}(H)$  satisfying the following conditions :

- for every point  $w \in \Omega$ ,  $(T - w)(H) = H$  and  $\dim \text{Ker}(T - w) = n$ ,
- the subspace  $\sum_{w \in \Omega} \text{Ker}(T - w)$  of  $H$  equals  $H$ , where  $\sum$  stands for the closed linear span.

We call  $\mathbf{B}_n(\Omega)$  the *Cowen-Douglas class of degree  $n$  on  $H$  with respect to  $\Omega$* , and its elements as the *Cowen-Douglas operators of degree  $n$  on  $H$  with respect to  $\Omega$* .

The following is essentially [9, Proposition 1.11 and Corollary 1.12].

**Proposition 1.1.** *Let  $n$  be a positive integer, and  $T$  be a Cowen-Douglas operator of degree  $n$  on  $H$  with respect to  $\Omega$ . Let*

$$E_T := \{(w, x) \in \Omega \times H \mid Tx = wx\}, \tag{1}$$

and let  $\pi_T : E_T \rightarrow \Omega$  be the restriction of the canonical projection from  $\Omega \times H$  to  $\Omega$ . Then, there is a natural structure of a complex manifold on  $E_T$  such that the map  $\pi_T : E_T \rightarrow \Omega$  is a holomorphic vector bundle of rank  $n$ . Moreover, the function  $h_T : E_T \times_{\Omega} E_T \rightarrow \mathbb{C}$  given by

$$h_T((w, x), (w, x')) = \langle x, x' \rangle, \quad w \in E_T, x, x' \in H,$$

is a smooth Hermitian metric on this holomorphic vector bundle.

In view of the last result, one can define a map from the category of Cowen-Douglas class over an open connected subset  $\Omega$  of  $\mathbb{C}$  to the category of holomorphic vector bundles over  $\Omega$  (see [9, Theorem 1.14]). It has been recorded in [3, Pg 1] that this correspondence is not essentially surjective (see [8, Section 0.3]; see also Example 3.5 below). On the other hand, there are many coanalytic Toeplitz operators belonging to  $\mathbf{B}_m(\Omega')$  for some  $m$  and for some component  $\Omega'$  of  $\Omega \setminus \sigma_e(T)$ . Also, the degree  $m$  may vary for different components of

$\Omega \setminus \sigma_e(T)$ . The structure of these operators can be revealed by analysing their eigensheaves, and in turn, providing another motivation for the present work.

Here is the plan of this paper. In Section 2, we discuss the idea of the eigensheaf  $\mathcal{F}_T$  of an arbitrary bounded linear Hilbert space operator  $T$  and observe that the support of  $\mathcal{F}_T$  is always contained in the closure of the point spectrum of  $T$  (see Proposition 2.5). We also discuss the relation between the interior of the point spectrum of  $T$  and the support of the sheaf  $\mathcal{F}_T$  (see Example 2.7). We show that if  $T$  is hyponormal on a separable Hilbert space, then  $\mathcal{F}_T$  is necessarily trivial (see Corollary 2.8). Further, we investigate the class of operators with locally free eigensheaves and obtain a model for these operators (see Theorem 2.9). In Section 3, we exhibit a class of locally free analytic eigensheaves arising from certain coanalytic Toeplitz operators on the Hardy space of the unit disc (see Theorem 3.3). In Section 4, we discuss the relation between the Putinar's sheaf and the one investigated in this paper (see Proposition 4.1).

## 2. The eigensheaf of an operator

Let  $M$  be a complex manifold and let  $X$  be a complex Banach space. We say that a map  $f : M \rightarrow X$  is  $X$ -valued holomorphic if the function  $\alpha \circ f : M \rightarrow \mathbb{C}$  is holomorphic for every continuous linear functional  $\alpha : X \rightarrow \mathbb{C}$ . Note that if  $H$  is a separable complex Hilbert space and  $\{e_n\}_{n \geq 0}$  is an orthonormal basis for  $H$ , then  $f : M \rightarrow H$  is holomorphic if and only if  $\langle f(\cdot), e_n \rangle : M \rightarrow \mathbb{C}$  is holomorphic for every integer  $n \geq 0$  (refer to [2] for the basic theory of vector-valued holomorphic functions). Consider the sheaf  $\mathcal{O}_{M,H}$  of  $H$ -valued holomorphic functions on  $M$ . This sheaf  $\mathcal{O}_{M,H}$  has a canonical structure of an  $\mathcal{O}_M$ -module, where  $\mathcal{O}_M$  is the sheaf of  $\mathbb{C}$ -valued holomorphic functions on  $M$ . If  $M = \mathbb{C}$ , then  $\mathcal{O}_{M,H}$  is denoted by  $\mathcal{O}_H$ . The restriction of a sheaf  $\mathcal{F}$  to a non-empty open subset  $\Omega$  is denoted by  $\mathcal{F}|_\Omega$  (refer to [31, Chapter 2] for the definition of sheaf and related notions).

The idea of associating a sheaf with a bounded linear operator first appears in the work [26]. This provides a novel approach to the local spectral theory (see [12, 27, 28, 13]; refer also to [4, Section 1.3] for discussion on a coherent sheaf associated to the Cowen-Douglas operators of degree 1).

**Definition 2.1.** For  $T \in \mathcal{L}(H)$ , define an  $\mathcal{O}_\mathbb{C}$ -submodule  $\mathcal{F}_T$  of  $\mathcal{O}_H$  by

$$\mathcal{F}_T(U) = \{f \in \mathcal{O}_H(U) \mid T(f(w)) = wf(w) \text{ for all } w \in U\}$$

for every open subset  $U$  of  $\mathbb{C}$ . We will call  $\mathcal{F}_T$  the *eigensheaf* of  $T$ .

*Remark 2.2.* For open subsets  $U, V$  of  $\mathbb{C}$  such that  $V \subseteq U$ , the restriction morphism  $\text{res}_{U,V} : \mathcal{F}_T(U) \rightarrow \mathcal{F}_T(V)$  is given by

$$\text{res}_{U,V}(f) = f|_V, \quad f \in \mathcal{F}_T(U).$$

Clearly, for every open set  $U$  of  $\mathbb{C}$ ,  $\text{res}_{U,U}$  is the identity morphism on  $\mathcal{F}_T(U)$ . Also, for open subsets  $U, V, W$  of  $\mathbb{C}$  such that  $W \subseteq V \subseteq U$ ,  $\text{res}_{W,V} \circ \text{res}_{V,U} = \text{res}_{W,U}$ . Thus  $\mathcal{F}_T$  is a presheaf. If  $\{\mathcal{U}_i\}_{i \in I}$  is an open covering of an open set  $U$ , and if  $f, g \in \mathcal{F}_T(U)$  have the property  $f|_{\mathcal{U}_i} = g|_{\mathcal{U}_i}$  for each  $i \in I$ , then

$f = g$ . Moreover, if  $\{U_i\}_{i \in I}$  is an open covering of an open set  $U$ , and if for each  $i \in I$  and  $f_i \in \mathcal{F}_T(U_i)$  is given such that for each pair  $U_i, U_j$  of the covering sets  $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$ , then  $f = f_i$  on  $U_i, i \in I$ , defines a section  $f \in \mathcal{F}_T(U)$ . Thus  $\mathcal{F}_T$  is a sheaf. Finally, note that one can define the eigensheaf of a continuous linear operator on any Frechet space (recall that if  $X$  is a Frechet space, then  $\mathcal{O}_X$  is the projective tensor product  $\mathcal{O}_\mathbb{C} \hat{\otimes} X$ ; see [13, Appendix 1]).

Let  $\mathcal{B}$  denote the category of bounded linear Hilbert space operators. If  $T : H \rightarrow H$  and  $S : K \rightarrow K$  are two bounded linear operators then

$$\text{Hom}_{\mathcal{B}}(T, S) = \{u : H \rightarrow K : u \circ T = S \circ u\}.$$

Let  $\mathcal{O}_\mathbb{C}\text{-mod}$  denote the category of  $\mathcal{O}_\mathbb{C}$ -module over  $\mathbb{C}$  and  $\text{Hom}_{\mathcal{O}_\mathbb{C}}(\mathcal{F}_1, \mathcal{F}_2)$  denote the set of  $\mathcal{O}_\mathbb{C}$ -module homomorphisms between the  $\mathcal{O}_\mathbb{C}$ -modules  $\mathcal{F}_1$  and  $\mathcal{F}_2$ .

**Proposition 2.3.** *Consider the category  $\mathcal{B}$  of bounded linear Hilbert space operators and the category  $\mathcal{O}_\mathbb{C}\text{-mod}$  of  $\mathcal{O}_\mathbb{C}$ -module over  $\mathbb{C}$ . Define  $F(T) = \mathcal{F}_T$  for  $T \in \mathcal{B}$ . For  $u \in \text{Hom}_{\mathcal{B}}(T, S)$ , define  $F(u)$  as the  $\mathcal{O}_\mathbb{C}$ -module homomorphism  $F(u) : \mathcal{F}_T \rightarrow \mathcal{F}_S$  given by*

$$F(u)_U : \mathcal{F}_T(U) \rightarrow \mathcal{F}_S(U), \quad F(u)_U(f) = u \circ f$$

for every open subset  $U \subseteq \mathbb{C}$ . Then  $F$  defines a functor.

**Proof.** Note that  $u \circ f \in \mathcal{F}_S(U)$  for every  $u \in \text{Hom}_{\mathcal{B}}(T, S)$ . Indeed, since  $f \in \mathcal{F}_T(U)$ ,  $T(f(w)) = wf(w)$  for all  $w \in U$ , and hence

$$S(u \circ f(w)) = (S \circ u)(f(w)) = u \circ T(f(w)) = u(w(f(w))) = w(u \circ f)(w).$$

It is easy to see that  $F(u)$  is a  $\mathcal{O}_\mathbb{C}$ -module homomorphism. Since  $F(\mathbf{1}_H) = \mathbf{1}_{\mathcal{F}_T}$  (note that  $\mathbf{1}_H$  is the identity morphism of  $T$  in the category  $\mathcal{B}$ ) and  $F(u \circ v) = F(u) \circ F(v)$  for every  $u \in \text{Hom}_{\mathcal{B}}(S, T)$  and  $v \in \text{Hom}_{\mathcal{B}}(T, R)$ ,  $F$  is a functor.  $\square$

Since  $F$  is a functor, if  $u$  is a Hilbert space isomorphism then  $F(u)$  is a  $\mathcal{O}_\mathbb{C}$ -module isomorphism. Let us see a particular instance in which the last proposition is applicable. Consider the subspace  $K$  of  $H$  given by

$$K = \sum_{w \in \mathbb{C}} \text{Ker}(T - w).$$

Define  $S : K \rightarrow K$  be  $S = T|_K$ . Clearly,  $S$  is a bounded linear operator on  $K$ . Further,  $T \circ u = u \circ S$ , where  $u : K \hookrightarrow H$  denotes the inclusion map. This yields a natural  $\mathcal{O}_\mathbb{C}$ -module homomorphism  $\Phi : \mathcal{F}_T \rightarrow \mathcal{F}_S$ . Moreover, since  $j$  is injective, so is  $\Phi$ . Note that the above observation is applicable to any closed  $T$ -invariant subspace  $K$  of  $H$ .

By the correspondence between vector bundles and locally free sheaves (see [31]), the eigensheaf of a Cowen-Douglas operator of degree  $n$  is locally free sheaf of rank  $n$  (see [24, Definition 2.1.35]). This yields an analogue of Proposition 1.1 formulated in the language of eigensheaves. Recall that an  $\mathcal{O}_X$ -module  $\mathcal{F}$  is *locally free* if for every point  $x$  in a complex manifold  $X$ , there exist an open neighborhood  $U$  of  $x$  and a set  $I_x$  such that  $\mathcal{F}|_U$  is isomorphic to the direct sum

$\mathcal{O}_X^{(I_x)}|_U$  as an  $\mathcal{O}_X|_U$ -module. For a nonnegative integer  $r$ , we say  $\mathcal{F}$  is *locally free of rank  $r$*  if for every  $x \in X$ , the index set  $I_x$  has the cardinality  $r$ .

**Proposition 2.4.** *Let  $\Omega$  be a connected open subset of  $\mathbb{C}$ ,  $n$  be a positive integer, and  $T$  be a Cowen-Douglas operator of degree  $n$  on  $H$  with respect to  $\Omega$ . Then the eigensheaf  $\mathcal{F}_T|_\Omega$  of  $T$  is locally free of rank  $n$ .*

**Proof.** Note that the  $\mathcal{O}_\Omega$ -module  $\mathcal{F}_T|_\Omega$  is the analytic sheaf of holomorphic sections of the holomorphic vector bundle  $\pi_T : E_T \rightarrow \Omega$  (see (1)), and hence by Proposition 1.1 and the preceding discussion, the eigensheaf  $\mathcal{F}_T|_\Omega$  is locally free of rank  $n$ .  $\square$

Note that  $\mathcal{F}_T$  is an  $\mathcal{O}_\mathbb{C}$ -submodule of  $\mathcal{O}_H$  obtained by imposing the condition on a section  $f$  that  $w$  is an eigenvalue of  $T$  whenever its evaluation  $f(w)$  at  $w$  is non-zero. This motivates one to explore the relation between the set of eigenvalues of  $T$  and the support of  $\mathcal{F}_T$ . Recall that if  $\mathcal{F}$  denotes a sheaf of abelian groups on a topological space  $X$ , then the *support of  $\mathcal{F}$*  is given by

$$\text{Supp}(\mathcal{F}) = \{x \in X \mid \mathcal{F}_x \neq 0\},$$

where  $\mathcal{F}_x$  stands for the stalk of  $\mathcal{F}$  at  $x$ . The following result describes the support of  $\mathcal{F}_T$  (cf. [26, Lemma 2.1] and Corollary 4.3 below).

**Proposition 2.5.** *For  $T \in \mathcal{L}(H)$ , let  $\mathcal{F}_T$  be the eigensheaf of  $T$ . Then the following statements are valid:*

(i) *Supp( $\mathcal{F}_T$ ) is an open subset of  $\mathbb{C}$  such that*

$$\text{Supp}(\mathcal{F}_T) \subseteq \overline{\sigma_p(T)}, \quad (2)$$

- (ii) *for every open neighbourhood  $V$  of  $z_0 \in \text{Supp}(\mathcal{F}_T)$ , there exists an open subset  $W$  of  $V$  contained in  $\sigma_p(T)$ ,*  
 (iii) *if  $\text{Supp}(\mathcal{F}_T) \neq \emptyset$ , then  $\sigma_p(T)$  has non-empty interior,*  
 (iv) *if there exists an open connected subset  $U$  of  $\mathbb{C}$  such that  $\mathcal{F}_T(U) \neq \{0\}$ , then  $U \subseteq \overline{\sigma_p(T)}$ .*

*In particular, the support of  $\mathcal{F}_T$  is contained in the approximate point spectrum of  $T$ .*

**Proof.** Let  $z_0 \in \text{Supp}(\mathcal{F}_T)$ . Then,  $(\mathcal{F}_T)_{z_0} \neq 0$ , so there is a nonzero element  $\gamma$  of  $(\mathcal{F}_T)_{z_0}$ . Choose a connected open neighbourhood  $U$  of  $z_0$  and a section  $f$  of  $\mathcal{F}_T(U)$  such that  $\gamma = f_{z_0}$ . Then,  $f_z \neq 0$  for all  $z \in U$ . For, if  $f_z = 0$  for some  $z \in U$ , then  $f$  vanishes on an open neighbourhood of  $z$  in  $U$ , and hence by the vector-valued identity theorem (see [2, Proposition A.2]),  $f = 0$ , which is not possible since  $f_{z_0} \neq 0$ . In particular,  $(\mathcal{F}_T)_z \neq 0$  for all  $z \in U$ , and hence  $U \subseteq \text{Supp}(\mathcal{F}_T)$ . This also shows that if  $V$  is any neighbourhood of  $z_0$ , then there exist an open neighbourhood  $U$  of  $z_0$  and a point  $z' \in U$  such that  $f(z') \neq 0$ . By the continuity of  $f$ ,  $f$  is non-vanishing in some neighborhood  $W \subseteq V$  of  $z'$ . As  $f \in \mathcal{F}_T(U)$ , we have  $(T - w')(f(w)) = 0$  for every  $w \in W$ . Thus,  $W \subseteq \overline{\sigma_p(T)}$ . This yields (ii) and also shows that  $z_0 \in \overline{\sigma_p(T)}$ , completing the proof of (i).

Clearly, (iii) follows from (ii). To see (iv), let  $f : U \rightarrow H$  be a non-zero element of  $\mathcal{F}_T(U)$ . If  $Z(f)$  denotes the zero set of  $f$ , then  $U \setminus Z(f) \subseteq \sigma_p(T)$ , and by the identity theorem,  $U \setminus Z(f)$  is dense in  $U$ . It follows that

$$U = \overline{U \setminus Z(f)}^U \subseteq \overline{U \setminus Z(f)} \subseteq \overline{\sigma_p(T)},$$

where  $\overline{W}^U$  denotes the closure of the subset  $W$  of  $U$  in  $U$ . Finally, since  $\sigma_p(T) \subseteq \sigma_{ap}(T)$  and  $\sigma_{ap}(T)$  is closed in  $\mathbb{C}$ , the remaining part is immediate from (2).  $\square$

*Remark 2.6.* If  $\mathcal{F}_T|_U$  is of finite type, then its support is closed in  $U$  (see [16, Chapter 0, Section 5.2.2], [24, Corollary 2.1.17]).

In general, the point spectrum of  $T$  need not be contained in the support of  $\mathcal{F}_T$ . For instance, if  $T$  is a diagonal operator with diagonal entries  $\{1/n\}_{n \geq 1}$  then  $\sigma_p(T) \neq \emptyset$ . However, by Proposition 2.5(iii),  $\mathcal{F}_T = \{0\}$ . Although equality does not hold in (2) in general, the natural question arises is whether one can recover the support of  $\mathcal{F}_T$  from the point spectrum  $\sigma_p(T)$  of  $T$ ? By Proposition 2.5(iii), if the interior  $\sigma_p(T)^\circ$  of  $\sigma_p(T)$  is empty then the support of  $\mathcal{F}_T$  is also empty. So perhaps the correct guess would be the following inclusion:

$$\sigma_p(T)^\circ \subseteq \text{Supp}(\mathcal{F}_T). \tag{3}$$

This certainly holds if eigenfunctions of  $T$  are holomorphic functions on the interior of point spectrum (for instance, this happens for operators  $T$  in the Cowen-Douglas class). Here is a slightly different example (cf. Theorem 3.3).

*Example 2.7.* Consider the positive definite kernel  $\kappa(z, w)$  given by

$$\kappa(z, w) = \frac{(z^2 + \bar{z} + 2)(\bar{w}^2 + w + 2)}{1 - z\bar{w}}, \quad z, w \in \mathbb{D}.$$

By [25, Theorem 5.21],  $z$  is a multiplier of the reproducing kernel Hilbert space  $H_\kappa$  associated with  $\kappa$ . Let  $M_z$  denote the operator of multiplication by  $z$  on  $H_\kappa$ . Then any  $w$  in the open unit disc  $\mathbb{D}$  is an eigenvalue of  $T := M_z^*$  with corresponding eigenfunction  $\kappa(\cdot, \bar{w}) \in H_\kappa$ , which is clearly not holomorphic in  $w$ . Still, the holomorphic section  $w \mapsto \frac{z^2 + \bar{z} + 2}{1 - zw}$  belongs to  $\mathcal{F}_T|_{\mathbb{D}}$ , and hence the inclusion (3) holds.  $\diamond$

For the definition of the single-valued extension property (for short, SVEP), see [1, Definition 2.3]. The following is certainly well-known. Indeed, it is a consequence of [1, Theorem 3.96] and the discussion prior to it.

**Corollary 2.8.** *Let  $H$  be a separable Hilbert space. For  $T \in \mathcal{L}(H)$ , let  $\mathcal{F}_T$  be the eigensheaf of  $T$ . If  $T$  is hyponormal, that is,  $T^*T - TT^* \geq 0$ , then  $\mathcal{F}_T = \{0\}$ . In particular, a hyponormal operator on a separable Hilbert space has SVEP.*

**Proof.** Recall the well-known fact about a hyponormal operator that its eigenvectors corresponding to distinct eigenvalues are orthogonal [21]. Since  $H$  is separable,  $T$  can have at most countable distinct eigenvalues. The desired conclusion now follows from Proposition 2.5(iii). The remaining part now follows from the fact that  $T$  has the SVEP if and only if  $\mathcal{F}_T$  is trivial.  $\square$

In the remaining part of this section, we study operators with locally free eigensheaves. The following result sheds some light on the relation between Cowen-Douglas operators and operators with locally free eigensheaves over connected domains. The later one is generically an extension of the adjoint of the multiplication operator  $M_z$  on a reproducing kernel Hilbert space of vector-valued homomorphic functions. We make this statement precise in the following theorem. The proof presented below is an adaptation of the method of modelling Cowen-Douglas class to the present situation (see [9, Pg 194] for Cowen-Douglas class of degree 1, and [10, Theorem 4.12], [33, Theorem B], [14, Pg 279-280] for the general case).

**Theorem 2.9.** *Let  $\Omega$  be a bounded open connected subset of  $\mathbb{C}$  and let  $T \in \mathcal{L}(H)$ . If the eigensheaf  $\mathcal{F}_T|_{\Omega}$  of  $T$  is locally free of rank  $n$ , then there exists a reproducing kernel Hilbert space  $\mathcal{H}$  of holomorphic  $\mathbb{C}^n$ -valued functions on  $\Omega^*$  and a surjective partial isometry  $V : H \rightarrow \mathcal{H}$  such that  $VT^* = M_zV$ , where  $M_z$  denotes the operator of multiplication by the coordinate function  $z$  acting on  $\mathcal{H}$ .*

**Proof.** Suppose that  $\mathcal{F}_T|_{\Omega}$  is locally free of rank  $n$ . By a theorem of Grauert, any holomorphic vector bundle over an open subset of  $\mathbb{C}$  is holomorphically trivial, and hence it admits a global holomorphic frame ([15, Theorem 30.4]). Let  $\{f_1, \dots, f_n\}$  denote a global holomorphic frame for the holomorphic vector bundle associated with  $\mathcal{F}_T|_{\Omega}$ . Define  $V : H \rightarrow \mathcal{O}(\Omega^*, \mathbb{C}^n)$  by

$$(Vh)(w) = (\langle h, f_1(\bar{w}) \rangle, \dots, \langle h, f_n(\bar{w}) \rangle), \quad h \in H, w \in \Omega^*.$$

Note that  $Vh = 0$  if and only if  $h$  is orthogonal to

$$K := \sum_{w \in \Omega} \text{Ker}(T - w) = \sum \{f_j(w) : w \in \Omega, j = 1, \dots, n\}.$$

Thus  $\text{Ker}(V) = H \ominus K$ . Moreover, for any  $h \in H$  and  $w \in \Omega^*$ ,

$$\begin{aligned} (VT^*h)(w) &= (\langle T^*h, f_1(\bar{w}) \rangle, \dots, \langle T^*h, f_n(\bar{w}) \rangle) \\ &= (\langle h, \bar{w}f_1(\bar{w}) \rangle, \dots, \langle h, \bar{w}f_n(\bar{w}) \rangle) \\ &= (M_zVh)(w), \end{aligned}$$

where  $M_z : \mathcal{O}(\Omega^*, \mathbb{C}^n) \rightarrow \mathcal{O}(\Omega^*, \mathbb{C}^n)$  is given by  $(M_zF)(w) = wF(w)$ ,  $F \in \mathcal{O}(\Omega^*, \mathbb{C}^n)$ . Consider the complex vector space  $\mathcal{H} := \{Vh : h \in H\}$  endowed with the inner product

$$\langle Vh, Vh' \rangle_{\mathcal{H}} = \langle P_K h, P_K h' \rangle_H, \quad h, h' \in H, \quad (4)$$

where  $P_K$  denotes the orthogonal projection of  $H$  onto  $K$ . Thus  $\tilde{V} : H \rightarrow \mathcal{H}$  given by  $\tilde{V}h = Vh$ ,  $h \in H$ , is a partial isometry with the initial space  $K$  and the final space  $\mathcal{H}$ . Therefore  $\mathcal{H}$  is a Hilbert space with the natural inner product induced by the norm  $\|\cdot\|_{\mathcal{H}}$ . The intertwining relation  $VT^* = M_zV$  shows that  $M_z$  is a bounded linear operator on  $\mathcal{H}$ . It now suffices to check that  $\mathcal{H}$  is a reproducing kernel Hilbert space. To see that, let  $L(\mathbb{C}^n, K)$  denote the vector

space of linear operators from  $\mathbb{C}^n$  into  $K$ , and define  $\gamma : \Omega^* \rightarrow L(\mathbb{C}^n, K)$  by

$$\gamma(w)(\alpha) = \sum_{j=1}^n \alpha_j f_j(\bar{w}), \quad \alpha = (\alpha_1, \dots, \alpha_n), \quad w \in \Omega^*.$$

Consider the positive definite kernel  $\kappa : \Omega^* \times \Omega^* \rightarrow L(\mathbb{C}^n)$  given by  $\kappa(z, w) = \gamma(w)^* \gamma(z)$ . It is easy to see that  $\gamma(w)^* h = (Vh)(w)$  for every  $w \in \Omega^*$  and  $h \in K$ . It now follows from (4) that for  $h \in H$ ,  $w \in \Omega^*$  and  $\alpha \in \mathbb{C}^n$ ,

$$\begin{aligned} \langle Vh, \kappa(\cdot, w)(\alpha) \rangle_{\mathcal{H}} &= \langle Vh, \gamma(\cdot)^* \gamma(w)(\alpha) \rangle_{\mathcal{H}} \\ &= \langle Vh, V(\gamma(w)(\alpha)) \rangle_{\mathcal{H}} \\ &= \langle P_K h, P_K \gamma(w)(\alpha) \rangle_H \\ &= \langle (Vh)(w), \alpha \rangle_{\mathbb{C}^n}. \end{aligned}$$

This completes the proof. □

We conclude the section with a brief discussion on the class  $\mathbf{S}_{m,n}(\Omega_1, \Omega_2)$  of operators  $T$  for which  $\mathcal{F}_T|_{\Omega_1}$  and  $\mathcal{F}_{T^*}|_{\Omega_2}$  are locally free of ranks  $m$  and  $n$  respectively, where  $\Omega_1, \Omega_2$  are bounded, open connected subsets of  $\mathbb{C}$ . This class appears to be closely related to the class  $\mathbf{B}_{m,n}(\Omega_1, \Omega_2)$  as investigated by M. Cowen in [8]. To see this, let  $T$  be an operator on  $H$  in  $\mathbf{S}_{m,n}(\Omega_1, \Omega_2)$ , and consider closed subspaces  $K_1, K_2$  of  $H$  given by

$$K_1 := \sum_{w_1 \in \Omega_1} \text{Ker}(T - w_1), \quad K_2 := \sum_{w_2 \in \Omega_2} \text{Ker}(T^* - w_2).$$

If  $H = \sum(K_1 \cup K_2)$ , then  $T$  admits the decomposition

$$T = \begin{pmatrix} T_1 & X \\ 0 & T_2 \end{pmatrix} \text{ on } H = K_1 \oplus K_2,$$

where  $T_1, T_2$  are bounded linear operators on  $K_1, K_2$  such that  $\mathcal{F}_{T_1}|_{\Omega_1}, \mathcal{F}_{T_2^*}|_{\Omega_2}$  are locally free of ranks  $m, n$  respectively, and  $X$  is a bounded linear transformation from  $K_1$  into  $K_2$  (cf. [8, Proposition 1.6]). The fact that  $K_1$  and  $K_2$  are mutually orthogonal follows from the following general fact.

**Proposition 2.10.** *For  $T \in \mathcal{L}(H)$ , let  $\mathcal{F}_T$  be the eigensheaf of  $T$ . If  $U, V$  are open subsets of  $\mathbb{C}$ , then for every  $f \in \mathcal{F}_T(U)$  and  $g \in \mathcal{F}_{T^*}(V)$ , the map  $\phi : U \times V \rightarrow \mathbb{C}$  given by*

$$\phi(u, v) = \langle f(u), g(v) \rangle, \quad u \in U, v \in V,$$

*is identically 0.*

**Proof.** For  $u \in U$  and  $v \in V$ , we have

$$u \langle f(u), g(v) \rangle = \langle Tf(u), g(v) \rangle = \langle f(u), T^*g(v) \rangle = \bar{v} \langle f(u), g(v) \rangle.$$

Thus  $\langle f(u), g(v) \rangle = 0$  whenever  $u \neq \bar{v}$ . By continuity of  $f$  and  $g$ , we get the same conclusion for every  $u \in U$  and  $v \in V$ . □

The classification of operators with locally free eigensheaves (in the spirit of [9, Theorem 1.14]) seems to be beyond reach at present. In this context, we conclude from Theorem 2.9 that any operator with locally free eigensheaf over a connected open set is necessarily an extension of the adjoint of multiplication operator  $M_z$  in a reproducing kernel Hilbert space.

### 3. Toeplitz operators with coanalytic symbols

The class of Toeplitz operators has been extensively studied in the literature, particularly, in the context of Cowen-Douglas theory (see [19, 20, 18, 32, 17]). It is worth noting that Cowen-Douglas class does not support Toeplitz operators with analytic symbols. Indeed, as a consequence of Corollary 2.8, one may conclude that the eigensheaf of the multiplication operator  $M_\phi$  on Hardy space  $H^2(\mathbb{D})$  of the unit disc with analytic symbol  $\phi$  is trivial. Recall that  $H^2(\mathbb{D})$  is the Hilbert space of holomorphic functions  $f(z) = \sum_{k=0}^{\infty} \hat{f}(k)z^k$  on the open unit disc  $\mathbb{D}$  endowed with the norm

$$\|f\| = \left( \sum_{k=0}^{\infty} |\hat{f}(k)|^2 \right)^{1/2}, \quad f \in H^2(\mathbb{D}).$$

It turns out that  $H^2(\mathbb{D})$  is a reproducing kernel Hilbert space with reproducing kernel  $\kappa$  given by

$$\kappa(z, w) = \frac{1}{1 - z\bar{w}}, \quad z, w \in \mathbb{D}. \quad (5)$$

This means that

$$f(w) = \langle f, \kappa(\cdot, w) \rangle, \quad w \in \mathbb{D}, f \in H^2(\mathbb{D}). \quad (6)$$

Recall further that for  $\phi \in H^\infty(\mathbb{D})$  (the space of bounded holomorphic functions on  $\mathbb{D}$ ), the multiplication (bounded linear) operator  $M_\phi$  is defined by

$$(M_\phi f)(z) = \phi(z)f(z), \quad f \in H^2(\mathbb{D}), z \in \mathbb{D}.$$

The Hardy space  $H^2(\mathbb{D})$  can be identified with the the closed subspace  $H^2(\mathbb{T}) = \{f \in L^2(\mathbb{T}) : \hat{f}(n) = 0 \text{ if } n < 0\}$  of  $L^2(\mathbb{T})$  by associating every  $f \in H^2(\mathbb{D})$  to its boundary function  $\tilde{f}$ . Here  $\mathbb{T}$  denotes the unit circle in  $\mathbb{C}$ . The Toeplitz operator  $T_\phi$  on  $H^2(\mathbb{T})$  with symbol  $\phi \in C(\mathbb{T})$  is defined by  $T_\phi = PN_\phi|_{H^2(\mathbb{T})}$ , where  $N_\phi$  denotes the operator of multiplication by  $\phi$  in  $L^2(\mathbb{T})$  and  $P$  is the orthogonal projection of  $L^2(\mathbb{T})$  onto  $H^2(\mathbb{T})$ . It is easy to see that if  $\phi \in H^\infty(\mathbb{D})$  then  $M_\phi$  is unitarily equivalent to  $T_{\tilde{\phi}}$  (refer to [22] for the basics of Toeplitz operators).

In this section, we describe the analytic sheaf  $\mathcal{F}_T$  for a class of Toeplitz operators  $T$  with coanalytic symbols (cf. [6, Theorem 1]). We begin with a couple of lemmas. The first of which is well-known (see, for instance, [11, Theorem 7.26] and [30, Theorem 10]).

**Lemma 3.1.** *For  $\phi \in H^\infty(\mathbb{D})$  with boundary function  $\tilde{\phi}$  on  $\mathbb{T}$ , the multiplication operator  $M_\phi$  (or equivalently  $M_\phi^*$ ) is Fredholm if and only if there exists  $M > 0$*

such that  $|\tilde{\phi}| > M$  almost everywhere on  $\mathbb{T}$  and  $\phi$  has finitely many zeros inside  $\mathbb{D}$ . In this case, the Fredholm index is given by

$$\text{ind } M_{\phi}^* = \dim \text{Ker}(M_{\phi}^*) = \text{the number of zeros of } \phi \text{ inside } \mathbb{D}.$$

The following lemma appears to be known (see [29, 9]). Since we could not locate it in the form we need in the sequel, we include it.

**Lemma 3.2.** *Let  $U$  be a bounded open subset of  $\mathbb{C}$  and let  $T \in \mathcal{L}(H)$ . If the range of  $T - w$  is closed for every  $w \in U$ , then  $\mathcal{F}_T|_U$  is locally free of rank  $n$  if and only if*

$$\dim \text{Ker}(T - w) = n \text{ for every } w \in U. \tag{7}$$

**Proof.** Suppose that the range of  $T - w$  is closed for every  $w \in U$ . If  $\mathcal{F}_T|_U$  is locally free of rank  $n$ , then  $U \subseteq \text{Supp}(\mathcal{F}_T)$ , and hence we obtain (7). To see the sufficiency part, assume that (7) holds. Note that  $T - w$  is left Fredholm for every  $w \in U$ . One may now argue as in the proof of [9, Proposition 1.11] (the situation is similar here except that possibly  $\text{ind}(T - w)$  could be  $-\infty$ ) to conclude that for every  $w_0 \in U$ , there exist  $H$ -valued holomorphic functions  $e_1(w), \dots, e_n(w)$  defined on some neighborhood  $\Delta \subseteq U$  of  $w_0$  such that  $\{e_1(w), \dots, e_n(w)\}$  forms a basis for  $\text{Ker}(T - w)$  for every  $w \in \Delta$ . In view of the correspondence between vector bundles and locally free sheaves,  $\mathcal{F}_T|_U$  is locally free of rank  $n$ .  $\square$

We now state the main result of this section (cf. [26, Section 5]).

**Theorem 3.3.** *Let  $p$  be a non-constant polynomial in the complex variable  $z$  of degree  $n$  and let  $T$  be the bounded linear operator  $M_p^*$  on  $H^2(\mathbb{D})$ . Then there exist clopen (possibly empty) subsets  $\Omega_1, \dots, \Omega_n$  of  $p(\mathbb{D})$  such that*

$$p(\mathbb{D}) \setminus p(\mathbb{T}) = \bigsqcup_{k=1}^n \Omega_k \text{ (disjoint union)}.$$

Moreover, the following statements are valid:

- (i) for every  $k = 1, \dots, n$ ,  $\mathcal{F}_T|_{\Omega_k}$  is locally free of rank  $k$  provided  $\Omega_k$  is non-empty,
- (ii) if, in addition,  $p$  has real coefficients, then

$$H^2(\mathbb{D}) = \sum_{\substack{w \in \Omega_k \\ k=1, \dots, n}} \text{Ker}(T - w).$$

**Proof.** For  $k = 1, \dots, n$ , define  $\Omega_k$  by

$$\Omega_k = \{\alpha \in p(\mathbb{D}) \setminus p(\mathbb{T}) : p(z) = \alpha \text{ has precisely } k \text{ roots in } \mathbb{D} \text{ (counted with multiplicities)}\}. \tag{8}$$

Let  $\alpha \in \Omega_k$  and note that  $\epsilon := \inf_{z \in \mathbb{T}} |p(z) - \alpha| > 0$ . In particular, for  $\beta \in \mathbb{C}$  such that  $|\beta - \alpha| < \epsilon$ ,

$$|(p(z) - \alpha) - (p(z) - \beta)| < \epsilon \leq |p(z) - \alpha|, \quad z \in \mathbb{T}.$$

By Rouché's Theorem [7],  $\beta$  must belong to  $\Omega_k$ , and hence  $\Omega_k$  is an open subset of  $p(\mathbb{D}) \setminus p(\mathbb{T})$ . We next check that each  $\Omega_k$  is relatively closed in  $p(\mathbb{D}) \setminus p(\mathbb{T})$ . By Lemma 3.1 (applied to  $\phi = p - \alpha$ ) and (8),

$$\Omega_k = \{\alpha \in p(\mathbb{D}) \setminus p(\mathbb{T}) : \dim \text{Ker}(T - \bar{\alpha}) = k\}. \quad (9)$$

Now, if  $\{\alpha_n\}_{n=1}^\infty \subseteq \Omega_k$  is such that  $\alpha_n \rightarrow \alpha$  for some  $\alpha \in p(\mathbb{D}) \setminus p(\mathbb{T})$ , then  $T - \alpha_n$  converges to  $T - \alpha$  in the operator norm, and hence by the continuity of the Fredholm index,  $\text{ind}(T - \bar{\alpha}) = k$ . Hence, by (9),  $\alpha \in \Omega_k$ . Also, since the range of  $T - \alpha$  is closed for every  $\alpha \in p(\mathbb{D}) \setminus p(\mathbb{T})$  (see Lemma 3.1), the desired conclusion in (i) is now immediate from Lemma 3.2.

To see (ii), let  $f \in H^2(\mathbb{D})$  be such that  $\langle f, g \rangle = 0$  for every  $g \in \text{Ker}(T - w)$ ,  $w \in \Omega_j$  and  $k = 1, \dots, n$ . Choose  $k \in \{1, \dots, n\}$  such that  $\Omega_k$  is non-empty (this is possible since  $p$  is non-constant). Let  $\alpha \in \Omega_k$  be such that  $p^{-1}\{\alpha\} \cap Z(p') = \emptyset$ , where  $Z(p')$  denotes the zero set of the derivative  $p'$  of  $p$ . We contend that for  $j = 1, \dots, k$ , there exist an open neighborhood  $V$  of  $\alpha$  and a holomorphic function  $g_j : V \rightarrow H^2(\mathbb{D})$  such that  $\{g_j(w) : j = 1, \dots, k\}$  forms a basis  $\text{Ker}(T - w)$  for every  $w \in V$ . Note that  $p$  is a covering map from  $\mathbb{D} \setminus Z(p')$  onto  $p(\mathbb{D} \setminus Z(p'))$ . Hence, one can choose a neighborhood  $V$  centered at  $\alpha$  such that

$$p^{-1}(V) \cap Z(p') = \emptyset, \quad p^{-1}(V) = \sqcup_{j=1}^k W_j$$

such that  $p|_{W_j}$  is a one-one holomorphic map onto  $V$ . Let  $q_j : V \rightarrow W_j$  be the holomorphic inverse of  $p|_{W_j}$ , and define  $g_j(w) = \kappa(\cdot, q_j(w))$ ,  $w \in V$  (see (5)). Assume now that  $p$  has real coefficients. Note that

$$\begin{aligned} Tg_j(w) &= M_p^* \kappa(\cdot, \overline{q_j(w)}) = \overline{p(q_j(w))} \kappa(\cdot, \overline{q_j(w)}) \\ &= p(q_j(w)) \kappa(\cdot, \overline{q_j(w)}) = wg_j(w), \quad w \in V, \quad j = 1, \dots, k. \end{aligned}$$

Thus  $g_j(w) \in \text{Ker}(T - w)$ , and hence by (6),

$$f \circ \overline{q_j(w)} = \langle f, g_j(w) \rangle = 0, \quad w \in V.$$

By the open mapping and the identity theorem (see [7]),  $f$  is identically zero. This completes the verification of part (ii).  $\square$

*Remark 3.4.* We make several remarks in order:

- (1) The part (ii) says that the eigenvectors of  $M_p^*$  corresponding to the points in the complement of the essential spectrum  $\sigma_e(T_p^*)$  of  $T_p^*$  are complete (see [6, Corollary 2], [32, Corollary 1] for variants).
- (2) The argument given for part (ii) actually yields the following stronger assertion: If  $\Omega_k$  is non-empty for some  $k = 1, \dots, n$ , then

$$H^2(\mathbb{D}) = \sum_{w \in \Omega_k} \text{Ker}(T - w).$$

- (3) The conclusions in (i) and (ii) hold true, with obvious modifications, if we replace  $\overline{p}$  the polynomial  $p$  by a bounded holomorphic function  $g$  satisfying  $g(\bar{z}) = g(z)$ ,  $z \in \mathbb{D}$ .

- (4) In the terminology of [19],  $p$  is an  $m$ -analytic cover of the connected component  $\Omega_m$  of  $p(\mathbb{D}) \setminus p(\mathbb{T})$  for every  $m = 1, \dots, n$ . In particular,  $M_p^*$  belongs to the Cowen-Douglas class  $\mathbf{B}_m(\Omega_m)$  provided  $\Omega_m \neq \emptyset$ .
- (5) It may be deduced from Oka's Coherence Theorem and Three Lemma for coherent sheaves (see [24, Corollary 2.1.31]) that  $\mathcal{F}_T|_{p(\mathbb{D}) \setminus p(\mathbb{T})}$  is a coherent sheaf (cf. [4, Proposition 1.4]).

In view of Remark 3.4(4), we certainly know that  $M_p^*$  is in  $\mathbf{B}_m(U)$  for some positive integer  $m$  and some domain  $U$ . The essential point in Theorem 3.3 is that it captures spectral information about  $M_p^*$  beyond  $U$ . This is achieved after examining the sheaf  $\mathcal{F}_{M_p^*}$ .

*Example 3.5.* We discuss here two examples.

- (1) If  $p(z) = az^n + b$  for a non-zero scalar  $a$ , then  $\Omega_j = \emptyset$  if and only if  $j \neq n$ , and hence  $p(\mathbb{D}) \setminus p(\mathbb{T}) = \Omega_n$ . In this case,  $\mathcal{F}_T|_{\Omega_n}$  is a locally free sheaf of rank  $n$ .
- (2) It may happen that  $\Omega_j$  is non-empty for more than one value of positive integer  $j$ . For instance, if  $p(z) = z(z^2 - \frac{z}{2} - 1)$ , then  $-\frac{1}{2} \in \Omega_1$  (the roots of  $p(z) = -1/2$  are  $\pm 1, \frac{1}{2}$ ) and  $0 \in \Omega_2$  (the roots of  $p(z) = 0$  are  $0, \frac{1}{4} \pm \frac{\sqrt{17}}{4}$ ). ◇

#### 4. Relation with the Putinar's sheaf

In this short section, we reveal the relation of the sheaf  $\mathcal{F}_T$  with the cokernel sheaf introduced in [26]. Recall that the *cokernel sheaf*  $\mathcal{G}_T$  associated to a bounded linear operator  $T$  on  $H$  is the sheaf associated with the presheaf

$$\mathcal{G}_T(U) = \mathcal{O}_H(U)/(T - w)\mathcal{O}_H(U), \quad U \subseteq \mathbb{C} \text{ open.}$$

This sheaf first appears in [26] in the context of local spectral theory. It turns out that if  $T$  has the Bishop's property ( $\beta$ ) (see [1, Definition 6.14]), then  $\mathcal{G}_T$  is a sheaf (see [26, Proposition 1.3], [13, P. 9]). To explain the precise relation between the cokernel sheaf and the eigensheaf, we need a definition. Following [9], we say that  $w_0 \in \mathbb{C}$  is a *point of stability* of a bounded linear operator  $T$  on  $H$  if  $T - w_0$  is Fredholm and  $\dim \text{Ker}(T - w)$  is constant in a neighbourhood of  $w_0$ .

**Proposition 4.1.** *For  $T \in \mathcal{L}(H)$ , let  $\mathcal{F}_T$  be the eigensheaf of  $T$ . If  $w_0$  is a point of stability for  $T$ , then there exists a neighbourhood  $N_{w_0}$  of  $w_0$  such that the following hold:*

- (i) *if  $T^t$  denotes the real transpose of  $T$ , then  $\mathcal{G}_{T^t}|_{N_{w_0}}$  is a sheaf,*
- (ii) *the sheaves  $\mathcal{G}_{T^t}|_{N_{w_0}}$  and  $\mathcal{F}_T|_{N_{w_0}}$  are isomorphic.*

*If (ii) holds, then  $w_0 \in \text{Supp}(\mathcal{G}_{T^t})$  if and only if  $w_0 \in \text{Supp}(\mathcal{F}_T)$ .*

**Proof.** By the proof of [9, Proposition 1.11], there exist a neighbourhood  $N_{w_0}$  of  $w_0$  outside the essential spectrum of  $T$  and a holomorphic function  $P : N_w \rightarrow \mathcal{L}(H)$  such that

$$\text{Ker}(T - w) = P(w)H, \quad w \in N_{w_0}. \quad (10)$$

By the closed range theorem, the range of  $T^t - w$  is closed for every  $w \in N_{w_0}$ . For any  $f \in \mathcal{O}_H(N_{w_0})$ , we now obtain the orthogonal decomposition of  $H$  when considered over the real field  $\mathbb{R}$  :

$$f(w) = f_1(w) + (T^t - w)f_2(w), \quad w \in N_{w_0}, \quad (11)$$

where  $f_1(w) \in \text{Ker}(T - w)$  and  $f_2(w) \in \text{Ker}(T^t - w)^\perp$ . By (10),  $w \mapsto f_1(w) \in \mathcal{O}_H(N_{w_0})$ . It follows that  $w \mapsto (T^t - w)f_2(w)$  is holomorphic on  $N_{w_0}$ . Thus, by the Chain rule and the continuity of  $T$ , for every  $w \in N_{w_0}$  and  $h \in H$ ,

$$\begin{aligned} \frac{\partial}{\partial \bar{w}} \langle (T^t - w)f_2(w), h \rangle &= \frac{\partial}{\partial \bar{w}} \langle f_2(w), (T^t)^*h \rangle - w \frac{\partial}{\partial \bar{w}} \langle f_2(w), h \rangle \\ &= \langle (T^t - w) \frac{\partial}{\partial \bar{w}} f_2(w), h \rangle, \end{aligned}$$

and hence by the Cauchy-Riemann equations,  $\frac{\partial}{\partial \bar{w}} f_2(w) \in \text{Ker}(T^t - w)$ . Since  $f_2(w) \in \text{Ker}(T^t - w)^\perp$ , by a routine limit argument,  $\frac{\partial f_2(w)}{\partial \bar{w}} \in \text{Ker}(T^t - w)^\perp$ . This implies that  $\frac{\partial f_2(w)}{\partial \bar{w}} = 0$ , or equivalently

$$f_2 \in \mathcal{O}_H(N_{w_0}). \quad (12)$$

For any non-empty open subset  $U$  of  $N_{w_0}$ , define an  $\mathcal{O}_{\mathbb{C}}(U)$ -module morphism  $\Phi(U) : \mathcal{G}_{T^t}|_U \rightarrow \mathcal{F}_T|_U$  by setting

$$\Phi(U)(f + (T^t - w)\mathcal{O}_H(U)) = f_1(w), \quad f \in \mathcal{O}_H(U).$$

Clearly,  $\Phi(U)$  is surjective. If  $f_1 = 0$ , then by (11) and (12),  $f$  belongs to  $(T^t - w)\mathcal{O}_H(U)$ , and hence  $\Phi(U)$  is injective. It is evident that this morphism is compatible with restrictions. Since isomorphic image of a sheaf is a sheaf, this yields (i) and (ii). The remaining part is now clear.  $\square$

*Remark 4.2.* If  $T$  is a hyponormal operator on a separable Hilbert space, then by Corollary 2.8,  $\mathcal{F}_T$  is trivial, and hence by Proposition 4.1,  $\mathcal{G}_{T^t}|_{N_{w_0}}$  is trivial.

Proposition 4.1 helps us in revealing the structure of the support of the sheaf  $\mathcal{F}_T$  under the assumption of finite cyclicity of the adjoint  $T^*$  of  $T$ .

**Corollary 4.3.** For  $T \in \mathcal{L}(H)$ , let  $\mathcal{F}_T$  be the eigensheaf of  $T$ . Suppose that there exist finitely many vectors  $h_1, \dots, h_k \in H$  such that  $H = \vee \{T^{*n}h_j : n \geq 0, j = 1, \dots, k\}$ . Then

$$\text{Supp}(\mathcal{F}_T) \setminus \sigma_r(T) = \sigma(T) \setminus \sigma_r(T),$$

where  $\sigma_r(T)$  denotes the right spectrum of  $T$ .

**Proof.** Since  $\sigma_{ap}(T^*) = \sigma_r(T)^*$ , applying [5, Lemma 5.3] to  $T^*$ , we conclude that  $\lambda$  is a point of stability of  $T$  for every  $\lambda \notin \sigma_r(T)$ . By Proposition 4.1,

$$\text{Supp}(\mathcal{F}_T) \setminus \sigma_r(T) = \text{Supp}(\mathcal{G}_{T^t}) \setminus \sigma_r(T).$$

Since  $\text{Supp}(\mathcal{G}_{T^t}) = \sigma(T^t) = \sigma(T)$  (see [26, Lemma 2.1]), the desired conclusion is now immediate.  $\square$

## Acknowledgment

We express our deep gratitude to N. Raghavendra for suggesting this topic including Definition 2.1 and Proposition 2.5. This paper would not have seen the present form without a number of helpful and stimulating conversations with him. Further, the first author is thankful to the faculty and the administrative unit of School of Mathematics, HRI, Allahabad and School of Mathematics and Statistics, University of Hyderabad for their warm hospitality during the preparation of the paper. We also convey our sincere thanks to G. Misra for several helpful suggestions concerning the subject of this paper. Finally, the authors would like to thank the anonymous referee for several valuable inputs improving the presentation of the paper.

## References

- [1] AIENA, PIETRO. Fredholm and local spectral theory, with applications to multipliers. *Kluwer Academic Publishers, Dordrecht*, (2004). xiv+444 pp. ISBN: 1-4020-1830-4. MR2070395, Zbl 1077.47001, doi: 10.1007/1-4020-2525-4. 873, 879
- [2] ARENDT, WOLFGANG; BATTY, CHARLES J. K.; HIEBER, MATTHIAS; NEUBRANDER, FRANK. Vector-valued Laplace transforms and Cauchy problems. Second edition. *Monographs in Mathematics*, 96. *Birkhäuser/Springer Basel AG, Basel*, 2011. xii+539 pp. ISBN: 978-3-0348-0086-0. MR2798103, Zbl 1226.34002, doi: 10.1007/978-3-0348-0087-7. 870, 872
- [3] BISWAS, INDRANIL; MISRA, GADADHAR.  $\widetilde{SL(2, R)}$ -homogeneous vector bundles. *Internat. J. Math.* **19** (2008), no. 1, 1–19. MR2380469, Zbl 1158.53043, doi: 10.1142/S0129167X08004534. 869
- [4] BISWAS, SHIBANANDA; MISRA, GADADHAR; PUTINAR, MIHAI. Unitary invariants for Hilbert modules of finite rank. *J. Reine Angew. Math.* **662** (2012), 165–204. MR2876263, Zbl 1243.46063, arXiv:0909.1902, doi: 10.1515/CRELLE.2011.091. 869, 870, 879
- [5] CHAVAN, SAMEER; STOCHEL, JAN. Weakly concave operators. Preprint, 2021. arXiv:2108.10228. 881
- [6] CLARK, DOUGLAS N. On Toeplitz operators with loops. *J. Operator Theory* **4** (1980), no. 1, 37–54. MR0587367, Zbl 0468.47014. 876, 878
- [7] CONWAY, JOHN B. Functions of one complex variable. Second edition. *Graduate Texts in Mathematics*, 11. *Springer-Verlag, New York-Berlin*, 1978. xiii+317 pp. ISBN: 0-387-90328-3. MR0503901, Zbl 0277.30001, doi: 10.1007/978-1-4612-6313-5. 878
- [8] COWEN, MICHAEL J. Fredholm operators with the spanning property. *INDIANA UNIV. MATH. J.* **35** (1986), no. 4, 855–895. MR0865431, Zbl 0655.47018, doi: 10.1512/iumj.1986.35.35044. 869, 875
- [9] COWEN, MICHAEL J.; DOUGLAS, RONALD G. Complex geometry and operator theory. *Acta Math.* **141** (1978), no. 3–4, 187–261. MR0501368, Zbl 0427.47017, doi: 10.1007/BF02545748. 868, 869, 874, 876, 877, 879, 880

- [10] CURTO, RAÚL E.; SALINAS, NORBERTO. Generalized Bergman kernels and the Cowen–Douglas theory. *Amer. J. Math.* **106** (1984), no. 2, 447–488. MR0737780, Zbl 0583.47037, doi: 10.2307/2374310. 874
- [11] DOUGLAS, RONALD G. Banach algebra techniques in operator theory. Pure and Applied Mathematics, 49. *Academic Press, New York-London*, 1972. xvi+216 pp. MR0361893, Zbl 0247.47001. 876
- [12] ESCHMEIER, JÖRG; PUTINAR, MIHAI. Spectral theory and sheaf theory. III. *J. Reine Angew. Math.* **354** (1984), 150–163. MR767577, Zbl 0539.47002. 869, 870
- [13] ESCHMEIER, JÖRG; PUTINAR, MIHAI. Spectral decompositions and analytic sheaves. London Mathematical Society Monographs. New Series, 10. Oxford Science Publications. *The Clarendon Press, Oxford University Press, New York*, 1996. x+362 pp. ISBN: 0-19-853667-4. MR1420618, Zbl 0855.47013. 869, 870, 871, 879
- [14] ESCHMEIER, JÖRG; SCHMITT, JOHANNES. Cowen–Douglas operators and dominating sets. *J. Operator Theory* **72** (2014), no. 1, 277–290. MR3246991, Zbl 1324.47016, doi: 10.7900/jot.2013jan21.1976. 874
- [15] FORSTER, OTTO. Lectures on Riemann surfaces. Graduate Texts in Mathematics, 81. *Springer-Verlag, New York*, 1991. viii+254 pp. ISBN: 0-387-90617-7. MR1185074, Zbl 0475.30002. 874
- [16] GROTHENDIECK, ALEXANDER. Éléments de géométrie algébrique. I. Le langage des schémas, *Inst. Hautes Études Sci. Publ. Math.* **4** (1960), 228pp. MR217083, Zbl 0118.36206, doi: 10.1007/BF02684778. 873
- [17] HWANG, IN SUNG; LEE, WOO YOUNG. Hyponormality of trigonometric Toeplitz operators. *Trans. Amer. Math. Soc.* **354** (2002), no. 6, 2461–2474. MR1888332, Zbl 1011.47023, doi: 10.1090/S0002-9947-02-02970-7. 876
- [18] KRANTZ, STEVEN G.; SALINAS, NORBERTO. Proper holomorphic mappings and the Cowen–Douglas class. *Proc. Amer. Math. Soc.* **117** (1993), no. 1, 99–105. MR1104397, Zbl 0802.47021, doi: 10.2307/2159703. 876
- [19] LIN, QING; SALINAS, NORBERTO. A characterization of coanalytical Toeplitz operators in Cowen–Douglas class. *Analysis and partial differential equations*, 519–527. Lecture Notes in Pure and Appl. Math, 122. *Dekker, New York*, 1990. MR1044806, Zbl 0703.47023. 876, 879
- [20] LIN, QING; SALINAS, NORBERTO. Proper holomorphic maps and analytic Toeplitz  $n$ -tuples. *Indiana Univ. Math. J.* **39** (1990), no. 3, 547–562. MR1078730, Zbl 0731.47027, doi: 10.1512/iumj.1990.39.39030. 876
- [21] MARTIN, MIRCEA; PUTINAR, MIHAI. Lectures on hyponormal operators. Operator Theory: Advances and Applications, 39. *Birkhäuser Verlag, Basel*, 1989. 304 pp. ISBN: 3-7643-2329-9. MR1028066, Zbl 0684.47018, doi: 10.1007/978-3-0348-7466-3. 873
- [22] MARTÍNEZ-AVENDAÑO, RUBÉN A.; ROSENTHAL, PETER. An introduction to operators on the Hardy–Hilbert space. Graduate Texts in Mathematics, 237. *Springer-Verlag, New York*, 2007. xii+220 pp. ISBN: 978-0-387-35418-7; 0-387-35418-2. MR2270722, Zbl 1116.47001, doi: 10.1007/978-0-387-48578-2. 876
- [23] MISRA, GADADHAR. Operators in the Cowen–Douglas class and related topics. *Handbook of analytic operator theory*, 87–137, CRC Press/Chapman Hall Handb. Math. Ser. *CRC Press, Boca Raton, FL*, 2019. MR3966616, Zbl 1447.47027, arXiv:1901.03801. 869
- [24] MORYE, ARCHANA S. On the Serre–Swan theorem, and on vector bundles over real abelian varieties. Thesis (Ph.D.)–Harish-Chandra Research Institute, Allahabad, 2011. 102 pp. [http://www.hri.res.in/~libweb/theses/softcopy/thes\\_archana\\_morye.pdf](http://www.hri.res.in/~libweb/theses/softcopy/thes_archana_morye.pdf). 871, 873, 879
- [25] PAULSEN, VERN I.; RAGHUPATHI, MRINAL. An introduction to the theory of reproducing kernel Hilbert spaces. Cambridge Studies in Advanced Mathematics, 152. *Cambridge University Press, Cambridge*, 2016. x+182 pp. ISBN: 978-1-107-10409-9. MR3526117, Zbl 1364.46004, doi: 10.1017/CBO97811316219232. 873

- [26] PUTINAR, MIHAI. Spectral theory and sheaf theory. I. *Dilation theory, Toeplitz operators, and other topics* (Timișoara/Herculane, 1982), 283–297. Oper. Theory Adv. Appl., 11. Birkhäuser, Basel, 1983. MR0789644, Zbl 0525.47010. 869, 870, 872, 877, 879, 881
- [27] PUTINAR, MIHAI. Spectral theory and sheaf theory. II. *Math. Z.* **192** (1986), no. 3, 473–490. MR845220, Zbl 0608.47011, doi: 10.1007/BF01164022. 869, 870
- [28] PUTINAR, MIHAI. Spectral theory and sheaf theory. IV. *Operator theory: operator algebras and applications, Part 2* (Durham, NH, 1988), 273–293. Proc. Sympos. Pure Math., 51, Part 2. Amer. Math. Soc., Providence, RI, 1990. MR1077448, Zbl 0778.47023, doi: 10.1090/pspum/051.2/1077448. 869, 870
- [29] ŠUBIN, MIKHAIL A. Factorization of matrix functions dependent on a parameter in normed rings, and related questions in the theory of Noetherian operators. *Mat. Sb.(N.S.)* **73 (115)** (1967), 610–629; translation in *Math. USSR Sb.*, **2** (1967), 543–560. MR0217638, Zbl 0182.17202, doi: 10.1070/SM1967v002n04ABEH002354. 877
- [30] VUKOTIĆ, DRAGAN. Analytic Toeplitz operators on the Hardy space  $H^p$ : a survey. *Bull. Belg. Math. Soc. Simon Stevin* **10** (2003), no. 1, 101–113. MR2032329, Zbl 1039.47016. 876
- [31] WELLS, RAYMOND O., JR. Differential analysis on complex manifolds. Second edition. Graduate Texts in Mathematics, 65. Springer-Verlag, New York-Berlin, 1979. x+260 pp. ISBN: 0-387-90419-0. MR0608414 (83f:58001), Zbl 0435.32004. 870, 871
- [32] YAKUBOVICH, DMITRY V. Dual piecewise analytic bundle shift models of linear operators. *J. Funct. Anal.* **136** (1996), no. 2, 294–330. MR1380657, Zbl 0867.47010, doi: 10.1006/jfan.1996.0032. 876, 878
- [33] ZHU, KEHE. Operators in Cowen–Douglas classes. *Illinois J. Math.* **44** (2000), no. 4, 767–783. MR1804320, Zbl 0972.47006, doi: 10.1215/ijm/1255984691. 874

(Sameer Chavan) DEPARTMENT OF MATHEMATICS AND STATISTICS, INDIAN INSTITUTE OF TECHNOLOGY KANPUR, INDIA  
chavan@iitk.ac.in

(Archana Morye) SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HYDERABAD, INDIA  
asmsm@uohyd.ac.in

This paper is available via <http://nyjm.albany.edu/j/2022/28-35.html>.