

On the Galois-Gauss sums of weakly ramified characters

Yu Kuang

ABSTRACT. Bley, Burns and Hahn used relative algebraic K -theory methods to formulate a precise conjectural link between the (second Adams-operator twisted) Galois-Gauss sums of weakly ramified Artin characters and the square root of the inverse different of finite, odd degree, Galois extensions of number fields. We provide concrete new evidence for this conjecture in the setting of extensions of odd prime-power degree by using a refined version of a well-known result of Ullom.

CONTENTS

1. Introduction	417
2. Preliminaries	421
3. The torsion subgroup of $K_0(\mathbb{Z}[\Gamma], \mathbb{Q}[\Gamma])$	425
4. Results for extensions of odd prime-power degree	428
5. The results of Vinatier	435
References	439

1. Introduction

The subject of ‘classical’ Galois module theory was developed by Fröhlich, M. Taylor and others during the 1970’s and 1980’s (see [Fro83]).

At the heart of this theory was the amazingly close interplay between arithmetic properties of the Galois-Gauss sums that are associated to tamely ramified Artin characters and the structure of Hermitian-Galois modules that are attached to rings of algebraic integers in tamely ramified extensions of number fields.

Much subsequent work has been dedicated to attempts to prove a natural extension of this theory in the setting of Artin characters that are not tamely ramified.

Nevertheless, even today, the arithmetic properties of the Galois-Gauss sums of such characters are still, in general, very poorly understood.

Received September 22, 2022.

2010 *Mathematics Subject Classification.* 11R33, 16E20, 11S15.

Key words and phrases. Galois-Gauss sums; Galois module structure; relative algebraic K -theory; weak ramification.

However, for one special class of wildly ramified characters, some important progress has been made.

To discuss this, we fix an *odd degree* Galois extension of number fields L/K , with $G := \text{Gal}(L/K)$, and we write $\mathcal{D}_{L/K}$ for the different of L/K . Then, by Hilbert's classical formula for the valuation of $\mathcal{D}_{L/K}$, there exists a unique G -stable fractional ideal $\mathcal{A}_{L/K}$ of L such that

$$(\mathcal{A}_{L/K})^2 = (\mathcal{D}_{L/K})^{-1}.$$

Further, if L/K is 'weakly ramified' in the sense of [Ere91] (i.e. the second ramification subgroup in G of every place of K is trivial, which, in particular, permits L/K to be wildly ramified), then Erez has shown that $\mathcal{A}_{L/K}$ is a projective G -module and has suggested that the Hermitian-Galois structure associated to this module should provide an appropriate analogue of the ring of integers for such extensions.

This aspect of the theory of $\mathcal{A}_{L/K}$ was developed in a series of articles of Erez and others and culminated in the article [ET92] of Erez and Taylor which showed that for *tamely* ramified extensions L/K of odd degree, the Hermitian-Galois structures that are attached to $\mathcal{A}_{L/K}$ are completely determined by the 'twisted' Galois-Gauss sums that are obtained by combining the classical Galois-Gauss sum with the action on characters of a second Adams operator.

It is, of course, natural to hope that such results could be extended to the setting of extensions that are weakly and wildly (rather than tamely) ramified.

This has, however, proved to be a very difficult, and rather technical, problem and the most impressive results to date are contained in a series of articles of Vinatier (see, for example, [Vin03] and [Vin05]), of Pickett and Vinatier [PV13], of Pickett and Thomas [PT16] in which weakly (wildly) ramified extensions satisfying a variety of restrictive hypotheses are considered.

Next we note that, taking inspiration from a very different direction (relating to leading term conjectures), Bley, Burns and Hahn [BBH20] (and see also the related PhD thesis [Hah16] of Hahn) have more recently introduced techniques of relative algebraic K -theory to this area of research in order to formulate a precise conjectural link between twisted Galois-Gauss sums of weakly ramified Artin characters and the square root of the inverse different.

In this way, they have, in particular, been able to explain how much of the theory developed by Erez, by Erez and Taylor and by Vinatier could be refined in a natural way.

To be more precise we write $\text{Cl}(\mathbb{Z}[G])$ for the reduced projective class group of $\mathbb{Z}[G]$ and $K_0(\mathbb{Z}[G], \mathbb{Q}^c[G])$ for the relative algebraic K_0 -group of the ring inclusion $\mathbb{Z}[G] \subset \mathbb{Q}^c[G]$.

Then in [BBH20], the authors describe a canonical pre-image $\mathfrak{a}_{L/K}$ of the stable-isomorphism class $[\mathcal{A}_{L/K}]$ of the (projective) module $\mathcal{A}_{L/K}$ in $\text{Cl}(\mathbb{Z}[G])$ under the (surjective) connecting homomorphism of relative K -theory $\partial_G : K_0(\mathbb{Z}[G], \mathbb{Q}^c[G]) \rightarrow \text{Cl}(\mathbb{Z}[G])$.

We recall that the kernel of the homomorphism ∂_G is very large: for example, if G is trivial, then $\text{Cl}(\mathbb{Z}[G])$ vanishes, whilst the kernel of ∂_G is isomorphic to the multiplicative group $\mathbb{Q}^{c^\times}/\{\pm 1\}$. For this reason, the study of $\mathfrak{a}_{L/K}$ is, at least in principle, likely to be much more difficult than that of $[\mathcal{A}_{L/K}]$.

Nevertheless, by using techniques of classical Galois module theory, Bley, Burns and Hahn were able to prove (in [BBH20, Th. 5.2]) that $\mathfrak{a}_{L/K}$ belongs to the torsion subgroup $K_0(\mathbb{Z}[G], \mathbb{Q}[G])_{\text{tor}}$ of the subgroup $K_0(\mathbb{Z}[G], \mathbb{Q}[G])$ of $K_0(\mathbb{Z}[G], \mathbb{Q}^c[G])$, that the elements $\mathfrak{a}_{L/K}$ have good functorial properties under change of extension L/K and that $\mathfrak{a}_{L/K}$ simultaneously controls both the Hermitian-Galois structures and metrized structures that arise naturally from the (self-dual) module $\mathcal{A}_{L/K}$.

To study the element $\mathfrak{a}_{L/K}$ further, the authors also define a canonical variant of the classical ‘unramified characteristic’ that has played a key role in the results of Fröhlich, Taylor et al. The elementary definition of this ‘idelic twisted unramified characteristic’ $\mathfrak{c}_{L/K}$ of L/K makes it clear that it also belongs to $K_0(\mathbb{Z}[G], \mathbb{Q}[G])_{\text{tor}}$ and enjoys the same functoriality properties under change of extension as does $\mathfrak{a}_{L/K}$.

When taken together with extensive numerical computations, these facts motivated Bley, Burns and Hahn to formulate the following conjecture.

Conjecture 1.1 ([BBH20, Conjecture 10.7]). *If L/K is any weakly ramified Galois extension of number fields of odd degree, then one has $\mathfrak{a}_{L/K} = \mathfrak{c}_{L/K}$.*

This concrete conjecture provides the motivation for much of the work that we undertake in this article, and is itself interesting for several reasons.

Firstly, we recall (from the introduction to [BBH20]) that the approach underlying the formulation of Conjecture 1.1 provides the first link between the theory of $\mathcal{A}_{L/K}$ and the very general framework of Tamagawa number conjectures that originates with Bloch and Kato in [BK90] and was subsequently refined to the relevant ‘equivariant’ setting by Burns and Flach in [BF01] (this link, though important, will however play no role in the results that we present in this article).

More concretely, we note that, since the element $\mathfrak{c}_{L/K}$ is both very elementary in nature and easy to compute explicitly, the above conjecture suggests that the link between the twisted Galois-Gauss sums of weakly ramified Artin characters of G and the invariants of $\mathcal{A}_{L/K}$ that are incorporated into the definition of $\mathfrak{a}_{L/K}$ should (or, at least, could!) be much finer than has previously been observed in the work of Erez, of Erez and Taylor or of Vinatier.

We recall that the strongest theoretical evidence in support of Conjecture 1.1 that is currently available is the following result (taken from [BBH20, Cor. 8.4]).

Proposition 1.2 (Bley, Burns and Hahn). *Let L/K be a weakly ramified finite Galois extension of number fields of odd degree. Then the following claims are valid.*

- (i) *The element $\mathfrak{a}_{L/K} - \mathfrak{c}_{L/K}$ belongs to $K_0(\mathbb{Z}[G], \mathbb{Q}[G])_{\text{tor}}$.*

- (ii) *Conjecture 1.1 is valid provided that every place v of K that is wildly ramified in L has the following three properties:*
- (a) *the decomposition subgroup in G of any place of L above v is abelian;*
 - (b) *the inertia subgroup in G of any place of L above v is cyclic;*
 - (c) *the completion of K at v is absolutely unramified.*

The proof of claim (i) above is reduced (by straightforward functoriality arguments, and a general result of Burns [Bur04] concerning $K_0(\mathbb{Z}[G], \mathbb{Q}[G])_{\text{tor}}$) to consideration of tamely ramified extensions.

The proof of claim (ii), however, depends heavily on detailed technical computations in certain families of wildly ramified extensions that are made by Pickett and Vinatier in [PV13]. For this reason, obtaining a full verification of Conjecture 1.1 for classes of extensions that go beyond those in Proposition 1.2(ii) seems to be a very difficult problem.

The main purpose of this paper is to provide further supporting evidence for Conjecture 1.1 by proving the following result.

Theorem 1.3. *Fix an odd prime p and a weakly ramified Galois extension of number fields L/K of p -power degree, and set $G := \text{Gal}(L/K)$. Suppose p is unramified in K and write $n(L/K)$ for the maximal order of a decomposition subgroup in G of a wildly ramified place. Then, in $K_0(\mathbb{Z}[G], \mathbb{Q}[G])$, one has*

$$\frac{n(L/K)}{p} \cdot \alpha_{L/K} = \frac{n(L/K)}{p} \cdot \epsilon_{L/K} = 0.$$

This result is of interest since there are many weakly ramified Galois extensions L/K of odd prime-power degree in which the decomposition subgroup of each wildly ramified place is non-abelian (we recall, for example, the explicit extensions constructed by Vinatier in [Vin02]).

Theorem 1.3 will be proved by combining (both parts of) Proposition 1.2 with a refined version, proved in Theorem 3.1, of a well-known result of Ullom [Ull74] concerning the structure of the kernel subgroup of $\text{Cl}(\mathbb{Z}[G])$ in the case that G has prime-power order. The latter result itself complements recent work of Bley and Wilson [BW09] concerning the explicit computation of relative algebraic K -groups and so is, we hope, of some independent interest.

We next recall that, if L/K is tamely ramified (and of odd degree), then in [Ere91, Th. 3] Erez used the techniques of classical Galois module theory to prove that $[\mathcal{A}_{L/K}]$ vanishes and hence, by Jacobinski's Cancellation Theorem (which is valid since G has odd order, see [CR87, Th. (51.3) and Th. (51.24)]), that $\mathcal{A}_{L/K}$ is a free $\mathbb{Z}[G]$ -module.

Motivated by this result, and several other factors (including extensive numerical computations), Vinatier was then led to make the following conjecture (cf. [Vin03, Conj.] and [CV16, §1.2]).

Conjecture 1.4 (Vinatier). *If L/K is a weakly ramified Galois extension of number fields of odd degree, then $\mathcal{A}_{L/K}$ is a free $\mathbb{Z}[G]$ -module (or equivalently, the class $[\mathcal{A}_{L/K}]$ vanishes).*

As evidence for this conjecture, Vinatier has proved explicit upper bounds on the order of the class $[\mathcal{A}_{L/K}]$ in the case that $K = \mathbb{Q}$ and G is of prime-power order (for details see the articles [Vin03] and [Vin05]).

If one projects the result of Theorem 1.3 into $\text{Cl}(\mathbb{Z}[G])$, then one obtains a bound on the order of $[\mathcal{A}_{L/K}]$ that is weaker than the corresponding bound established by Vinatier. The reason for this is that a key step in the approach of Vinatier cannot, as far as we can see, be repeated in the setting of relative algebraic K -groups (see Remark 5.2).

Nevertheless, we are able to show that the approach developed by Vinatier in [Vin03] can be used to obtain concrete new information about the relative element $\alpha_{L/K}$ of Bley, Burns and Hahn.

This result is proved as Theorem 5.1 and, whilst it does not (as far as we can currently see) provide more evidence in support of Conjecture 1.1 than is given by Theorem 1.3, it does certainly provide further evidence for the belief that the link between $\mathcal{A}_{L/K}$ and Galois-Gauss sums should be closer than has so far been observed.

Acknowledgements I would like to thank David Burns for suggesting the problem and many inspiring discussions. I am very grateful to Stéphane Vinatier for helpful, and encouraging, correspondence. Finally, I am very grateful to the anonymous referee for their careful review and valuable feedback, particularly, for having pointed out an error in an earlier version of the proof of Theorem 3.1 and for motivating me to obtain a sharper bound in the class group that is consistent with Vinatier's result.

2. Preliminaries

2.1. Notations. For a unital ring A , we write A^\times for the multiplicative group of invertible elements of A and $\zeta(A)$ for the centre of A .

Fix a finite group Γ , and a Dedekind domain R of characteristic zero, with field of fractions F , and E is an extension field of F . We let \mathcal{O}_E denote the integral closure of R in E . For any $R[\Gamma]$ -module M , we write $M_E := E \otimes_R M$, and $M_v := R_v \otimes_R M$, with R_v the completion of R at v , where v is a (non-zero) prime ideal of R .

Assume E/F is a finite Galois extension of fields. We let $\text{Gal}(E/F)$ denote the Galois group of E/F , and we write the action of $\text{Gal}(E/F)$ on E by $x \mapsto g(x)$ for $x \in E$ and $g \in \text{Gal}(E/F)$. We fix a separable closure F^c of F , and write Ω_F for the absolute Galois group $\text{Gal}(F^c/F)$ of F . For convenience, we take \mathbb{Q}^c to be the algebraic closure of \mathbb{Q} in \mathbb{C} .

For a finite group Γ , we write $\widehat{\Gamma}$ for the set of \mathbb{Q}^c -valued irreducible characters of Γ and R_Γ for the additive group generated by $\widehat{\Gamma}$. In particular, for each χ in $\widehat{\Gamma}$, we obtain a primitive idempotent of the centre $\zeta(\mathbb{Q}^c[\Gamma])$ of $\mathbb{Q}^c[\Gamma]$ by setting $e_\chi = \frac{\chi(1)}{|\Gamma|} \sum_{g \in \Gamma} \chi(g^{-1})g$ and so each element of $\zeta(\mathbb{Q}^c[\Gamma])$ (respectively $\zeta(\mathbb{Q}^c[\Gamma])^\times$) can

be written uniquely in the form

$$x = \sum_{\chi \in \hat{\Gamma}} e_{\chi} \cdot x_{\chi}, \quad \text{with } x_{\chi} \in \mathbb{Q}^c, \text{ respectively } x_{\chi} \in \mathbb{Q}^{c^{\times}}, \text{ for all } \chi. \quad (1)$$

2.2. Algebraic K-theory. In this section, we recall the definitions of relevant algebraic K -groups and the long exact sequences. Detailed descriptions of these definitions and results are given by Swan in [Swa68, Chap. 13 and 15] and also by Curtis and Reiner in [CR87, Chap. 5 and 6].

We will let $\mathcal{P}(E[\Gamma])$ denote the category of finitely generated projective $E[\Gamma]$ -modules and $\text{Aut}_{E[\Gamma]}(M)$ for the group of $E[\Gamma]$ -module automorphisms of M .

2.2.1. Whitehead groups. We write $K_1(E[\Gamma])$ for the Whitehead group of the ring $E[\Gamma]$. In particular, we recall that each element of $K_1(E[\Gamma])$ can be represented by a pair $[P, f]$ where $P \in \mathcal{P}(E[\Gamma])$ and $f \in \text{Aut}_{E[\Gamma]}(P)$. (We refer the reader to [CR87, §38B] for a detailed discussion of this group in terms of generators and relations.)

Suppose E is either a number field or a p -adic field (for some prime p). Then, for some finite index set I , there is a Wedderburn decomposition $E[\Gamma] = \prod_{i \in I} M_{n_i}(A_i)$ and each matrix ring $M_{n_i}(A_i)$ is a simple central algebra over the centre $\zeta(A_i)$ of A_i . In this case, one can compute in $K_1(E[\Gamma])$ by using the ‘reduced norm’ map discussed by Curtis and Reiner in [CR87, 45A]

$$\begin{aligned} \text{Nrd}_{E[\Gamma]} : K_1(E[\Gamma]) &= \bigoplus_{i \in I} K_1(M_{n_i}(A_i)) \\ &\cong \bigoplus_{i \in I} K_1(A_i) \xrightarrow{(\text{Nrd}_{A_i})_i} \bigoplus_i \zeta(A_i)^{\times} = \zeta(E[\Gamma])^{\times}, \quad (2) \end{aligned}$$

in which the isomorphism follows from the Morita equivalence (applied to each ring $M_{n_i}(A_i)$).

2.2.2. The relative K-group. We write $K_0(R[\Gamma], E[\Gamma])$ for the relative algebraic K -group associated to $R \subset E$ and recall that each element of this group is represented by a triple $[P, \phi, Q]$ where $P, Q \in \mathcal{P}(R[\Gamma])$ and $\phi \in \text{Is}_{E[\Gamma]}(P_E, Q_E)$, the set of $E[\Gamma]$ -module isomorphisms from P_E to Q_E . (For an explicit description of this group in terms of the generators and relations, see [Swa68, pp. 215].)

Then there is a canonical decomposition

$$K_0(R[\Gamma], F[\Gamma]) \xrightarrow{\sim} \bigoplus_v K_0(R_v[\Gamma], F_v[\Gamma]), \quad (3)$$

where v runs over all non-Archimedean places of F . This isomorphism is induced by the diagonal localisation homomorphism $(\pi_{\Gamma, v})_v$ (see, for example, discussion below [CR87, (49.12)]), where for each non-zero prime ideal v of R , we write

$$\pi_{\Gamma, v} : K_0(R[\Gamma], F[\Gamma]) \rightarrow K_0(R_v[\Gamma], F_v[\Gamma]) \quad (4)$$

for the homomorphism that sends the class of $[P, \phi, Q]$ to the class of $[P_v, F_v \otimes_F \phi, Q_v]$.

2.3. Equivariant Galois-Gauss sums. Suppose E/F is a finite Galois extension of ℓ -adic fields and set $\Gamma := \text{Gal}(E/F)$. In terms of (1), we define the following ‘equivariant’ elements in $\zeta(\mathbb{Q}^c[\Gamma])^\times$ as in [BBH20, §4A].

Definition 2.1.

- (i) The ‘equivariant Galois-Gauss sum’ of E/F is the element

$$\tau_{E/F} \in \zeta(\mathbb{Q}^c[\Gamma])^\times$$

defined by setting

$$\tau_{E/F} := \sum_{\chi \in \widehat{\Gamma}} e_\chi \cdot \tau(F, \chi).$$

where $\tau(F, \chi)$ is the (local) Galois Gauss sum discussed in [Fro83, Chap. III, §2, Th. 18 and Rem. 1].

- (ii) For each $\phi \in \widehat{\Gamma}$, the ‘unramified characteristic’ of ϕ is defined by setting (as in [Fro83, Chap. IV, §1, (1.1)])

$$y(F, \phi) = \begin{cases} 1, & \text{if } \phi \text{ is ramified;} \\ (-1)^{\phi(1)} \phi(\sigma), & \text{if } \phi \text{ is unramified,} \end{cases}$$

where σ is the Frobenius element in Γ/Γ_0 lifted to Γ and Γ_0 is the inertia subgroup of Γ .

- (iii) The ‘equivariant unramified characteristic’ of E/F is the element of $\zeta(\mathbb{Q}[\Gamma])^\times$ obtained by setting

$$y_{E/F} := \sum_{\chi \in \widehat{\Gamma}} e_\chi \cdot y(F, \chi).$$

- (iv) The ‘modified equivariant Galois-Gauss sum’ of E/F is the element

$$\tau'_{E/F} := \tau_{E/F} \cdot y_{E/F}^{-1} \in \zeta(\mathbb{Q}^c[\Gamma])^\times.$$

- (v) We set $\tau_F := \tau(\mathbb{Q}_\ell, \text{ind}_F^{\mathbb{Q}_\ell} \mathbf{1}_F)$ and define the ‘induced discriminant’ of E/F to be the element τ_F^Γ of $\zeta(\mathbb{Q}^c[\Gamma])^\times$ with

$$(\tau_F^\Gamma)_\chi = \tau_F^{\chi(1)}$$

for all $\chi \in \widehat{\Gamma}$.

Last, for each natural number k and each $\chi \in R_\Gamma$, we write ψ_k for the k -th Adams operator on R_Γ (cf. [CR81, Prop. (12.8)]). Then for each pair of integers m and n , and each natural number k , we define an endomorphism $m + n \cdot \psi_{k,*}$ of $\zeta(\mathbb{Q}^c[\Gamma])^\times$ (in terms of (1)) such that for each element x of $\zeta(\mathbb{Q}^c[\Gamma])^\times$, the element $(m + n \cdot \psi_{k,*})(x)$ is uniquely specified by $((m + n \cdot \psi_{k,*})(x))_\chi := (x_\chi)^m \cdot (x_{\psi_k(\chi)})^n$ for each irreducible character χ in $\widehat{\Gamma}$.

3. The torsion subgroup of $K_0(\mathbb{Z}[\Gamma], \mathbb{Q}[\Gamma])$

Fix a finite group Γ and let R denote either \mathbb{Z} or \mathbb{Z}_ℓ for a prime ℓ , and \mathbb{Q} denote the corresponding fields \mathbb{Q} or \mathbb{Q}_ℓ . In this section we study the torsion subgroup

$$DT(R[\Gamma]) := K_0(R[\Gamma], \mathbb{Q}[\Gamma])_{\text{tor}}$$

of $K_0(R[\Gamma], \mathbb{Q}[\Gamma])$.

We recall that these groups have already been studied in the article [BW09] of Bley and Wilson that contains both general results and numerical algorithms.

To start, we first note that the isomorphism (3) restricts to give a canonical direct sum decomposition

$$DT(\mathbb{Z}[\Gamma]) \cong \bigoplus_{\ell} DT(\mathbb{Z}_\ell[\Gamma]) \tag{9}$$

over all primes ℓ .

Next, for each prime ℓ , we fix a maximal order \mathcal{M}_ℓ in $\mathbb{Q}_\ell[\Gamma]$ that contains $\mathbb{Z}_\ell[\Gamma]$ and we note that \mathcal{M}_ℓ is hereditary (by [CR81, Th. (26.12)]). Then the results of [BW09, Th. 2.2(iv) and 2.4(i)] imply that

$$DT(\mathbb{Z}_\ell[\Gamma]) \cong \frac{\text{Nrd}_{\mathbb{Q}_\ell[\Gamma]}(\mathcal{M}_\ell^\times)}{\text{Nrd}_{\mathbb{Q}_\ell[\Gamma]}(\mathbb{Z}_\ell[\Gamma]^\times)}. \tag{10}$$

In the sequel, unless there is danger of confusion, we shall abbreviate $\text{Nrd}_{\mathbb{Q}_\ell[\Gamma]}$ to Nrd . This isomorphism then combines with the direct sum decomposition (9) to give an isomorphism

$$DT(\mathbb{Z}[\Gamma]) \cong \prod_{\ell} \frac{\text{Nrd}(\mathcal{M}_\ell^\times)}{\text{Nrd}(\mathbb{Z}_\ell[\Gamma]^\times)} = \prod_{\ell \in \Sigma(\Gamma)} \frac{\text{Nrd}(\mathcal{M}_\ell^\times)}{\text{Nrd}(\mathbb{Z}_\ell[\Gamma]^\times)}. \tag{11}$$

where $\Sigma(\Gamma)$ denotes the finite set of primes that divide the order of Γ (or equivalently, at which $\mathcal{M}_\ell \neq \mathbb{Z}_\ell[\Gamma]$). This isomorphism implies that $DT(\mathbb{Z}[\Gamma])$ is finite (see [BW09, Cor. 2.5]).

In the next result we prove finer results for $DT(\mathbb{Z}[\Gamma])$ in the case that Γ has prime power order. Claim (i) of this result is a version for relative K -groups of a well-known bound on the exponent of the kernel group $D(\mathbb{Z}[\Gamma])$ that is due to Ullom [Ull74] (see also [CR87, Th. (50.19)]).

Theorem 3.1. *Let p be an odd prime and Γ be a non-trivial group of order p^n for a natural number n . Then, the following claims are valid.*

- (i) *The exponent of $DT(\mathbb{Z}[\Gamma])$ is divisible by $p - 1$ and divides $(1 - 1/p)|\Gamma|$.*
- (ii) *Write $\Xi(\Gamma)$ for the set of normal subgroups of Γ that are given by the kernels of the irreducible characters of Γ . Then, the exponent of the kernel of the diagonal projection map*

$$DT(\mathbb{Z}[\Gamma]) \rightarrow \bigoplus_{\Delta \in \Xi(\Gamma)} DT(\mathbb{Z}[\Gamma/\Delta])$$

divides $|\Gamma|/p$.

(iii) *Claims (i) and (ii) remain valid if one replaces \mathbb{Z} by \mathbb{Z}_p .*

Proof. Fix a normal subgroup Δ of Γ and set $e_\Delta := |\Delta|^{-1} \sum_{g \in \Delta} g$. For each prime ℓ we write $\mathcal{M}_{\ell, \Gamma/\Delta} := e_\Delta \mathcal{M}_\ell$. Then, since the order \mathcal{M}_ℓ is maximal, the central idempotent e_Δ belongs to \mathcal{M}_ℓ , and so \mathcal{M}_ℓ decomposes as a direct sum $(1 - e_\Delta) \mathcal{M}_\ell \oplus \mathcal{M}_{\ell, \Gamma/\Delta}$. This decomposition implies that $\mathcal{M}_{\ell, \Gamma/\Delta}$ is a maximal order in $\mathbb{Q}_\ell[\Gamma/\Delta]$, and also that the natural map $\mathcal{M}_\ell^\times \rightarrow \mathcal{M}_{\ell, \Gamma/\Delta}^\times$ is surjective. By the isomorphism (11), we can therefore deduce that the natural projection map

$$\mathrm{DT}(\mathbb{Z}[\Gamma]) \rightarrow \mathrm{DT}(\mathbb{Z}[\Gamma/\Delta]) \quad (12)$$

is surjective.

We now assume that $|\Gamma| = p^n$. In this case, the set $\Sigma(\Gamma)$ is equal to $\{p\}$ and so (11) implies that the natural map $\mathrm{DT}(\mathbb{Z}[\Gamma]) \rightarrow \mathrm{DT}(\mathbb{Z}_p[\Gamma])$ is bijective. In particular, if we can prove claims (i) and (ii), then claim (iii) is clearly also true.

To do this, we first choose Δ to have index p (this can be done since, in this case, any subgroup of Γ of index p is normal), then $\mathrm{DT}(\mathbb{Z}[\Gamma/\Delta])$ is cyclic of order $p - 1$ (see, for example, [BW09, Cor. 8.2]), and so the surjection (12) implies that the exponent of $\mathrm{DT}(\mathbb{Z}[\Gamma])$ is divisible by $p - 1$, as stated in claim (i).

Next, to prove that the exponent of $\mathrm{DT}(\mathbb{Z}[\Gamma]) = \mathrm{DT}(\mathbb{Z}_p[\Gamma])$ divides $(p - 1)p^{n-1}$, we recall that, by Schilling's Theorem (cf. [CR87, Th. (74.17)]), for a suitable natural number t , the Wedderburn decomposition of $\mathbb{Q}_p[\Gamma]$ has the form

$$\mathbb{Q}_p[\Gamma] \cong \mathbb{Q}_p \times \prod_{i=1}^{i=t} M_{n_i}(F_i), \quad (13)$$

where each n_i is a natural number and each F_i is a p -adic cyclotomic field of the form $\mathbb{Q}_p(\omega_i)$ for a suitable non-trivial p -power root of unity ω_i . In the sequel we set $F_0 := \mathbb{Q}_p$, and for each i , we write \mathcal{O}_i for the valuation ring of F_i and \mathfrak{p}_i for the maximal ideal of \mathcal{O}_i .

In this way, by [BW09, Th. 2.2(iv)] we have that

$$\mathrm{Nrd}(\mathcal{M}_p^\times) = \prod_{i=0}^{i=t} \mathcal{O}_i^\times,$$

where the right hand side is the unit group of the integral closure of \mathbb{Z}_p in the centre $\prod_{i=0}^{i=t} F_i$ of $\mathbb{Q}_p[\Gamma]$. Therefore, each element x of $\mathrm{Nrd}(\mathcal{M}_p^\times)$ can be written in the form

$$x = (u_0, u_1, \dots, u_t), \quad (14)$$

with $u_i \in \mathcal{O}_i^\times$ for each $0 \leq i \leq t$. Moreover, since each extension F_i/\mathbb{Q}_p is totally ramified, the natural map $\mathbb{Z}_p/(p) \rightarrow \mathcal{O}_i/\mathfrak{p}_i$ is bijective, and so one has $u_i^{p-1} \equiv 1 \pmod{\mathfrak{p}_i}$.

It follows that x^{p-1} is equal to a tuple (v_0, v_1, \dots, v_t) in which each element v_i belongs to \mathcal{O}_i^\times and is such that $v_i \equiv 1$ modulo \mathfrak{p}_i . In view of the isomorphism (10), in order to prove claim (i), it is therefore enough to show that the p^{n-1} -st

power of any such tuple belongs to $\text{Nrd}(\mathbb{Z}_p[\Gamma]^\times)$. This is exactly what is proved by the argument of Ullom that is discussed on pp. 257-259 of [CR87] (in which our tuple corresponds to the element α), and so this concludes the proof of claim (i).

To prove claim (ii), we note that the components in the Wedderburn decomposition (13) are in bijective correspondence with a set of representatives for the orbits of the natural action of $\text{Gal}(\mathbb{Q}_p^c/\mathbb{Q}_p)$ on the set $\widehat{\Gamma}_p$ of irreducible \mathbb{Q}_p^c -valued characters of Γ . More precisely, we fix a subset $\{\psi_j : 0 \leq j \leq t\}$ of $\widehat{\Gamma}_p$ such that ψ_0 is the trivial character and set $F_0 := \mathbb{Q}_p$, $\mathcal{O}_0 = \mathbb{Z}_p$ and $n_0 = 1$ for $j = 0$. For each character ψ_j , we let $T_{j,*}$ denote the \mathbb{Q}_p -linear map $\mathbb{Q}_p[\Gamma] \rightarrow M_{n_j}(F_j)$ that is induced by a choice of a representation $T_j : \Gamma \rightarrow \text{GL}_{n_j}(F_j)$ of ψ_j . In this way, for each j , the projection map

$$\mathbb{Q}_p[\Gamma] \rightarrow M_{n_j}(F_j)$$

that is induced by (13) coincides with $T_{j,*}$. We fix an embedding $\mathbb{Q}^c \rightarrow \mathbb{Q}_p^c$ and we shall identify $\widehat{\Gamma}$ with $\widehat{\Gamma}_p$ via the choice of the embedding. Then, since Γ is nilpotent of odd order, the results of Roquette in [Roq58] (or see the more general results of Cliff, Ritter and Weiss in [CRW92]) imply that one can choose a T_j (of ψ_j) such that it takes values in $\text{GL}_{n_j}(\mathcal{O}_j)$ and this is what we shall do.

Set $\Delta_j := \ker(\psi_j) \in \Xi(\Gamma)$ and $\Gamma_j := \Gamma/\Delta_j$, we note that $T_{j,*}$ factors through the projection map $\pi_j : \mathbb{Q}_p[\Gamma] \rightarrow \mathbb{Q}_p[\Gamma_j]$. We also note that $T_{0,*}$ is the natural augmentation map $\varepsilon_\Gamma : \mathbb{Q}_p[\Gamma] \rightarrow \mathbb{Q}_p$.

Now, we fix an element y in \mathcal{M}_p^\times . Then, for each j , the element u_j that occurs in the decomposition (14) of $x = \text{Nrd}(y)$ belongs to \mathcal{O}_j^\times and can be computed as $\det(T_{j,*}(y))$.

To prove claim (ii), we claim that it is enough to show that if the image \bar{x} of x in $\text{DT}(\mathbb{Z}_p[\Gamma])$ belongs to the kernel of the projection map $\pi_{i,*} : \text{DT}(\mathbb{Z}_p[\Gamma]) \rightarrow \text{DT}(\mathbb{Z}_p[\Gamma_i])$ for any given i with $1 \leq i \leq t$, then one has

$$\det(T_{i,*}(y)) \equiv (T_{0,*}(y))^{\psi_i(1)} \pmod{\mathfrak{p}_i}. \tag{15}$$

To see that the above result implies claim (ii), we assume that these congruences are valid for each $1 \leq i \leq t$. Set $v_i := u_0^{-\psi_i(1)} u_i$, we note that such element satisfies $v_i \equiv 1$ modulo \mathfrak{p}_i for all i . In particular, since u_0 belongs to $\mathbb{Z}_p^\times \subseteq \mathbb{Z}_p[\Gamma]^\times$, these congruences and the argument of claim (i) combine to imply that the element

$$\begin{aligned} x^{p^{n-1}} &= (u_0, u_1, \dots, u_t)^{p^{n-1}} \\ &= (u_0, u_0^{\psi_1(1)}, \dots, u_0^{\psi_t(1)})^{p^{n-1}} \times (1, v_1, \dots, v_t)^{p^{n-1}} \\ &= \text{Nrd}(u_0)^{p^{n-1}} \times (1, v_1, \dots, v_t)^{p^{n-1}} \end{aligned}$$

belongs to $\mathbb{Z}_p[\Gamma]^\times$, and hence that $\bar{x}^{p^{n-1}} = 0$, as required.

To prove the stated congruence (15), we assume that \bar{x} belongs to the kernel of $\pi_{i,*}$. Then, the isomorphism (10) implies that there exists an element $z_i =$

$\sum_{\gamma \in \Gamma_i} c_{i,\gamma} \gamma \in \mathbb{Z}_p[\Gamma_i]^\times$ such that

$$\pi_{i,*}(\text{Nrd}_{\mathbb{Q}_p[\Gamma]}(y)) = \text{Nrd}_{\mathbb{Q}_p[\Gamma_i]}(z_i) \quad (16)$$

and hence,

$$\det(T_{i,*}(y)) = \det(T_{i,*}(z_i)) = \det\left(\sum_{\gamma \in \Gamma_i} c_{i,\gamma} \cdot T_i(\gamma)\right). \quad (17)$$

Now, we note that (by the choice of T_i as above) each matrix $T_i(\gamma)$ belongs to $\text{GL}_{n_i}(\mathcal{O}_i)$. Upon setting $m_i := |\Gamma_i|$, the matrix $T_i(\gamma)^{m_i} = T_i(\gamma^{m_i})$ is equal to the $n_i \times n_i$ identity matrix I_{n_i} . Hence, since m_i belongs to \mathfrak{p}_i , the binomial theorem implies that there are congruences modulo $M_{n_i}(\mathfrak{p}_i)$ of the form

$$\begin{aligned} \left(\sum_{\gamma \in \Gamma_i} c_{i,\gamma} \cdot T_i(\gamma)\right)^{m_i} &\equiv \sum_{\gamma \in \Gamma_i} c_{i,\gamma}^{m_i} \cdot I_{n_i} \\ &\equiv \left(\sum_{\gamma \in \Gamma_i} c_{i,\gamma}^{m_i}\right) \cdot I_{n_i} \\ &\equiv \left(\sum_{\gamma \in \Gamma_i} c_{i,\gamma}\right)^{m_i} \cdot I_{n_i} \\ &\equiv (T_{0,*}(y))^{m_i} \cdot I_{n_i} \quad (\text{mod } M_{n_i}(\mathfrak{p}_i)), \end{aligned}$$

where the last congruence is true since $T_{0,*}$ is the natural augmentation map and so, in terms of the expression (16), we have that $T_{0,*}(y)$ is equal to $\sum_{\gamma \in \Gamma_i} c_{i,\gamma}$.

Since $n_i = \psi_i(1)$, these congruences and the equality (17) combine to imply that

$$\begin{aligned} \det(T_{i,*}(y))^{m_i} &= \det\left(\sum_{\gamma \in \Gamma_i} c_{i,\gamma} \cdot T_i(\gamma)\right)^{m_i} \\ &\equiv \det\left((T_{0,*}(y))^{m_i} \cdot I_{n_i}\right) \quad (\text{mod } \mathfrak{p}_i) \\ &\equiv \left((T_{0,*}(y))^{\psi_i(1)}\right)^{m_i} \quad (\text{mod } \mathfrak{p}_i). \end{aligned}$$

Finally, since $\det(T_{i,*}(x))$ and $(T_{0,*}(x))^{\psi_i(1)}$ both belong to \mathcal{O}_i^\times (and the order of $(\mathcal{O}_i/\mathfrak{p}_i)^\times$ is prime to m_i), the above congruence then implies the required congruence (15). This completes the proof of the claim. \square

Remark 3.2.

- (i) The result of Theorem 3.1(ii) is of interest only if $\Xi(\Gamma)$ does not contain the trivial subgroup of Γ . This condition is equivalent to requiring that Γ has no faithful irreducible characters and such groups are completely classified by Gaschutz in [Gas54].
- (ii) If Γ is abelian, then the set $\Xi(\Gamma)$ is equal to the set of subgroups Δ of Γ with the property that the quotient group Γ/Δ is cyclic.

4. Results for extensions of odd prime-power degree

In this section, we try to provide some new evidence for Conjecture 1.1 in the case of extensions of odd prime-power degree.

4.1. The canonical relative elements of Bley, Burns and Hahn. Firstly, we recall the key definition of the canonical relative elements of Bley, Burns and Hahn from [BBH20].

4.1.1. The local element. In this section, we fix a finite odd degree Galois extension E/F of local fields of residue characteristic ℓ and set $\Gamma := \text{Gal}(E/F)$. We follow Bley, Burns and Hahn [BBH20, §7A] in defining the local relative elements.

We write $\Sigma(E)$ for the set of \mathbb{Q}_ℓ -linear field embeddings $E \rightarrow \mathbb{Q}_\ell^c$. We then define a (free) $\mathbb{Z}_\ell[\Gamma]$ -module by setting

$$H_E := \prod_{\Sigma(E)} \mathbb{Z}_\ell,$$

upon which Γ acts via its usual pre-composition action on $\Sigma(E)$, and we consider the isomorphism of $\mathbb{Q}_\ell^c[\Gamma]$ -modules

$$\kappa_E : \mathbb{Q}_\ell^c \otimes_{\mathbb{Q}_\ell} E \rightarrow \prod_{\Sigma(E)} \mathbb{Q}_\ell^c = \mathbb{Q}_\ell^c \otimes_{\mathbb{Z}_\ell} H_E$$

that sends $x \otimes l$, for every $x \in \mathbb{Q}_\ell^c$ and l in E to $(\sigma(l)x)_{\sigma \in \Sigma(E)}$.

Suppose E/F is weakly ramified. Then, by the result [Ere91, Th. 1] of Erez, $\mathcal{A}_{E/F}$ is a full projective $\mathbb{Z}_\ell[\Gamma]$ -sublattice of E and so, following [BBH20, §7A], we obtain a well-defined element of $K_0(\mathbb{Z}_\ell[\Gamma], \mathbb{Q}_\ell^c[\Gamma])$ by setting

$$\Delta(\mathcal{A}_{E/F}) := [\mathcal{A}_{E/F}, \kappa_E, H_E].$$

For the next definition, we use the following notation: for each prime ℓ and embedding of fields $j_\ell^c : \mathbb{Q}^c \rightarrow \mathbb{Q}_\ell^c$, we also write j_ℓ^c for the induced homomorphism of rings $\mathbb{Q}^c[G] \rightarrow \mathbb{Q}_\ell^c[G]$ and consider the homomorphism of abelian groups

$$\begin{aligned} j_{\ell,*}^c : K_0(\mathbb{Z}[\Gamma], \mathbb{Q}^c[\Gamma]) &\rightarrow K_0(\mathbb{Z}_\ell[\Gamma], \mathbb{Q}_\ell^c[\Gamma]), \\ [P, \phi, Q] &\mapsto [P_\ell, \mathbb{Q}_\ell^c \otimes_{\mathbb{Q}^c, j_\ell^c} \phi, Q_\ell]. \end{aligned} \tag{18}$$

Definition 4.1. We set

$$T_{E/F}^{(2)} := \tau_F^\Gamma \cdot (\psi_{2,*} - 1)(\tau'_{E/F}) \in \zeta(\mathbb{Q}^c[\Gamma])^\times,$$

where the elements τ_F^Γ and $\tau'_{E/F}$ are as in §2.3.

We then define an element of $K_0(\mathbb{Z}_\ell[\Gamma], \mathbb{Q}_\ell^c[\Gamma])$ by setting

$$\mathfrak{a}_{E/F} := \Delta(\mathcal{A}_{E/F}) - \delta_{\mathbb{Z}_\ell, \mathbb{Q}_\ell^c, \Gamma}(j_\ell^c(T_{E/F}^{(2)})) - U_{E/F},$$

where $U_{E/F}$ is the canonical ‘unramified’ element of $K_0(\mathbb{Z}_\ell[\Gamma], \mathbb{Q}_\ell^c[\Gamma])$ that is defined by Breuning in [Bre04b].

In particular, we note (by [BBH20, Prop. 7.1]) that $\mathfrak{a}_{E/F}$ is independent of the choice of embedding j_ℓ^c and belongs to $K_0(\mathbb{Z}_\ell[\Gamma], \mathbb{Q}_\ell^c[\Gamma])$.

4.1.2. The global element. Let L/K be a weakly ramified finite Galois extension of number fields of odd degree with Galois group $G := \text{Gal}(L/K)$. In [BBH20, §2A3 and §5], Bley, Burns and Hahn define a global canonical relative element of $K_0(\mathbb{Z}[G], \mathbb{Q}^c[G])$

$$\mathfrak{a}_{L/K} := \Delta(\mathcal{A}_{L/K}) - \delta_{\mathbb{Z}, \mathbb{Q}^c, G}(T_{L/K}^{(2)}),$$

where $\Delta(\mathcal{A}_{L/K})$ and $T_{L/K}^{(2)}$ are the global analogues of the elements defined above.

Fix a place w of L above a place v of K , we write G_w for the decomposition subgroup of w in G , and for each $\chi \in R_G$, we let χ_w denote the restriction of χ to G_w (therefore regard χ_w as characters of $\text{Gal}(L_w/K_v)$ via identifying $G_w = \text{Gal}(L_w/K_v)$). In this article, we will often use the decomposition property of the relative element (taken from Theorem 7.6 of loc.cit.) that

$$\mathfrak{a}_{L/K} = \sum_{\ell} \sum_{v|\ell} i_{G_w, \mathbb{Q}_{\ell}}^{G, *}(\mathfrak{a}_{L_w/K_v}), \quad (19)$$

where the first sum runs over all rational primes ℓ and the second over all places v of K of residue characteristic ℓ .

Remark 4.2. Since (by [BBH20, Prop. 5.5]) $\mathfrak{a}_{L/K}$ belongs to $K_0(\mathbb{Z}[G], \mathbb{Q}[G])$, the sum on the right hand side of the formula in (19) can only have finitely many non-zero ℓ -primary components. In fact, Bley, Burns and Hahn [BBH20, Th. 8.1] have shown that \mathfrak{a}_{L_w/K_v} vanishes if the extension L_w/K_v is tamely ramified (which is true for all v that do not divide the order of G).

Next we recall a variant of the classical ‘unramified characteristic’ that is introduced in loc. cit.

Definition 4.3.

- (i) Let E/F be a finite odd Galois extension of ℓ -adic fields and set $\Gamma := \text{Gal}(E/F)$. Then, the ‘twisted unramified characteristic’ of E/F is the element of $K_0(\mathbb{Z}_{\ell}[\Gamma], \mathbb{Q}_{\ell}[\Gamma])$ that is obtained by setting

$$\mathfrak{c}_{E/F} := \delta_{\mathbb{Z}_{\ell}, \mathbb{Q}_{\ell}, \Gamma}((1 - \psi_{2, *})(y_{E/F})),$$

where $y_{E/F}$ is the equivariant unramified characteristic defined in §2.3. In particular, we recall from [BBH20, Rem. 7.5] that $\mathfrak{c}_{E/F} = 0$ if E/F is tamely ramified.

- (ii) Let L/K be a finite odd Galois extension of number fields and set $G := \text{Gal}(L/K)$. Then, the ‘idelic twisted unramified characteristic’ of L/K is defined to be the element

$$\mathfrak{c}_{L/K} = \sum_{\ell} \sum_{v|\ell} i_{G_w, \mathbb{Q}_{\ell}}^{G, *}(\mathfrak{c}_{L_w/K_v}) \in K_0(\mathbb{Z}[G], \mathbb{Q}[G]).$$

Remark 4.4. Conjecture 1.1 is compatible with the change of extension functors. To see this, we recall from [BBH20, Th. 6.1 and Rem. 8.9] that, for any subgroup J of G , the following equalities are valid.

- (i) $\rho_J^{G,*}(\mathfrak{a}_{L/K}) = \mathfrak{a}_{L/L'}$ and $\rho_J^{G,*}(\mathfrak{c}_{L/K}) = \mathfrak{c}_{L/L'}$, where $\rho_J^{G,*}$ denotes the restriction map $K_0(\mathbb{Z}[G], \mathbb{Q}[G]) \rightarrow K_0(\mathbb{Z}[J], \mathbb{Q}[J])$.
- (ii) If J is normal to G , with $Q := G/J$, then $q_Q^{G,*}(\mathfrak{a}_{L/K}) = \mathfrak{a}_{L'/K}$ and $q_Q^{G,*}(\mathfrak{c}_{L/K}) = \mathfrak{c}_{L'/K}$, where $q_Q^{G,*}$ denotes the coinflation map $K_0(\mathbb{Z}[G], \mathbb{Q}[G]) \rightarrow K_0(\mathbb{Z}[Q], \mathbb{Q}[Q])$.

4.2. Proof of Theorem 1.3. The motivation for this section is the fact that there are many weakly ramified extensions L/K of p -power degree in which the decomposition subgroup of every p -adic place is non-abelian, so Proposition 1.2(ii) does not apply directly. (For example, in [Vin02, Th. 1.4], Vinatier has constructed an infinite family of weakly ramified Galois extensions of $K = \mathbb{Q}$ of degree 27 in which the decomposition subgroup of each 3-adic place of L is non-abelian.)

However, by combining Proposition 1.2 with the result of Theorem 3.1, we are able to obtain an upper bound on the order of the element $\mathfrak{a}_{L/K}$ and can thereby provide evidence in support of Conjecture 1.1 for a more general class of extensions.

Taking account of the decomposition of $\mathfrak{a}_{L/K}$ given in (19), the result of Bley, Burns and Hahn recalled in Remark 4.2 and the definition of $\mathfrak{c}_{L/K}$ as an explicit sum of local terms (see Definition 4.3(ii)), it is enough for us to show that the stated equalities hold for each (p -adic) place v of K that ramifies wildly in L .

Proposition 4.5. *Fix an odd prime p . Suppose that L/K is a weakly ramified Galois extension of number fields of p -power degree and that p is unramified in K . If v is any p -adic place of K that is wildly ramified in L , we write $G_w := \text{Gal}(L_w/K_v)$ for some place w of L above v and $n_w := |G_w|$. Then, in $K_0(\mathbb{Z}_p[G_w], \mathbb{Q}_p[G_w])$, one has*

$$\frac{n_w}{p} \cdot \mathfrak{a}_{L_w/K_v} = \frac{n_w}{p} \cdot \mathfrak{c}_{L_w/K_v} = 0. \tag{20}$$

In the sequel, we fix an embedding $j_p^c : \mathbb{Q}^c \rightarrow \mathbb{Q}_p^c$. For a finite group Γ , we shall use j_p^c to identify $\widehat{\Gamma}$ with $\widehat{\Gamma}_p$, the set of \mathbb{Q}_p^c -valued irreducible characters of Γ (via the isomorphism $\chi \mapsto \chi^j$ where $\chi^j(g) := j_p^c(\chi(g))$ for all $\chi \in \widehat{\Gamma}$ and $g \in \Gamma$). We also write j_p^c for the induced embedding $\zeta(\mathbb{Q}^c[\Gamma])^\times \rightarrow \zeta(\mathbb{Q}_p^c[\Gamma])^\times$. Let \mathcal{O}_p^t denote the valuation ring of the maximal tamely ramified extension of \mathbb{Q}_p , we consider the associated homomorphism of relative K -groups,

$$J_{p,*}^t : K_0(\mathbb{Z}_p[\Gamma], \mathbb{Q}_p^c[\Gamma]) \rightarrow K_0(\mathcal{O}_p^t[\Gamma], \mathbb{Q}_p^c[\Gamma]), \tag{21}$$

$$[P, \phi, Q] \mapsto [\mathcal{O}_p^t \otimes_{\mathbb{Z}_p} P, \phi, \mathcal{O}_p^t \otimes_{\mathbb{Z}_p} Q].$$

To prove the equalities in (20), we need to show some preliminary results.

For the next three results, we fix a weakly ramified Galois extension of p -adic fields E/F of arbitrary odd degree, set $\Gamma := \text{Gal}(E/F)$ and write Δ for the inertia subgroup of Γ . We also write Γ_1 (resp. Γ_2) for the 1-st (resp. 2-nd) ramification subgroup (in the lower numbering) of Γ . The following result is a consequence of the hypothesis that E/F is weakly ramified (i.e. Γ_2 is trivial).

Lemma 4.6.

- (i) *The group Γ_1 is abelian and of exponent dividing p .*
- (ii) *The inertia subgroup Δ is abelian if and only if it is either a p -group (and hence, equal to Γ_1) or cyclic and of order prime-to- p .*

Proof. At the outset, we note that the group Γ_2 is a normal subgroup of Γ_1 , and the quotient group Γ_1/Γ_2 is abelian and of exponent dividing p (by [Ser79, Chap. IV, §2, Cor. 3 to Prop. 7]). Then, claim (i) follows directly from the hypothesis that E/F is weakly ramified.

Turning to claim (ii). If Δ has order prime-to- p , then it is cyclic (since, in this case, E/F is tamely ramified). If Δ is a p -group, then it is equal to Γ_1 and hence abelian (by claim (i)).

Now, to show the converse, we assume Δ is abelian and set $e_0 := |\Delta/\Gamma_1|$. It is enough for us to consider the case that $e_0 > 1$. In this case, upon applying the result of [Ser79, Chap. IV, §2, Cor. 2 to Prop. 9] to the abelian extension E/E^Δ , one can deduce that $\Gamma_1 = \Gamma_2$ is trivial. It follows that E/F is tamely ramified, and hence that Δ is cyclic. This completes the proof of claim (ii). \square

The second of the required equalities (20) follows directly from the next result.

Lemma 4.7. *If E/F is as above, then in $K_0(\mathbb{Z}_p[\Gamma], \mathbb{Q}_p[\Gamma])$, one has $|\Gamma/\Delta| \cdot \mathbf{c}_{E/F} = 0$.*

Proof. Fix an element σ of Γ that projects to give the Frobenius automorphism in Γ/Δ .

By [BBH20, Rem. 7.5], the element $\mathbf{c}_{E/F}$ is equal to the image under $\delta_{\mathbb{Z}_p, \mathbb{Q}_p, \Gamma}$ of the element

$$(1 - \psi_{2,*})(y_{E/F}) = (1 - e_\Delta) + \sigma^{-1}e_\Delta \in \zeta(\mathbb{Q}_p[\Gamma])^\times. \quad (22)$$

Then, the claimed result is true since,

$$((1 - e_\Delta) + \sigma^{-1}e_\Delta)^{|\Gamma/\Delta|} = (1 - e_\Delta) + (\sigma^{-1}e_\Delta)^{|\Gamma/\Delta|} = (1 - e_\Delta) + e_\Delta = 1.$$

\square

In the next result, we write B for the maximal unramified extension of F in E (so that $\Delta = \text{Gal}(E/B)$). We also recall from [Fro83, Chap. I, §4 and Chap. III, §3, (3.1)] that the *resolvent* element and the (local) *norm resolvent* is defined by setting, for each element a in E generating a normal basis of E/F , and each character χ of representation $T_\chi : \Gamma \rightarrow \text{GL}_n(\mathbb{Q}_p^c)$,

$$(a|\chi) := \det\left(\sum_{g \in \Gamma} g(a)T_\chi(g^{-1})\right), \quad \mathcal{N}_{F/\mathbb{Q}_p}(a|\chi) = \prod_{\omega} (a|\chi^{\omega^{-1}})^{\omega},$$

where for the second product ω runs through a transversal of Ω_F in $\Omega_{\mathbb{Q}_p}$.

Proposition 4.8. *In $K_0(\mathbb{Z}_p[\Gamma], \mathbb{Q}_p[\Gamma])$, one has $|\Gamma/\Delta| \cdot \mathbf{a}_{E/F} = i_{\Delta, \mathbb{Q}_p}^{\Gamma,*}(\mathbf{a}_{E/B})$.*

Proof. By Taylor’s Fixed Point Theorem (cf. [Mar84, Chap. 8, §1]), one knows that the restriction to $K_0(\mathbb{Z}_p[\Gamma], \mathbb{Q}_p[\Gamma])$ of the homomorphism $j_{p,*}^t$ (defined in (21)) is injective, so it is enough to prove that the images in $K_0(\mathcal{O}_p^t[\Gamma], \mathbb{Q}_p^c[\Gamma])$ of the respective elements $|\Gamma/\Delta| \cdot \mathbf{a}_{E/F}$ and $i_{\Delta, \mathbb{Q}_p}^{\Gamma,*}(\mathbf{a}_{E/B})$ coincide.

For $L \in \{B, F\}$, with $\Gamma_L := \text{Gal}(E/L)$, we set $\delta_{\Gamma_L, p} := \delta_{\mathbb{Z}_p, \mathbb{Q}_p^c, \Gamma_L}$ and

$$\mathbf{a}'_{E/L} := \Delta(\mathcal{A}_{E/L}) - \delta_{\Gamma_L, p}(j_p^c(T_{E/L}^{(2)})) \in K_0(\mathbb{Z}_p[\Gamma_L], \mathbb{Q}_p^c[\Gamma_L]).$$

Then, the result [Bre04a, Prop. 4.4] of Breuning that $U_{E/L} \in \ker(j_{p,*}^t)$ implies that the images in $K_0(\mathcal{O}_p^t[\Gamma_L], \mathbb{Q}_p^c[\Gamma_L])$ of the elements $\mathbf{a}_{E/L}$ and $\mathbf{a}'_{E/L}$ coincide. It is therefore enough for us to prove that the images in $K_0(\mathcal{O}_p^t[\Gamma], \mathbb{Q}_p^c[\Gamma])$ of the elements $|\Gamma/\Delta| \cdot \mathbf{a}'_{E/F}$ and $i_{\Delta, \mathbb{Q}_p}^{\Gamma,*}(\mathbf{a}'_{E/B})$ coincide.

To do this, we fix a \mathbb{Z}_p -basis $\{a_\sigma\}_{\sigma \in \Sigma(L)}$ of \mathcal{O}_L and set

$$\delta_L := \det(\tau(a_\sigma))_{\tau, \sigma \in \Sigma(L)} \in \mathbb{Q}_p^c \tag{23}$$

and $u_L := \delta_L/j_p^c(\tau_L)$. We also fix an element a_L of E such that $\mathcal{A}_{E/L} = \mathcal{O}_L[\Gamma_L] \cdot a_L$. Then, by the explicit formula [Bre04a, Lem. 4.16] of Breuning, one has

$$\begin{aligned} \mathbf{a}'_{E/L} &= \delta_{\Gamma_L, p} \left((j_p^c(T_{E/L}^{(2)}))^{-1} \sum_{\chi \in \widehat{\Gamma}_L} (\delta_L^{\chi(1)} \cdot \mathcal{N}_{L/\mathbb{Q}_p}(a_L|\chi)) e_\chi \right) \\ &= \delta_{\Gamma_L, p} \left(\sum_{\chi \in \widehat{\Gamma}_L} \left((\delta_L/j_p^c(\tau_L))^{\chi(1)} \cdot \frac{\mathcal{N}_{L/\mathbb{Q}_p}(a_L|\chi)}{j_p^c(\tau(L, \psi_2(\chi) - \chi)y(L, \chi - \psi_2(\chi)))} \right) e_\chi \right) \\ &= \delta_{\Gamma_L, p}(x_L) + \delta_{\Gamma_L, p} \left(\sum_{\chi \in \widehat{\Gamma}_L} \frac{\mathcal{N}_{L/\mathbb{Q}_p}(a_L|\chi)}{j_p^c(\tau(L, \psi_2(\chi) - \chi) \cdot y(L, \chi - \psi_2(\chi)))} e_\chi \right), \end{aligned}$$

with $x_L := \sum_{\chi \in \widehat{\Gamma}_L} u_L^{\chi(1)} e_\chi = \text{Nrd}_{\mathbb{Q}_p^c[\Gamma_L]}(u_L)$.

In the sequel, we omit the occurrence of j_p^c in each of our notations. Since u_L is a unit of \mathcal{O}_p^t (by [Bre04a, Lem. 4.29]), the image in $K_0(\mathcal{O}_p^t[\Gamma_L], \mathbb{Q}_p^c[\Gamma_L])$ of $\delta_{\Gamma_L, p}(x_L)$ vanishes.

In addition, one has $y(B, \chi - \psi_2(\chi)) = 1$ for every $\chi \in \widehat{\Delta}$ since the extension E/B is totally ramified (and so there is no non-trivial unramified character). One can also compute (directly from Definition 2.1(ii)) that $y(F, \chi - \psi_2(\chi))^{|\Gamma/\Delta|} = 1$ for every $\chi \in \widehat{\Gamma}$.

In view of (8), one can write

$$\begin{aligned}
 i_{\Delta, \mathbb{Q}_p}^{\Gamma, *}\circ\delta_{\Delta, p}\left(\sum_{\phi\in\widehat{\Delta}}\frac{\mathcal{N}_{B/\mathbb{Q}_p}(a_B|\phi)}{j_p^c(\tau(B, \psi_2(\phi) - \phi))}e_\phi\right) &= \delta_{\Gamma, p}\circ i_{\Delta}^{\Gamma}\left(\sum_{\phi\in\widehat{\Delta}}\frac{\mathcal{N}_{B/\mathbb{Q}_p}(a_B|\phi)}{j_p^c(\tau(B, \psi_2(\phi) - \phi))}e_\phi\right) \\
 &= \delta_{\Gamma, p}\left(\sum_{\chi\in\widehat{\Gamma}}\frac{\mathcal{N}_{B/\mathbb{Q}_p}(a_B|\text{res}_\Delta^\Gamma\chi)}{\tau(B, \text{res}_\Delta^\Gamma(\psi_2(\chi) - \chi))}e_\chi\right),
 \end{aligned}
 \tag{24}$$

where the second equality follows from the definition of the induction map (7) and the fact that $\tau(B, \psi_2(\text{res}_\Delta^\Gamma\chi) - \text{res}_\Delta^\Gamma\chi) = \tau(B, \text{res}_\Delta^\Gamma(\psi_2(\chi) - \chi))$ (cf. [Ere91, Prop.-Def. 3.5]).

In order to show that the images in $K_0(\mathcal{O}_p^t[\Gamma], \mathbb{Q}_p^c[\Gamma])$ of the elements $|\Gamma/\Delta| \cdot \mathfrak{a}'_{E/F}$ and $i_{\Delta, \mathbb{Q}_p}^{\Gamma, *}\left(\mathfrak{a}'_{E/B}\right)$ coincide, it is therefore enough for us to show that the elements

$$\sum_{\chi\in\widehat{\Gamma}}\frac{(\mathcal{N}_{F/\mathbb{Q}_p}(a_F|\chi))^{[B:F]}}{\tau(F, \psi_2(\chi) - \chi)^{[B:F]}}e_\chi \quad \text{and} \quad \sum_{\chi\in\widehat{\Gamma}}\frac{\mathcal{N}_{B/\mathbb{Q}_p}(a_B|\text{res}_\Delta^\Gamma\chi)}{\tau(B, \text{res}_\Delta^\Gamma(\psi_2(\chi) - \chi))}e_\chi \tag{25}$$

of $\zeta(\mathbb{Q}_p^c[\Gamma])^\times$ differ by an element of $\text{Nrd}_{\mathbb{Q}_p^c[\Gamma]}(\mathbb{Z}_p[\Gamma]^\times)$.

To prove this, we note that, by the argument of [Fro83, Th. 25 and the following Remark] (which is valid even for extensions that are wildly ramified, as observed by Erez in [Ere91, §6]), for each χ in $\widehat{\Gamma}$, one has

$$\tau(F, \psi_2(\chi) - \chi)^{[B:F]} = \tau(B, \text{res}_\Delta^\Gamma(\psi_2(\chi) - \chi)),$$

since $\psi_2(\chi) - \chi$ has degree zero, and for a suitable choice of generating element a_B , one also has

$$(\mathcal{N}_{F/\mathbb{Q}_p}(a_F|\chi))^{[B:F]} = \mathcal{N}_{B/\mathbb{Q}_p}(a_B|\text{res}_\Delta^\Gamma\chi) \cdot \text{Det}_\chi(\lambda),$$

where the element λ of $\mathbb{Z}_p[\Gamma]^\times$ is independent of χ . This completes the proof of the claimed result. \square

We are now ready to prove Proposition 4.5.

For v and w as in (20), we now write I_w for the inertia subgroup of G_w and B_w for the fixed field of I_w in L_w .

Then, in view of the last result (for $E/F = L_w/K_v$) and the obvious equality $|G_w|/p = |I_w|/p \times |G_w/I_w|$, to prove the first equality in (20), it is therefore enough for us to show that the element \mathfrak{a}_{L_w/B_w} of $K_0(\mathbb{Z}_p[I_w], \mathbb{Q}_p[I_w])$ is annihilated by $|I_w|/p$.

To do this, we write L_0 for the fixed field of I_w in L and w_0 for the place of L_0 obtained by restricting w . Then, since p is unramified in K , the extension $L_{0, w_0}/\mathbb{Q}_p$ is unramified and so L_0 cannot contain a non-trivial p -th root of unity. By the result [Neu79, Cor. 2, p. 156] of Neukirch, we can therefore fix a finite Galois extension L' of L_0 with both of the following properties:

- (i) L' has a unique place w' above w_0 and the completion $L'_{w'}/L_{0,w_0}$ is isomorphic to L_w/B_w ;
- (ii) if v' is any place of L_0 which divides $|I_w|$, and $v' \neq w_0$, then v' is totally split in L'/L_0 .

These conditions imply that the extension L'/L_0 is weakly ramified and that the group $G' := \text{Gal}(L'/L_0)$ identifies with $\text{Gal}(L'_{w'}/L_{0,w_0}) \cong \text{Gal}(L_w/B_w) = I_w$. In particular, since w_0 is the only place of L_0 that ramifies wildly in L' , in this case the decomposition result in (19) combines with the vanishing of \mathfrak{a}_{L'/L_0} for tamely ramified extensions of ℓ -adic fields (see Remark 4.2 with $L/K = L'/L_0$) to imply that

$$\mathfrak{a}_{L'/L_0} = \mathfrak{a}_{L'_{w'}/L_{0,w_0}} = \mathfrak{a}_{L_w/B_w}.$$

It is therefore sufficient for us to show that \mathfrak{a}_{L'/L_0} is annihilated by $|G'|/p$. To do this we note that Proposition 1.2(i) implies that \mathfrak{a}_{L'/L_0} belongs to $\text{DT}(\mathbb{Z}[G'])$.

In addition, Lemma 4.6(ii) implies that the group $G' (\cong I_w)$ is abelian and so the set $\Xi(G')$ in Theorem 3.1(ii) is equal to the set of normal subgroups H of G' with the property that G'/H is cyclic (see Remark 3.2(ii)).

Now, we fix H in $\Xi(G')$ and set $\tilde{L}' := (L')^H$ and $\tilde{G} := G'/H \cong \text{Gal}(\tilde{L}'/L_0)$. Then, Remark 4.4(ii) implies that $q_{\tilde{G}}^{G',*}(\mathfrak{a}_{L'/L_0}) = \mathfrak{a}_{\tilde{L}'/L_0}$. In addition, since \tilde{G} is cyclic of p -power order and L_{0,w_0} is an unramified extension of \mathbb{Q}_p , Proposition 1.2(ii) implies that $\mathfrak{a}_{\tilde{L}'/L_0} = \mathfrak{c}_{\tilde{L}'/L_0}$. Finally, we note that, since w_0 is totally ramified in \tilde{L}' , the expression (22) implies that $\mathfrak{c}_{\tilde{L}'/L_0}$ vanishes.

This argument shows that \mathfrak{a}_{L'/L_0} belongs to the kernel of the diagonal map that occurs in Theorem 3.1(ii) (with $\Gamma = G'$) and so the latter result implies that \mathfrak{a}_{L'/L_0} is annihilated by $|G'|/p$, as required.

This finishes the proof of Proposition 4.5 and hence, Theorem 1.3.

Remark 4.9. Alternatively, one can complete the argument of Proposition 4.5 by proving an analogous result to [BBH20, Th. 6.1] of the functoriality properties of the local element defined in §4.1.1. In this way, for any $H \in \Xi(I_w)$ with $L'_w := (L_w)^H$, one has $q_{I_w/H}^{I_w}(\mathfrak{a}_{L_w/B_w}) = \mathfrak{a}_{L'_w/B_w}$. Moreover, by [BBH20, Th. 8.1] one has $\mathfrak{a}_{L'_w/B_w} = \mathfrak{c}_{L'_w/B_w} = 0$. Hence the result that $\mathfrak{a}_{L'_w/B_w}$ is annihilated by $|I_w|/p$ now follows from Theorem 3.1(ii) and (iii).

5. The results of Vinatier

If L/K is any extension as in Theorem 1.3, the result of Theorem 1.3 can be combined with the result [BBH20, Th. 5.2(iv)] of Bley, Burns and Hahn to deduce that

$$\frac{n(L/K)}{p} \cdot [\mathcal{A}_{L/K}] = 0$$

in $\text{Cl}(\mathbb{Z}[G])$.

In the special case that $K = \mathbb{Q}$, Vinatier [Vin03] has given a much better upper bound on the order of $[\mathcal{A}_{L/\mathbb{Q}}]$, with additional improvements in the case $p = 3$ obtained in [Vin05]. The essential part in proving this bound is that, by

working in $\text{Cl}(\mathbb{Z}[G])$, Vinatier is able to prove a ‘global’ analogue of the induction formula in (25) in which the exponent $|\Gamma/\Delta|$ is replaced by p . (We note that the second expression of (25) is an induction in the sense of (24).)

In the next result, we describe a consequence of Vinatier’s approach for the relative element $\Delta(\mathcal{A}_{L/\mathbb{Q}})$.

We note that while (as far as we can see) this result does not directly provide evidence in support of Conjecture 1.1, it does provide further evidence for the general belief that the elements $\Delta(\mathcal{A}_{L/K})$ should be closely controlled by analytic invariants.

Theorem 5.1. *Fix an odd prime p and a wildly and weakly ramified Galois extension L of \mathbb{Q} of p -power degree and set $G := \text{Gal}(L/\mathbb{Q})$. We fix a p -adic place w of L and write $e(w)$ for its absolute ramification degree and H for its decomposition subgroup in G and identify the latter subgroup with $\text{Gal}(L_w/\mathbb{Q}_p)$. Let i_w^* denote the induction map $K_0(\mathbb{Z}[H], \mathbb{Q}^c[H]) \rightarrow K_0(\mathbb{Z}[G], \mathbb{Q}^c[G])$ and δ_w for the composite homomorphism $\partial_{\mathbb{Z}, \mathbb{Q}^c, H}^1 \circ (\text{Nrd}_{\mathbb{Q}^c[H]})^{-1}$. Then the element*

$$\mathfrak{a}_{L/\mathbb{Q}} + i_w^*(\delta_w((\psi_{2,*} - 1)(\tau'_{L_w/\mathbb{Q}_p})))$$

of $K_0(\mathbb{Z}[G], \mathbb{Q}^c[G])$ is annihilated by $(p - 1)e(w)$.

Proof. We set $\delta := \delta_{\mathbb{Z}, \mathbb{Q}^c, G}$, $\tilde{\tau}_w := (\psi_{2,*} - 1)(\tau'_{L_w/\mathbb{Q}_p})$ and abbreviate the map $\tilde{i}_H^G : \zeta(\mathbb{Q}^c[H])^\times \rightarrow \zeta(\mathbb{Q}^c[G])^\times$ to \tilde{i}_w . Then, since $i_w^* \circ \delta_w = \delta \circ \tilde{i}_w$ (see (8)), the stated element in the claim is equal to

$$\xi := \mathfrak{a}_{L/\mathbb{Q}} + \delta(\tilde{i}_w(\tilde{\tau}_w)).$$

First, we note that the horizontal isomorphism in [Fro83, Chap. II, §1, Lem. 1.6] implies that, in terms of the decomposition (1), for every finite prime ℓ , $a \in \text{GL}_n(\mathbb{Q}_\ell^c[G])$ and χ in \widehat{G} , one has that $\text{Nrd}_{\mathbb{Q}_\ell^c[G]}(a)_\chi = \text{Det}_\chi(a)$, where Det denotes the generalized determinant discussed in [Fro83, Chap. I, §2]. In this way $\text{Nrd}_{\mathbb{Q}_\ell^c[G]}(a)$ can be considered as $\text{Det}(a) : \chi \mapsto \text{Det}_\chi(a)$, an element of $\text{Hom}(R_G, \mathbb{Q}_\ell^{c \times})$. Write \mathcal{M} for the maximal order in $\mathbb{Q}[G]$ that contains $\mathbb{Z}[G]$, we then deduce from [Fro83, Prop. 2.2, pp. 23] that,

$$\prod_{\ell} \text{Nrd}_{\mathbb{Q}_\ell[G]}(\mathcal{M}_\ell^\times) = \text{Hom}(R_G, U_f(\mathbb{Q}^c))^{\Omega_{\mathbb{Q}}},$$

where the product runs over all finite primes ℓ and $U_f(\mathbb{Q}^c)$ denotes the group of unit ideles (i.e. the ideles u whose ℓ -components are units for all finite primes ℓ).

Then, the fact that the values of the unramified characteristic are roots of unity in \mathbb{Q}^c (cf. [Fro83, Th. 29(i)]) and the definition of the induction map (7) combine to imply that $\tilde{i}_w((\psi_{2,*} - 1)(y_{L_w/\mathbb{Q}_p}^{-1}))$ belongs to $\prod_{\ell} \text{Nrd}_{\mathbb{Q}_\ell[G]}(\mathcal{M}_\ell^\times)$. In addition, the result [Vin03, Prop. 2.5] of Vinatier implies that the p -th power of the element $\tilde{i}_w((\psi_{2,*} - 1)(\tau_{L_w/\mathbb{Q}_p}))$ also belongs to $\prod_{\ell} \text{Nrd}_{\mathbb{Q}_\ell[G]}(\mathcal{M}_\ell^\times)$. In view

of the isomorphisms (9) and (10), we can therefore deduce that the element

$$p \cdot \delta(\tilde{i}_w(\tilde{\tau}_w)) = \delta(\tilde{i}_w((\psi_{2,*} - 1)(\tau_{L_w/\mathbb{Q}_p}))^p) + \delta(\tilde{i}_w((\psi_{2,*} - 1)(y_{L_w/\mathbb{Q}_p}^{-1}))^p)$$

belongs to $\text{DT}(\mathbb{Z}[G]) = \text{DT}(\mathbb{Z}_p[G])$ (since $\mathcal{M}_\ell = \mathbb{Z}_\ell[G]$ for all $\ell \neq p$, see (10) and (11)).

Since $\mathfrak{a}_{L/\mathbb{Q}} \in \text{DT}(\mathbb{Z}_p[G])$ (by Proposition 1.2(i)), it follows that the element $p \cdot \xi$ also belongs to the subgroup $\text{DT}(\mathbb{Z}_p[G])$ of $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$. For each prime ℓ , we identify $K_0(\mathbb{Z}_\ell[G], \mathbb{Q}_\ell[G])$ with a subgroup of $K_0(\mathbb{Z}[G], \mathbb{Q}[G])$ by means of the decomposition (4) and recall the homomorphism $j_{p,*}^c$ from (18). In this way the image $j_{p,*}^c(p \cdot \xi)$ is equal to $p \cdot \xi$ in $K_0(\mathbb{Z}_p[G], \mathbb{Q}_p[G])$.

Set $\delta_{w,p} := \delta_{\mathbb{Z}_p, \mathbb{Q}_p^c, H}^1$ and let $i_{w,p}^*$ denote the induction homomorphism

$$K_0(\mathbb{Z}_p[H], \mathbb{Q}_p^c[H]) \rightarrow K_0(\mathbb{Z}_p[G], \mathbb{Q}_p^c[G]).$$

We note that the decomposition result in (19) and Remark 4.2 combine to imply that $\mathfrak{a}_{L/\mathbb{Q}} = i_{w,p}^*(\mathfrak{a}_{L_w/\mathbb{Q}_p})$. Then,

$$\begin{aligned} p \cdot \xi &= p \cdot (j_{p,*}^c(\mathfrak{a}_{L/\mathbb{Q}} + \delta(\tilde{i}_w(\tilde{\tau}_w)))) \\ &= p \cdot i_{w,p}^*(\mathfrak{a}_{L_w/\mathbb{Q}_p} + \delta_{w,p}(j_p^c(\tilde{\tau}_w))) \\ &= p \cdot i_{w,p}^*(\Delta(\mathcal{A}_{L_w/\mathbb{Q}_p}) - \delta_{w,p}(j_p^c(T_{L_w/\mathbb{Q}_p}^{(2)})) - U_{L_w/\mathbb{Q}_p} + \delta_{w,p}(j_p^c(\tilde{\tau}_w))) \\ &= p \cdot i_{w,p}^*(\Delta(\mathcal{A}_{L_w/\mathbb{Q}_p}) - U_{L_w/\mathbb{Q}_p}). \end{aligned} \quad (26)$$

Here, the second equality follows from (8) that $\delta \circ \tilde{i}_w = i_w^* \circ \delta_w$, the fact that $j_{p,*}^c \circ i_w^* = i_{w,p}^* \circ j_p^c$ (directly from the definitions (6) and (18)) and [BB03, pp. 580] that $j_{p,*}^c \circ \delta_w = \delta_{w,p} \circ j_p^c$. The third equality follows directly from the explicit definition of $\mathfrak{a}_{L_w/\mathbb{Q}_p}$ (see Definition 4.1), and the final one is true since $T_{L_w/\mathbb{Q}_p}^{(2)} = \tilde{\tau}_w$ as $\tau_{\mathbb{Q}_p}^H = 1$.

Now, since $p \cdot \xi$ belongs to $\text{DT}(\mathbb{Z}_p[G])$, Theorem 3.1(i) implies that

$$(p-1)|G| \cdot \xi = (1-1/p)|G| \cdot (p \cdot \xi) = 0.$$

As $e(w)$ is a non-trivial power of p , in order to prove that $(p-1)e(w) \cdot \xi = 0$, by Taylor's Fixed Point Theorem, it is therefore enough to show that the image in $K_0(\mathcal{O}_p^t[G], \mathbb{Q}_p^c[G])$ of the element $e(w) \cdot \xi = (e(w)/p) \cdot (p \cdot \xi)$ is torsion and of order prime to p .

Since the image of U_{L_w/\mathbb{Q}_p} in $K_0(\mathcal{O}_p^t[H], \mathbb{Q}_p^c[H])$ vanishes (see [Bre04a, Lem. 4.4]), the expression (26) reduces to showing that the image in $K_0(\mathcal{O}_p^t[H], \mathbb{Q}_p^c[H])$ of $e(w) \cdot \Delta(\mathcal{A}_{L_w/\mathbb{Q}_p})$ is torsion and of order prime to p .

To prove this, we fix an element a_p of L_w such that $\mathcal{A}_{L_w/\mathbb{Q}_p} = \mathbb{Z}_p[H] \cdot a_p$. Then, since the element δ_F (defined in (23)) is equal to 1 when $F = \mathbb{Q}_p$, the formula [Bre04a, Lem. 4.16] of Breuning implies that, in $K_0(\mathbb{Z}_p[H], \mathbb{Q}_p^c[H])$,

one has

$$\Delta(\mathcal{A}_{L_w/\mathbb{Q}_p}) = \delta_{w,p} \left(\sum_{\chi \in \hat{H}_p} (a_p | \chi) \cdot e_\chi \right).$$

We next note that, by the result [Vin03, Prop. 2.7] of Viatier, there exists a finite unramified extension E of \mathbb{Q}_p such that if q is the order of the residue field of E , for all $\chi \in \hat{H}_p$, one has

$$(a_p | \chi)^{(q-1)e(w)} \in \text{Det}_\chi(\mathcal{O}_E[H]^\times).$$

It follows that, in terms of [Fro83, Chap. II, §2, Lem. 2.1 and §1, Lem. 1.6], one has

$$\left(\sum_{\chi \in \hat{H}_p} (a_p | \chi) \cdot e_\chi \right)^{(q-1)e(w)} \in \text{Nrd}_{\mathbb{Q}_p^c[H]}(K_1(\mathcal{O}_E[H])).$$

In particular, since the image of this element under $\delta_{w,p}$ represents $(q-1)e(w) \cdot \Delta(\mathcal{A}_{L_w/\mathbb{Q}_p})$, and the composite homomorphism

$$K_1(\mathcal{O}_E[H]) \rightarrow K_1(\mathbb{Q}_p^c[H]) \rightarrow K_0(\mathcal{O}_p^t[H], \mathbb{Q}_p^c[H])$$

is zero (as E/\mathbb{Q}_p is unramified), we deduce that the image in $K_0(\mathcal{O}_p^t[H], \mathbb{Q}_p^c[H])$ of $e(w) \cdot \Delta(\mathcal{A}_{L_w/\mathbb{Q}_p})$ is torsion, and of order dividing $q-1$.

Since $q-1$ is prime to p , this implies the result we desire and hence, completes the proof. \square

We shall finish this article by making some remarks concerning the class of $[\mathcal{A}_{L/\mathbb{Q}}]$ in the class group.

Remark 5.2. If we project the result of Theorem 5.1 into the class group, the bound we obtain on the order of $[\mathcal{A}_{L/\mathbb{Q}}]$ is weaker than the one proved by Viatier in [Vin03, Th. 1]. This is due to the fact that, in the class group, one knows that the class of $[\mathcal{A}_{L/\mathbb{Q}}]$ belongs to the kernel subgroup $D(\mathbb{Z}[G])$, which is again a p -group if G has p -powered order (cf. [Vin03, pp. 404]). However, this result is not shared by the torsion subgroup $DT(\mathbb{Z}[G])$, and therefore we are unable to obtain a sharper bound in $K_0(\mathbb{Z}[G], \mathbb{Q}^c[G])$.

Corollary 5.3. *Under the hypothesis of Theorem 5.1, one has that $e(w) \cdot [\mathcal{A}_{L/\mathbb{Q}}] = 0$ in the class group. Moreover, if L_w/\mathbb{Q}_p is totally ramified, then the order of $[\mathcal{A}_{L/\mathbb{Q}}]$ is bounded by $e(w)/p$.*

Proof. The first claim will be proved by combining the results of Theorem 1.3 and 5.1. We first note that $i_w^* \circ \delta_w = \delta \circ \tilde{i}_w = \partial_{\mathbb{Z}, \mathbb{Q}^c, G}^1 \circ (\text{Nrd}_{\mathbb{Q}^c[G]})^{-1} \circ \tilde{i}_w$. Then the image in $\text{Cl}(\mathbb{Z}[G])$ of the correction term $i_w^*(\delta_w((\psi_{2,*} - 1)(\tau'_{L_w/\mathbb{Q}_p})))$ in Theorem 5.1 vanishes since $\partial_{\mathbb{Z}, \mathbb{Q}^c, G}^0 \circ \partial_{\mathbb{Z}, \mathbb{Q}^c, G}^1$ is a zero homomorphism (by the exactness of (5)). We also recall from [BBH20, Th. 5.2(iv)] that $\partial_{\mathbb{Z}, \mathbb{Q}^c, G}^0(\mathfrak{a}_{L/\mathbb{Q}}) = [\mathcal{A}_{L/\mathbb{Q}}]$ and therefore deduce that, under the surjective map $K_0(\mathbb{Z}[G], \mathbb{Q}^c[G]) \rightarrow \text{Cl}(\mathbb{Z}[G])$, the order of $[\mathcal{A}_{L/\mathbb{Q}}]$ is bounded by

$$\gcd(n(L/K)/p, (p-1)e(w)) = e(w).$$

In this way, we obtain the result of Vinatier.

Now suppose that $|H| = e(w)$, the argument above Remark 4.9 implies that $e(w)/p \cdot \mathfrak{a}_{L/\mathbb{Q}} = 0$ in $K_0(\mathbb{Z}[G], \mathbb{Q}^c[G])$ and so claim (ii) is true. \square

References

- [BB03] BLEY, W.; BURNS, D. Equivariant epsilon constants, discriminants and étale cohomology. *Proc. London Math. Soc.* **87** (2003) 545-590. MR2005875, Zbl 1056.11071, doi: 10.1112/S0024611503014217. 423, 437
- [BBH20] BLEY, WERNER; BURNS, DAVID; HAHN, CARL. On refined metric and hermitian structures in arithmetic, I: Galois-Gauss sums and weak ramification. *Ann. K-theory* **5** (2020) 79-140. MR4078226, Zbl 1451.11121, arXiv:1806.02235, doi: 10.2140/akt.2020.5.79. 418, 419, 424, 429, 430, 432, 435, 438
- [BW09] BLEY, WERNER; WILSON, STEPHEN M. J. Computations in relative algebraic K -groups. *London Math. Soc. J. Comput. Math.* **12** (2009) 166-194. MR2564571, Zbl 1225.16003, doi: 10.1112/S146115700001480. 420, 425, 426
- [BK90] BLOCH, SPENCER; KATO, KAZUYA. L -functions and Tamagawa numbers of motives. The Grothendieck Festschrift, Vol. I. *Prog. in Math* **86** (1990) 333-400. MR1086888, Zbl 0768.14001, doi: 10.1007/978-0-8176-4574-8_9. 419
- [Bre04a] BREUNING, MANUEL. Epsilon Constants for Galois extensions of Number Fields and p -adic fields. PhD thesis. *King's College London*, 2004. Available at <https://sites.google.com/site/manuelbreuning/research>. 433, 437
- [Bre04b] BREUNING, MANUEL. Equivariant local epsilon constants and étale cohomology. *J. London Math. Soc.* (2) **70** (2004) 289-306. MR2078894, Zbl 1068.11075, doi: 10.1112/s002461070400554x. 429
- [Bur04] BURNS, DAVID. Equivariant Whitehead torsion and refined Euler characteristics. *CRM Proc. Lecture Notes* **36** (2004) 35-59. MR2076565, Zbl 1070.19003, doi: 10.1090/crpm/036/04. 420
- [BF01] BURNS, D.; FLACH, M. Tamagawa numbers for motives with (non-commutative) coefficients. *Doc. Math.* **6** (2001) 501-570. MR1884523, Zbl 1052.11077, doi: 10.4171/dm/113. 419
- [CV16] CAPUTO, LUCA; VINATIER, STÉPHANE. Galois module structure of the square root of the inverse different in even degree tame extensions of number fields. *J. Algebra* **468** (2016) 103-154. MR3550860, Zbl 1407.11129, doi: 10.1016/j.jalgebra.2016.06.035. 420
- [CRW92] CLIFF, GERALD; RITTER, JÜRGEN; WEISS, ALFRED. Group representations and integrality. *J. Reine Angew. Math.* **426** (1992) 193-202. MR1155753, Zbl 0739.20002, doi: 10.1515/crll.1992.426.193. 427
- [CR81] CURTIS, CHARLES W.; REINER, IRVING. Methods of representation theory. Vol. I. *John Wiley & Sons, Inc., New York*, 1981. xxi+819 pp. ISBN: 0-471-18994-4. MR632548, Zbl 469.20001. 423, 424, 425
- [CR87] CURTIS, CHARLES W.; REINER, IRVING. Methods of representation theory. Vol. II. *John Wiley & Sons, Inc., New York*, 1987. xviii+951 pp. ISBN: 0-471-88871-0. MR892316, Zbl 0616.20001. 420, 422, 423, 425, 426, 427
- [Ere91] EREZ, BOAS. The Galois structure of the square root of the inverse different. *Math. Z.* **208** (1991) 239-255. MR1128708, Zbl 0713.11078, doi: 10.1007/BF02571523. 418, 420, 429, 434
- [ET92] EREZ, B.; TAYLOR, M. J. Hermitian modules in Galois extensions of number fields and Adams operations. *Ann. Math.* (2) **135** (1992) 271-296. MR1154594, Zbl 0756.11035, doi: 10.2307/2946590. 418
- [Fro83] FRÖHLICH, ALBRECHT. Galois Structure of Algebraic Integers. *Ergebnisse Math.* 3. *Springer-Verlag, Berlin*, 1983. x+262 pp. ISBN: 3-540-11920-5. MR717033, Zbl 0501.12012, doi: 10.1007/978-3-642-68816-4. 417, 424, 432, 434, 436, 438

- [Gas54] GASCHÜTZ, WOLFGANG. Endliche Gruppen mit treuen absolut-irreduziblen Darstellungen. *Math. Nachr.* **12** (1954) 253-255. MR67115, Zbl 0056.25703, doi: 10.1002/mana.19540120308. 428
- [Hah16] HAHN, CARL. On the square root of the inverse different via relative algebraic K-theory. PhD thesis. *King's College London*, 2016. 418
- [Neu79] NEUKIRCH, JÜRGEN. On solvable number fields. *Invent. Math.* **53** (1979) 135-164. MR560411, Zbl 0447.12008, doi: 10.1007/BF01390030. 434
- [PT16] PICKETT, ERIK JARL; THOMAS, LARA. Formal group exponentials and Galois modules in Lubin-Tate extensions. *Int. Math. Res. Not. IMRN* (2016) 3205-3236. MR3556410, Zbl 1404.11128, arXiv:1201.4023, doi: 10.1093/imrn/rnv224. 418
- [PV13] PICKETT, ERIK JARL; VINATIER, STÉPHANE. Self-dual integral normal bases and Galois module structure. *Compos. Math.* **149** (2013) 1175-1202. MR3078643, Zbl 1292.11128, doi: 10.1112/S0010437X12000851. 418, 420
- [Roq58] ROQUETTE, PETER. Realisierung von Darstellungen endlicher nilpotenter Gruppen. *Arch. Math. (Basel)* **9** (1958) 241-250. MR97452, Zbl 0083.25002, doi: 10.1007/BF01900587. 427
- [Ser79] SERRE, JEAN-PIERRE. Local Fields. Translated from the French by Marvin Jay Greenberg. Graduate Texts in Math., 67, *Springer, New York*, 1979. viii+241 pp. ISBN: 0-387-90424-7. MR554237, Zbl 0423.12016, doi: 10.1007/978-1-4757-5673-9. 432
- [Swa68] SWAN, R. G. Algebraic K-theory. Lect. Notes in Math., 76, *Springer-Verlag, Berlin-New York*, 1968. iv+262 pp. ISSN: 0075-8434. MR0245634, Zbl 0193.34601, doi: 10.1007/bfb0080281. 422, 423
- [Mar84] TAYLOR, MARTIN. Classgroups of group rings. London Mathematical Society Lecture Note Series, 91, *Cambridge University Press, Cambridge*, 1984. xiii+119 pp. ISBN: 0-521-27870-8. MR748670, Zbl 0597.13002, doi: 10.1017/CBO9781107325555. 433
- [Ull74] ULLOM, STEPHEN V. The exponent of class groups. *J. Algebra* **29** (1974) 124-132. MR338144, Zbl 0278.20016, doi: 10.1016/0021-8693(74)90117-3. 420, 425
- [Vin02] VINATIER, STÉPHANE. Une famille infinie d'extensions faiblement ramifiées. *Math. Nachr.* **243** (2002) 165-187. MR1923839, Zbl 1056.11061, doi: 10.1002/1522-2616(200209)243:1<165::AID-MANA165>3.0.CO;2-K. 420, 431
- [Vin03] VINATIER, STÉPHANE. Sur la racine carrée de la codifférente. *J. Théor. Nombres Bordeaux* **15** (2003) 393-410. MR2019023, Zbl 1048.11088, doi: 10.5802/jtnb.409. 418, 420, 421, 435, 436, 438
- [Vin05] VINATIER, STÉPHANE. Galois module structure in weakly ramified 3-extensions. *Acta Arith.* **119** (2005) 171-186. MR2167720, Zbl 1075.11071, doi: 10.4064/aa119-2-3. 418, 421, 435

(Yu Kuang) 1135 JIUZHOU DADA0, ZHUHAI, CHINA
 yu.kuang.3@gmail.com

This paper is available via <http://nyjm.albany.edu/j/2023/29-18.html>.