

On a twisted Jacquet module of $\mathrm{GL}(6, F)$ over a finite field

Kumar Balasubramanian and Himanshi Khurana

ABSTRACT. Let F be a finite field and $G = \mathrm{GL}(6, F)$. In this paper, we give an explicit description of a certain twisted Jacquet module of an irreducible cuspidal representation of G .

CONTENTS

1. Introduction	874
2. Preliminaries	876
3. Dimension of the twisted Jacquet module	877
4. Main theorem	882
Acknowledgements	909
References	909

1. Introduction

Let F be a finite field and $G = \mathrm{GL}(n, F)$. Let P be a parabolic subgroup of G with Levi decomposition $P = MN$. Let π be any irreducible finite dimensional complex representation of G and ψ be an irreducible representation of N . Let $\pi_{N, \psi}$ be the sum of all irreducible representations of N inside π , on which π acts via ψ . It is easy to see that $\pi_{N, \psi}$ is a representation of the subgroup M_ψ of M , consisting of those elements in M which leave the isomorphism class of ψ invariant under the inner conjugation action of M on N . The space $\pi_{N, \psi}$ is called the *twisted Jacquet module* of the representation π . It is an interesting question to understand for which irreducible representations π , the twisted Jacquet module $\pi_{N, \psi}$ is non-zero and to understand its structure as a module for M_ψ .

In an earlier work of ours [1], inspired by the work of Prasad in [4], we studied the structure of a certain twisted Jacquet module of a cuspidal representation of $\mathrm{GL}(4, F)$. Recently, we have realised that the structure of the module for $\mathrm{GL}(2n, F)$ can also be studied using the more sophisticated theory of

Received January 21, 2023.

2020 *Mathematics Subject Classification.* 20G40 (Primary); 20G05 (Secondary).

Key words and phrases. Cuspidal representations, Twisted Jacquet module.

Research of the corresponding author Kumar Balasubramanian is supported by the SERB grant: MTR/2019/000358.

derivatives of the p -adic group $\mathrm{GL}(n)$ studied in the classical paper of Bernstein-Zelevinskii (see [2]). In this paper, we continue our study of the twisted Jacquet module for a cuspidal representation of $\mathrm{GL}(6, F)$ and use elementary methods from linear algebra to calculate its structure. The calculations in the $\mathrm{GL}(6, F)$ case are much more involved than in the case of $\mathrm{GL}(4, F)$ and we hope that some of these ideas could be used to give an alternative proof in the $\mathrm{GL}(2n, F)$ case as well, using simple techniques from linear algebra. We refer the reader to Section 1 in [1] for a more elaborate introduction to the problem.

Before we state our result, we set up some notation. Let $G = \mathrm{GL}(6, F)$ and P be the maximal parabolic subgroup of G with Levi decomposition $P = MN$, where $M \simeq \mathrm{GL}(3, F) \times \mathrm{GL}(3, F)$ and $N \simeq \mathrm{M}(3, F)$. We write F_6 for the unique field extension of F of degree 6. Let ψ_0 be a fixed non-trivial additive character of F . Let

$$A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

and $\psi_A : N \rightarrow \mathbb{C}^\times$ be the character of N given by

$$\psi_A \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) = \psi_0(\mathrm{Tr}(AX)). \quad (1.1)$$

Let $H_A = M_1 \times M_2$ where M_1 is the Mirabolic subgroup of $\mathrm{GL}(3, F)$ and $M_2 = w_0 M_1^\top w_0^{-1}$ where $w_0 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$. Let U be the subgroup of unipotent matrices in $\mathrm{GL}(6, F)$ and $U_A = U \cap H_A$. Clearly, we have $U_A \simeq U_1 \times U_2$ where U_1 and U_2 are the upper triangular unipotent subgroups of $\mathrm{GL}(3, F)$. For $k = 1, 2$, let $\mu_k : U_k \rightarrow \mathbb{C}^\times$ be the non-degenerate character of U_k given by

$$\mu_k \left(\begin{bmatrix} 1 & x_{12} & x_{13} \\ 0 & 1 & x_{23} \\ 0 & 0 & 1 \end{bmatrix} \right) = \psi_0(x_{12} + x_{23}).$$

Let $\mu : U_A \rightarrow \mathbb{C}^\times$ be the character of U_A given by

$$\mu(u) = \mu_1(u_1)\mu_2(u_2)$$

$$\text{where } u = \begin{bmatrix} u_1 & 0 \\ 0 & u_2 \end{bmatrix}.$$

Theorem 1.1. *Let θ be a regular character of F_6^\times and $\pi = \pi_\theta$ be an irreducible cuspidal representation of G . Then*

$$\pi_{N, \psi_A} \simeq \theta|_{F^\times} \otimes \mathrm{ind}_{U_A}^{H_A} \mu$$

as M_{ψ_A} modules.

We establish the above isomorphism by explicitly calculating the characters of π_{N,ψ_A} and $\theta|_{F^\times} \otimes \text{ind}_{U_A}^{H_A}(\mu)$, and showing that they are equal at any arbitrary element of M_{ψ_A} .

2. Preliminaries

In this section, we mention some preliminary results that we need in our paper.

2.1. Character of a cuspidal representation. Let F be the finite field of order q and $G = \text{GL}(m, F)$. Let F_m be the unique field extension of F of degree m . A character θ of F_m^\times is called a “regular” character, if under the action of the Galois group of F_m over F , θ gives rise to m distinct characters of F_m^\times . It is a well known fact that the cuspidal representations of $\text{GL}(m, F)$ are parametrized by the regular characters of F_m^\times . To avoid introducing more notation, we mention below only the relevant statements on computing the character values that we have used. We refer the reader to Section 6 in [3] for more precise statements on computing character values.

Theorem 2.1. *Let θ be a regular character of F_m^\times . Let $\pi = \pi_\theta$ be an irreducible cuspidal representation of $\text{GL}(m, F)$ associated to θ . Let Θ_θ be its character. If $g \in \text{GL}(m, F)$ is such that the characteristic polynomial of g is not a power of a polynomial irreducible over F . Then, we have*

$$\Theta_\theta(g) = 0.$$

Theorem 2.2. *Let θ be a regular character of F_m^\times . Let $\pi = \pi_\theta$ be an irreducible cuspidal representation of $\text{GL}(m, F)$ associated to θ . Let Θ_θ be its character. Suppose that $g = s.u$ is the Jordan decomposition of an element g in $\text{GL}(m, F)$. If $\Theta_\theta(g) \neq 0$, then the semisimple element s must come from F_m^\times . Suppose that s comes from F_m^\times . Let z be an eigenvalue of s in F_m and let t be the dimension of the kernel of $g - z$ over F_m . Then*

$$\Theta_\theta(g) = (-1)^{m-1} \left[\sum_{\alpha=0}^{d-1} \theta(z^{q^\alpha}) \right] (1 - q^d)(1 - (q^d)^2) \cdots (1 - (q^d)^{t-1}).$$

where q^d is the cardinality of the field generated by z over F , and the summation is over the distinct Galois conjugates of z .

See Theorem 2 in [4] for this version.

2.2. Twisted Jacquet module. In this section, we recall the character and the dimension formula of the twisted Jacquet module of a representation π .

Let $G = \text{GL}(k, F)$ and $P = MN$ be a parabolic subgroup of G . Let ψ be a character of N . For $m \in M$, let ψ^m be the character of N defined by $\psi^m(n) = \psi(mnm^{-1})$. Let

$$V(N, \psi) = \text{Span}_{\mathbb{C}} \{ \pi(n)v - \psi(n)v \mid n \in N, v \in V \}$$

and

$$M_\psi = \{m \in M \mid \psi^m(n) = \psi(n), \forall n \in N\}.$$

Clearly, M_ψ is a subgroup of M and it is easy to see that $V(N, \psi)$ is an M_ψ -invariant subspace of V . Hence, we get a representation $(\pi_{N, \psi}, V/V(N, \psi))$ of M_ψ . We call $(\pi_{N, \psi}, V/V(N, \psi))$ the twisted Jacquet module of π with respect to ψ . We write $\Theta_{N, \psi}$ for the character of $\pi_{N, \psi}$.

Proposition 2.3. *Let (π, V) be a representation of $\mathrm{GL}(k, F)$ and Θ_π be the character of π . We have*

$$\Theta_{N, \psi}(m) = \frac{1}{|N|} \sum_{n \in N} \Theta_\pi(mn) \overline{\psi(n)}.$$

We refer the reader to Proposition 2.3 in [1] for a proof.

Remark 2.4. Taking $m = 1$, we get the dimension of $\pi_{N, \psi}$. To be precise, we have

$$\dim_{\mathbb{C}}(\pi_{N, \psi}) = \frac{1}{|N|} \sum_{n \in N} \Theta_\pi(n) \overline{\psi(n)}.$$

2.3. Character of the induced representation. In this section, we recall the character formula for the induced representation of a group G . For a proof, we refer the reader to Chapter 3, Theorem 12 in [5].

Proposition 2.5. *Let G be a finite group and H be a subgroup of G . Let (π, V) be a representation of H and χ_π be the character of π . Then for each $s \in G$, the character of $\mathrm{ind}_H^G(\pi)$ is given by*

$$\chi_{\mathrm{ind}_H^G(\pi)}(s) = \frac{1}{|H|} \sum_{\substack{t \in G \\ t^{-1}st \in H}} \chi_\pi(t^{-1}st).$$

3. Dimension of the twisted Jacquet module

Let $\pi = \pi_\theta$ be an irreducible cuspidal representation of G corresponding to the regular character θ of F_6^\times and Θ_θ be its character. Throughout, we write $M(n, m, r, q)$ for the set of $n \times m$ matrices of rank r over the finite field $F = F_q$. In this section, we calculate the dimension of π_{N, ψ_A} . Before we continue, we record some preliminary lemmas that we need.

Lemma 3.1. *Let $r \in \{0, 1, 2, 3\}$ and $X \in M(3, 3, r, q)$. We have*

$$\Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) = \begin{cases} (q-1)(q^2-1)(q^3-1)(q^4-1)(q^5-1), & \text{if } r=0 \\ -(q-1)(q^2-1)(q^3-1)(q^4-1), & \text{if } r=1 \\ (q-1)(q^2-1)(q^3-1), & \text{if } r=2 \\ -(q-1)(q^2-1), & \text{if } r=3 \end{cases}$$

Proof. The result follows from Theorem 2.2 above. \square

Let

$$X = \begin{bmatrix} a & d & g \\ b & e & h \\ c & f & k \end{bmatrix}, A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, AX = \begin{bmatrix} a & d & g \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

For $\alpha \in F$ and $r \in \{0, 1, 2, 3\}$, consider the subset $Y_{3,r}^\alpha$ of $M(3, F)$ given by

$$Y_{3,r}^\alpha = \{X \in M(3, F) \mid \text{Rank}(X) = r, \text{Tr}(AX) = \alpha\}.$$

Lemma 3.2. *Let $r \in \{1, 2, 3\}$ and $\alpha, \beta \in F^\times$. Then we have*

$$\#Y_{3,r}^\alpha = \#Y_{3,r}^\beta.$$

Proof. Consider the map $\phi : Y_{3,r}^\alpha \rightarrow Y_{3,r}^\beta$ given by

$$\phi(X) = \alpha^{-1}\beta X.$$

Suppose that $\phi(X) = \phi(Y)$. Since $\alpha^{-1}\beta \neq 0$, it follows that ϕ is injective. For $Y \in Y_{3,r}^\beta$, let $X = \alpha\beta^{-1}Y$. Clearly, we have $\text{Tr}(AX) = \alpha$ and $\text{Rank}(X) = \text{Rank}(Y) = r$. Thus, ϕ is surjective and hence the result. \square

Theorem 3.3. *Let θ be a regular character of F_6^\times and $\pi = \pi_\theta$ be an irreducible cuspidal representation of $\text{GL}(6, F)$. We have*

$$\dim_{\mathbb{C}}(\pi_{N, \psi_A}) = (q - 1)^2(q^2 - 1)^2.$$

Proof. It is easy to see that the dimension of π_{N, ψ_A} is given by

$$\dim_{\mathbb{C}}(\pi_{N, \psi_A}) = \frac{1}{q^9} \sum_{X \in M(3, F)} \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \overline{\psi_0(\text{Tr}(AX))}. \quad (3.1)$$

We calculate the following sums

$$(a) \quad S_1 = \sum_{X \in M(3, 3, 0, q)} \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \overline{\psi_0(\text{Tr}(AX))}$$

$$(b) \quad S_2 = \sum_{\substack{X \in M(3, 3, 1, q) \\ \text{Tr}(AX)=0}} \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \overline{\psi_0(\text{Tr}(AX))}$$

$$+ \sum_{\substack{X \in M(3, 3, 1, q) \\ \text{Tr}(AX)=\alpha \neq 0}} \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \overline{\psi_0(\text{Tr}(AX))}$$

$$(c) \quad S_3 = \sum_{\substack{X \in M(3, 3, 2, q) \\ \text{Tr}(AX)=0}} \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \overline{\psi_0(\text{Tr}(AX))}$$

$$+ \sum_{\substack{X \in M(3, 3, 2, q) \\ \text{Tr}(AX)=\alpha \neq 0}} \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \overline{\psi_0(\text{Tr}(AX))}$$

$$(d) \quad S_4 = \sum_{\substack{X \in M(3,3,3,q) \\ \mathrm{Tr}(AX)=0}} \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \overline{\psi_0(\mathrm{Tr}(AX))}$$

$$+ \sum_{\substack{X \in M(3,3,3,q) \\ \mathrm{Tr}(AX)=\alpha \neq 0}} \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \overline{\psi_0(\mathrm{Tr}(AX))}$$

separately to compute the dimension of π_{N,ψ_A} .

For a fixed $r \in \{0, 1, 2, 3\}$ and $\alpha \in \{0, 1\}$, we find a partition of $Y_{3,r}^\alpha$ into certain subsets, and compute the cardinality of each of these subsets to find the cardinality of $Y_{3,r}^\alpha$. We record the necessary information in the tables below.

For (a), we clearly have

$$S_1 = \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \overline{\psi_0(0)}$$

$$= (q-1)(q^2-1)(q^3-1)(q^4-1)(q^5-1).$$

For (b), we have

TABLE 1. $\mathrm{Rank}(X) = 1$

Partition of $Y_{3,1}^0$	Cardinality	Partition of $Y_{3,1}^1$	Cardinality
$\left\{ \begin{bmatrix} 0 & 0 & 0 \\ b & \lambda b & \beta b \\ c & \lambda c & \beta c \end{bmatrix} \right\}$	$(q^2-1)q^2$	$\left\{ \begin{bmatrix} 1 & \lambda & \beta \\ b & \lambda b & \beta b \\ c & \lambda c & \beta c \end{bmatrix} \right\}$	q^4
$\left\{ \begin{bmatrix} 0 & d & \lambda d \\ 0 & e & \lambda e \\ 0 & f & \lambda f \end{bmatrix} \right\}$	$(q^3-1)q$	-	-
$\left\{ \begin{bmatrix} 0 & 0 & g \\ 0 & 0 & h \\ 0 & 0 & k \end{bmatrix} \right\}$	q^3-1	-	-

Thus,

$$S_2 = \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \left(\sum_{\substack{X \in M(3,3,1,q) \\ \mathrm{Tr}(AX)=0}} \overline{\psi_0(0)} + \sum_{\substack{X \in M(3,3,1,q) \\ \mathrm{Tr}(AX)=\alpha \neq 0}} \overline{\psi_0(\alpha)} \right)$$

$$\begin{aligned}
&= \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) (\#Y_{3,1}^0 - \#Y_{3,1}^1) \\
&= -(q-1)(q^2-1)(q^3-1)(q^4-1)((q^2-1)q^2 + (q^3-1)q + (q^3-1) - q^4) \\
&= -(q-1)^2(q^2-1)^2(q^8 + 2q^7 + 2q^6 + q^5 - 2q^4 - 3q^3 - 4q^2 - 2q - 1).
\end{aligned}$$

For (d), we have

TABLE 2. $\text{Rank}(X) = 3$

Partition of $Y_{3,3}^0$	Cardinality	Partition of $Y_{3,3}^1$	Cardinality
$\left\{ \begin{bmatrix} 0 & d & g \\ b & e & h \\ c & f & k \end{bmatrix} \right\}$	$(q^2-1)(q^3-q)(q^3-q^2)$	$\left\{ \begin{bmatrix} 1 & d & g \\ b & e & h \\ c & f & k \end{bmatrix} \right\}$	$q^2(q^3-q)(q^3-q^2)$

Thus,

$$\begin{aligned}
S_4 &= \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \left(\sum_{\substack{X \in M(3,3,3,q) \\ \text{Tr}(AX)=0}} \overline{\psi_0(0)} + \sum_{\substack{X \in M(3,3,3,q) \\ \text{Tr}(AX)=\alpha \neq 0}} \overline{\psi_0(\alpha)} \right) \\
&= \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) (\#Y_{3,3}^0 - \#Y_{3,3}^1) \\
&= -(q-1)(q^2-1)((q^2-1)(q^3-q)(q^3-q^2) - q^2(q^3-q)(q^3-q^2)) \\
&= (q-1)^2(q^2-1)^2q^3.
\end{aligned}$$

For (c), we let $X' = \begin{bmatrix} e & h \\ f & k \end{bmatrix}$. For $\alpha \in \{0, 1\}$, we partition the set $Y_{3,2}^\alpha$ according to the rank of X' and count the cardinalities of each of these subsets. For $\text{Rank}(X') \in \{0, 1, 2\}$ and $\alpha \in \{0, 1\}$, we record the cardinality of such subsets of $Y_{3,2}^\alpha$ in the following tables.

TABLE 3. $\mathrm{Rank}(X) = 2, \mathrm{Rank}(X') = 0$

Partition of $Y_{3,2}^0$	Cardinality	Partition of $Y_{3,2}^1$	Cardinality
$B_1 = \left\{ \begin{bmatrix} 0 & d & g \\ b & 0 & 0 \\ c & 0 & 0 \end{bmatrix} \mid (d, g) \neq (0, 0) \right\}$	$(q^2 - 1)^2$	$B_2 = \left\{ \begin{bmatrix} 1 & d & g \\ b & 0 & 0 \\ c & 0 & 0 \end{bmatrix} \mid (d, g) \neq (0, 0) \right\}$	$(q^2 - 1)^2$

TABLE 4. $\mathrm{Rank}(X) = 2, \mathrm{Rank}(X') = 2$

Partition of $Y_{3,2}^0$	Cardinality	Partition of $Y_{3,2}^1$	Cardinality
$C_1 = \left\{ \begin{bmatrix} 0 & e & h \\ \lambda e + \beta h & e & h \\ \lambda f + \beta k & f & k \end{bmatrix} \right\}$	$(q^2 - 1)(q^2 - q)(q)$	$C_4 = \left\{ \begin{bmatrix} 1 & d & 0 \\ d^{-1}e + \beta h & e & h \\ d^{-1}f + \beta k & f & k \end{bmatrix}, \begin{bmatrix} 1 & 0 & g \\ \lambda e + g^{-1}h & e & h \\ \lambda f + g^{-1}k & f & k \end{bmatrix} \right\}$	$2(q^2 - 1)(q^2 - q)^2$
$C_2 = \left\{ \begin{bmatrix} 0 & d & 0 \\ \beta h & e & h \\ \beta k & f & k \end{bmatrix}, \begin{bmatrix} 0 & e & g \\ \beta e & f & h \\ \beta f & g & k \end{bmatrix} \right\}$	$2(q^2 - 1)(q^2 - q)^2$	$C_5 = \left\{ \begin{bmatrix} 1 & d & g \\ d^{-1}(1 - \beta g)e + \beta h & e & h \\ d^{-1}(1 - \beta g)f + \beta k & f & k \end{bmatrix} \right\}$	$(q^2 - 1)(q^3 - q^2)(q - 1)^2$
$C_3 = \left\{ \begin{bmatrix} 0 & d & g \\ \beta(-gd^{-1}e + h) & e & h \\ \beta(-gd^{-1}f + k) & f & k \end{bmatrix} \right\}$	$(q^2 - 1)(q^2 - q)^2(q - 1)$	-	-

TABLE 5. $\mathrm{Rank}(X) = 2, \mathrm{Rank}(X') = 1$

Partition of $Y_{3,2}^0$	Cardinality	Partition of $Y_{3,2}^1$	Cardinality
$E_1 = \left\{ \begin{bmatrix} 0 & 0 & 0 \\ b & e & 0 \\ c & f & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ b & 0 & h \\ c & 0 & k \end{bmatrix} \right\}$	$2(q^2 - 1)(q^2 - q)$	$F_1 = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ b & 0 & h \\ c & 0 & k \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ b & e & 0 \\ c & f & 0 \end{bmatrix} \right\}$	$2(q^2 - 1)q^2$
$E_2 = \left\{ \begin{bmatrix} 0 & d & 0 \\ b & e & 0 \\ c & f & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & g \\ b & 0 & h \\ c & 0 & k \end{bmatrix} \right\}$	$2(q^2 - 1)^2(q - 1)$	$F_2 = \left\{ \begin{bmatrix} 1 & d & 0 \\ \beta h & 0 & h \\ \beta k & 0 & k \end{bmatrix}, \begin{bmatrix} 1 & 0 & g \\ \beta e & 0 & h \\ \beta f & f & 0 \end{bmatrix} \right\}$	$2(q^2 - 1)(q^2 - q)$
$E_3 = \left\{ \begin{bmatrix} 0 & 0 & g \\ \lambda e & e & 0 \\ \lambda f & f & 0 \end{bmatrix}, \begin{bmatrix} 0 & d & 0 \\ \lambda h & 0 & h \\ \lambda k & 0 & k \end{bmatrix} \right\}$	$2(q^2 - 1)(q^2 - q)$	$F_3 = \left\{ \begin{bmatrix} 1 & 0 & g \\ b & 0 & h \\ c & 0 & k \end{bmatrix} \mid (b, c) \neq (g^{-1}h, g^{-1}k), \begin{bmatrix} 1 & d & 0 \\ b & e & 0 \\ c & f & 0 \end{bmatrix} \mid (b, c) \neq (d^{-1}e, d^{-1}f) \right\}$	$2(q^2 - 1)^2(q - 1)$
$E_4 = \left\{ \begin{bmatrix} 0 & d & g \\ -d^{-1}\beta ge & e & h \\ -d^{-1}\beta gf & f & 0 \end{bmatrix}, \begin{bmatrix} 0 & d & g \\ -g^{-1}\beta dh & 0 & h \\ -g^{-1}\beta dk & 0 & k \end{bmatrix} \right\}$	$2(q^3 - q)(q - 1)^2$	$F_4 = \left\{ \begin{bmatrix} 1 & d & g \\ \beta h & 0 & h \\ \beta k & 0 & k \end{bmatrix}, \begin{bmatrix} 1 & d & g \\ \beta e & e & 0 \\ \beta f & f & 0 \end{bmatrix} \right\}$	$2q(q^2 - 1)(q - 1)^2$
$E_5 = \left\{ \begin{bmatrix} 0 & 0 & 0 \\ b & \lambda h & h \\ c & \lambda k & k \end{bmatrix} \mid \lambda \neq 0 \right\}$	$(q^2 - 1)(q - 1)(q^2 - q)$	$F_5 = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ b & e & \lambda e \\ c & f & \lambda f \end{bmatrix} \mid \lambda \neq 0 \right\}$	$(q^2 - 1)q^2(q - 1)$
$E_6 = \left\{ \begin{bmatrix} 0 & d & 0 \\ \beta h & \lambda h & h \\ \beta k & \lambda k & k \end{bmatrix} \mid \lambda \neq 0 \right\}$	$2q(q^2 - 1)(q - 1)^2$	$F_6 = \left\{ \begin{bmatrix} 1 & d & 0 \\ d^{-1}e + \beta h & e & \lambda e \\ d^{-1}f + \beta k & f & \lambda f \end{bmatrix}, \begin{bmatrix} 1 & 0 & g \\ \delta e + g^{-1}\lambda e & 0 & \lambda e \\ \delta f + g^{-1}\lambda f & f & \lambda f \end{bmatrix} \mid \lambda \neq 0 \right\}$	$2q(q^2 - 1)(q - 1)^2$
$E_7 = \left\{ \begin{bmatrix} 0 & d & g \\ -d^{-1}\beta g\lambda h + \beta h & \lambda h & h \\ -d^{-1}\beta g\lambda k + \beta k & \lambda k & k \end{bmatrix} \mid d \neq \lambda g, \lambda \neq 0 \right\}$	$(q^2 - 1)(q - 1)^2(q^2 - 2q)$	$F_7 = \left\{ \begin{bmatrix} 1 & d & g \\ d^{-1}(1 - \beta g)e + \beta h & e & \lambda e \\ d^{-1}(1 - \beta g)f + \beta k & f & \lambda f \end{bmatrix} \mid g \neq \lambda d, \lambda \neq 0 \right\}$	$(q^2 - 1)(q - 1)^2(q^2 - 2q)$
$E_8 = \left\{ \begin{bmatrix} 0 & \lambda g & g \\ b & \lambda h & h \\ c & \lambda k & k \end{bmatrix} \mid \lambda \neq 0 \right\}$	$(q^2 - 1)^2(q - 1)^2$	$F_8 = \left\{ \begin{bmatrix} 1 & d & \lambda d \\ b & e & \lambda e \\ c & f & \lambda f \end{bmatrix} \mid \lambda \neq 0 \right\}$	$(q^2 - 1)^2(q - 1)^2$

We have,

$$Y_{3,2}^0 = B_1 \bigsqcup_{i=1}^3 C_i \bigsqcup_{j=1}^8 E_j$$

and

$$Y_{3,2}^1 = B_2 \bigsqcup_{i=4}^5 C_i \bigsqcup_{j=1}^8 F_j.$$

Thus,

$$\begin{aligned} S_3 &= \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) \left(\sum_{\substack{X \in M(3,3,2,q) \\ \text{Tr}(AX)=0}} \overline{\psi_0(0)} + \sum_{\substack{X \in M(3,3,2,q) \\ \text{Tr}(AX)=\alpha \neq 0}} \overline{\psi_0(\alpha)} \right) \\ &= \Theta_\theta \left(\begin{bmatrix} 1 & X \\ 0 & 1 \end{bmatrix} \right) (\#Y_{3,2}^0 - \#Y_{3,2}^1) \\ &= (q-1)(q^2-1)(q^3-1)(q^6-q^5-2q^4+q^2+q) \\ &= (q-1)^2(q^2-1)^2(q^6-q^4-3q^3-2q^2-q). \end{aligned}$$

From equation (3.1), it follows that

$$\begin{aligned} \dim_{\mathbb{C}}(\pi_{N,\psi_A}) &= \frac{1}{q^9} \{S_1 + S_2 + S_3 + S_4\} \\ &= \frac{1}{q^9} (q-1)^2(q^2-1)^2 q^9 \\ &= (q-1)^2(q^2-1)^2. \end{aligned}$$

□

Remark 3.4. Suppose that $B = Aw_0$. It is easy to see that $\Theta_{N,\psi_A} \left(\begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \right) = \Theta_{N,\psi_B} \left(\begin{bmatrix} w_0 m_1 w_0 & 0 \\ 0 & m_2 \end{bmatrix} \right)$. Thus, we have that $\dim(\pi_{N,\psi_A}) = \dim(\pi_{N,\psi_B})$.

4. Main theorem

In this section, we prove the main result of this paper. Hereafter, we take

$$A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

For the sake of completeness, we recall the statement below.

Theorem 4.1. Let θ be a regular character of F_6^\times and $\pi = \pi_\theta$ be an irreducible cuspidal representation of G . Then

$$\pi_{N,\psi_A} \simeq \theta|_{F^\times} \otimes \mathrm{ind}_{U_A}^{H_A} \mu$$

as M_{ψ_A} modules.

The key idea of the proof is to compute the characters of the representations $\rho = \theta|_{F^\times} \otimes \mathrm{ind}_{U_A}^{H_A} \mu$ and π_{N,ψ_A} and show that they are equal at any arbitrary element in M_{ψ_A} . Before we continue, we set up some notation and record a few lemmas that we need.

Lemma 4.2. Let $M_{\psi_A} = \{m \in M \mid \psi_A^m(n) = \psi_A(n), \forall n \in N\}$. Then we have

$$M_{\psi_A} = \left\{ \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & a \\ & & \begin{bmatrix} a & y_{12} & y_{13} \\ 0 & y_{22} & y_{23} \\ 0 & y_{32} & y_{33} \end{bmatrix} \end{bmatrix} \mid a \in F^\times \right\}.$$

Proof. Let $g = \begin{bmatrix} g_1 & 0 \\ 0 & g_2 \end{bmatrix} \in M$. Then $g \in M_{\psi_A}$ if and only if $Ag_1 = g_2A$. It follows that $g \in M_{\psi_A}$ if and only if $g_1 = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & a \end{bmatrix}$ and $g_2 = \begin{bmatrix} a & y_{12} & y_{13} \\ 0 & y_{22} & y_{23} \\ 0 & y_{32} & y_{33} \end{bmatrix}$. \square

Lemma 4.3. Let $Z = Z(G)$ be the center of G . Then, we have

$$M_{\psi_A} \simeq Z \times H_A.$$

Proof. Trivial. \square

4.1. Character calculation for ρ . Let $\rho_1 = \mathrm{ind}_{U_1}^{M_1} \mu_1$ and $\rho_2 = \mathrm{ind}_{U_2}^{M_2} \mu_2$. In this section, we calculate the character of the representation

$$\rho = \theta|_{F^\times} \otimes \mathrm{ind}_{U_A}^{H_A} \mu \simeq \theta|_{F^\times} \otimes (\rho_1 \otimes \rho_2).$$

4.1.1. Character computation of ρ_1 . Let μ_1 be same as above. Consider the representation

$$\rho_1 = \mathrm{ind}_{U_1}^{M_1} \mu_1$$

of M_1 . Let χ_{ρ_1} be the character of ρ_1 . Let

$$S_1 = \left\{ \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1 \end{bmatrix} \mid a, b \in F^\times \right\} \text{ and } S_2 = \left\{ \begin{bmatrix} p & q & 0 \\ r & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mid p \in F, q, r \in F^\times \right\}.$$

It is easy to see that $S = S_1 \cup S_2$ is a set of left coset representatives of U_1 in M_1 .

Lemma 4.4. *Let*

$$m = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \in M_1 \text{ and } t = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1 \end{bmatrix} \in S_1.$$

Then, $t^{-1}mt \in U_1$ if and only if $a_{11} = a_{22} = 1$ and $a_{21} = 0$. In particular, for $m \in M_1$ with $a_{11} = a_{22} = 1$ and $a_{21} = 0$, we have

$$\sum_{t \in S_1} \mu_1(t^{-1}mt) = \sum_{a,b \in F^\times} \psi_0(a^{-1}ba_{12} + b^{-1}a_{23}).$$

Proof. For $m \in M_1$ and $t \in S_1$, we have

$$t^{-1}mt = \begin{bmatrix} a_{11} & a^{-1}ba_{12} & a^{-1}a_{13} \\ b^{-1}aa_{21} & a_{22} & b^{-1}a_{23} \\ 0 & 0 & 1 \end{bmatrix}.$$

Thus, it follows that $t^{-1}mt \in U_1$ if and only if $a_{11} = a_{22} = 1$ and $a_{21} = 0$. Clearly, we have

$$\sum_{t \in S_1} \mu_1(t^{-1}mt) = \sum_{a,b \in F^\times} \psi_0(a^{-1}ba_{12} + b^{-1}a_{23}),$$

hence the result. \square

Lemma 4.5. *Let*

$$m = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \in M_1 \text{ and } t = \begin{bmatrix} p & q & 0 \\ r & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \in S_2.$$

Suppose that $a_{21} = 0$. Then,

$$t^{-1}mt \in U_1 \text{ if and only if } a_{11} = a_{22} = 1 \text{ and } a_{12} = 0.$$

In particular, for $m \in M_1$ with $a_{11} = a_{22} = 1$ and $a_{21} = a_{12} = 0$, we have

$$\sum_{t \in S_2} \mu_1(t^{-1}mt) = \sum_{\substack{p \in F \\ r,q \in F^\times}} \psi_0(-pq^{-1}r^{-1}a_{23} + q^{-1}a_{13}).$$

Proof. Let $m \in M_1$ and $t \in S_2$. If $a_{21} = 0$, we have,

$$t^{-1}mt = \begin{bmatrix} a_{22} & 0 & r^{-1}a_{23} \\ pq^{-1}(a_{11} - a_{22}) + rq^{-1}a_{12} & a_{11} & q^{-1}a_{13} - pq^{-1}r^{-1}a_{23} \\ 0 & 0 & 1 \end{bmatrix}.$$

Then, $t^{-1}mt \in U_1$ if and only if $a_{22} = a_{11} = 1$ and $a_{12} = 0$. Clearly, we have

$$\sum_{t \in S_2} \mu_1(t^{-1}mt) = \sum_{\substack{p \in F \\ r,q \in F^\times}} \psi_0(-pq^{-1}r^{-1}a_{23} + q^{-1}a_{13}).$$

Hence the result. \square

Lemma 4.6. *Let*

$$m = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \in M_1 \text{ and } t = \begin{bmatrix} p & q & 0 \\ r & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \in S_2.$$

Suppose that $a_{21} \neq 0$.

- a) If $p = 0$, then $t^{-1}mt \in U_1$ if and only if $a_{11} = a_{22} = 1$ and $a_{12} = 0$. In particular, we have

$$\sum_{t \in S_2} \mu_1(t^{-1}mt) = \sum_{r,q \in F^\times} \psi_0(qr^{-1}a_{21} + q^{-1}a_{13}).$$

- b) If $p \neq 0$, then $t^{-1}mt \in U_1$ if and only if $a_{11} + a_{22} = 2$, $a_{12} = \frac{-(a_{11}-1)^2}{a_{21}}$ and $r = (\frac{pa_{21}}{a_{11}-1})$. In particular, we have

$$\sum_{t \in S_2} \mu_1(t^{-1}mt) = - \sum_{q \in F^\times} \psi_0(q^{-1}(-\delta + a_{13})),$$

where $\delta = a_{21}^{-1}a_{23}(a_{11} - 1)$.

Proof. Let

$$m = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \in M_1$$

and suppose that $a_{21} \neq 0$. In case a), since $p = 0$, we have

$$t^{-1}mt = \begin{bmatrix} a_{22} & qr^{-1}a_{21} & r^{-1}a_{23} \\ rq^{-1}a_{12} & a_{11} & q^{-1}a_{13} \\ 0 & 0 & 1 \end{bmatrix}.$$

Thus, it follows that $t^{-1}mt \in U_1$ if and only if $a_{22} = a_{11} = 1$ and $a_{12} = 0$. In particular, we have

$$\sum_{t \in S_2} \mu_1(t^{-1}mt) = \sum_{r,q \in F^\times} \psi_0(qr^{-1}a_{21} + q^{-1}a_{13}).$$

In case b), since $p \neq 0$, we see that $t^{-1}mt$ is equal to

$$\begin{bmatrix} a_{22} + pr^{-1}a_{21} & qr^{-1}a_{21} & r^{-1}a_{23} \\ pq^{-1}(a_{11} - a_{22}) + rq^{-1}a_{12} - p^2q^{-1}r^{-1}a_{21} & a_{11} - pr^{-1}a_{21} & q^{-1}a_{13} - pq^{-1}r^{-1}a_{23} \\ 0 & 0 & 1 \end{bmatrix}.$$

Clearly, we have $t^{-1}mt \in U_1$ if and only if $a_{11} = 1 + pr^{-1}a_{21}$, $a_{22} = 1 - pr^{-1}a_{21}$ and $pq^{-1}(a_{11} - a_{22}) + rq^{-1}a_{12} - p^2q^{-1}r^{-1}a_{21} = 0$. Using $a_{11} - a_{22} = 2pr^{-1}a_{21}$, $a_{11} + a_{22} = 2$ and $\det(t^{-1}mt) = 1$, it follows that $a_{12} = \frac{-(a_{11}-1)^2}{a_{21}}$ and $r = (\frac{pa_{21}}{a_{11}-1})$.

In particular, taking $\delta = a_{21}^{-1}a_{23}(a_{11} - 1)$ we have

$$\sum_{t \in S_2} \mu_1(t^{-1}mt) = - \sum_{q \in F^\times} \psi_0(q^{-1}(-a_{21}^{-1}a_{23}(a_{11} - 1) + a_{13})).$$

$$= - \sum_{q \in F^\times} \psi_0(q^{-1}(-\delta + a_{13})),$$

hence the result. \square

We summarize the character values of ρ_1 in the table below.

TABLE 6. Character of ρ_1

Type of m	m	$\chi_{\rho_1}(m)$	Type of m	m	$\chi_{\rho_1}(m)$
Type-1	$\left\{ \begin{bmatrix} 1 & a_{12} & a_{13} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mid a_{12} \in F^\times \right\}$	$(1-q)$	Type-6	$\left\{ \begin{bmatrix} 1 & 0 & a_{13} \\ 0 & 1 & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \mid a_{13}, a_{23} \in F^\times \right\}$	$(1-q)$
Type-2	$\left\{ \begin{bmatrix} 1 & a_{12} & a_{13} \\ 0 & 1 & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \mid a_{12}, a_{23} \in F^\times \right\}$	1	Type-7	$\left\{ \begin{bmatrix} 1 & 0 & 0 \\ a_{21} & 1 & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \mid a_{21} \in F^\times \right\}$	$(1-q)$
Type-3	$\left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \mid a_{23} \in F^\times \right\}$	$(1-q)$	Type-8	$\left\{ \begin{bmatrix} 1 & 0 & a_{13} \\ a_{21} & 1 & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \mid a_{21}, a_{13} \in F^\times \right\}$	1
Type-4	$\left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right\}$	$(1-q)(1-q^2)$	Type-9	$\left\{ \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \mid \begin{array}{l} a_{21} \in F^\times, a_{13} = \delta \\ a_{12} = -a_{21}^{-1}(a_{11} - 1)^2 \end{array} \right\}$	$(1-q)$
Type-5	$\left\{ \begin{bmatrix} 1 & 0 & a_{13} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mid a_{13} \in F^\times \right\}$	$(1-q)$	Type-10	$\left\{ \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \mid \begin{array}{l} a_{21} \in F^\times, a_{13} \neq \delta \\ a_{12} = -a_{21}^{-1}(a_{11} - 1)^2 \end{array} \right\}$	1

If $m \in M_1$ is not one of the types mentioned in Table 6, then $\chi_{\rho_1}(m) = 0$.

4.1.2. Character computation of ρ_2 . Let μ_2 be same as above. Consider the representation

$$\rho_2 = \text{ind}_{U_2}^{M_2} \mu_2$$

of M_2 . Let χ_{ρ_2} be the character of ρ_2 . Let

$$S_3 = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & b \end{bmatrix} \mid a, b \in F^\times \right\}$$

and

$$S_4 = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & p & q \\ 0 & r & 0 \end{bmatrix} \mid p \in F, q, r \in F^\times \right\}.$$

It is easy to see that $S = S_3 \cup S_4$ is a set of left coset representatives of U_2 in M_2 .

Lemma 4.7. *Let*

$$m = \begin{bmatrix} 1 & y_{12} & y_{13} \\ 0 & y_{22} & y_{23} \\ 0 & y_{32} & y_{33} \end{bmatrix} \in M_2, t = \begin{bmatrix} 1 & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & b \end{bmatrix} \in S_3.$$

Then, $t^{-1}mt \in U_2$ if and only if $y_{22} = y_{33} = 1$ and $y_{32} = 0$. In particular, for $m \in M_2$ with $y_{22} = y_{33} = 1$ and $y_{32} = 0$, we have

$$\sum_{t \in S_3} \mu_2(t^{-1}mt) = \sum_{a,b \in F^\times} \psi_0(ay_{12} + ba^{-1}y_{23}).$$

Proof. Similar to Lemma 4.4. □

Lemma 4.8. *Let*

$$m = \begin{bmatrix} 1 & y_{12} & y_{13} \\ 0 & y_{22} & y_{23} \\ 0 & y_{32} & y_{33} \end{bmatrix} \in M_2 \text{ and } t = \begin{bmatrix} 1 & 0 & 0 \\ 0 & p & q \\ 0 & r & 0 \end{bmatrix} \in S_4.$$

Suppose that $y_{32} = 0$. Then,

$$t^{-1}mt \in U_2 \text{ if and only if } y_{22} = y_{33} = 1 \text{ and } y_{23} = 0.$$

In particular, for $m \in M_2$ with $y_{22} = y_{33} = 1$ and $y_{32} = y_{23} = 0$, we have

$$\sum_{t \in S_4} \mu_2(t^{-1}mt) = \sum_{\substack{p \in F \\ r,q \in F^\times}} \psi_0(py_{12} + ry_{13}).$$

Proof. Similar to Lemma 4.5. □

Lemma 4.9. *Let*

$$m = \begin{bmatrix} 1 & y_{12} & y_{13} \\ 0 & y_{22} & y_{23} \\ 0 & y_{32} & y_{33} \end{bmatrix} \in M_2 \text{ and } t = \begin{bmatrix} 1 & 0 & 0 \\ 0 & p & q \\ 0 & r & 0 \end{bmatrix} \in S_4.$$

Suppose that $y_{32} \neq 0$.

- a) *If $p = 0$, then $t^{-1}mt \in U_2$ if and only if $y_{22} = y_{33} = 1$ and $y_{23} = 0$. In particular, we have*

$$\sum_{t \in S_4} \mu_2(t^{-1}mt) = \sum_{r,q \in F^\times} \psi_0(qr^{-1}y_{32} + ry_{13}).$$

- b) *If $p \neq 0$, then $t^{-1}mt \in U_2$ if and only if $y_{22} + y_{33} = 2$, $y_{23} = -\frac{(y_{22}-1)^2}{y_{32}}$ and $r = \left(\frac{y_{32}p}{y_{22}-1}\right)$. In particular, we have*

$$\sum_{t \in S_4} \mu_2(t^{-1}mt) = - \sum_{p \in F^\times} \psi_0(p(\gamma + y_{12})),$$

$$\text{where } \gamma = y_{32}y_{13}(y_{22} - 1)^{-1}.$$

Proof. Similar to Lemma 4.6. □

We record the character values of ρ_2 in the following table.

TABLE 7. Character of ρ_2

Type of m	m	$\chi_{\rho_2}(m)$	Type of m	m	$\chi_{\rho_2}(m)$
Type-1	$\left\{ \begin{bmatrix} 1 & 0 & y_{13} \\ 0 & 1 & y_{23} \\ 0 & 0 & 1 \end{bmatrix} \middle y_{23} \in F^\times \right\}$	$(1 - q)$	Type-6	$\left\{ \begin{bmatrix} 1 & y_{12} & y_{13} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \middle y_{12}, y_{13} \in F^\times \right\}$	$(1 - q)$
Type-2	$\left\{ \begin{bmatrix} 1 & y_{12} & y_{13} \\ 0 & 1 & y_{23} \\ 0 & 0 & 1 \end{bmatrix} \middle y_{12}, y_{23} \in F^\times \right\}$	1	Type-7	$\left\{ \begin{bmatrix} 1 & y_{12} & 0 \\ 0 & 1 & 0 \\ 0 & y_{32} & 1 \end{bmatrix} \middle y_{32} \in F^\times \right\}$	$(1 - q)$
Type-3	$\left\{ \begin{bmatrix} 1 & y_{12} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \middle y_{12} \in F^\times \right\}$	$(1 - q)$	Type-8	$\left\{ \begin{bmatrix} 1 & y_{12} & y_{13} \\ 0 & 1 & 0 \\ 0 & y_{32} & 1 \end{bmatrix} \middle y_{13}, y_{32} \in F^\times \right\}$	1
Type-4	$\left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right\}$	$(1 - q)(1 - q^2)$	Type-9	$\left\{ \begin{bmatrix} 1 & y_{12} & y_{13} \\ 0 & y_{22} & y_{23} \\ 0 & y_{32} & y_{33} \end{bmatrix} \middle \begin{array}{l} y_{32} \in F^\times, y_{12} = -y, \\ y_{23} = -y_{32}^{-1}(y_{22} - 1)^2 \end{array} \right\}$	$(1 - q)$
Type-5	$\left\{ \begin{bmatrix} 1 & 0 & y_{13} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \middle y_{13} \in F^\times \right\}$	$(1 - q)$	Type-10	$\left\{ \begin{bmatrix} 1 & y_{12} & y_{13} \\ 0 & y_{22} & y_{23} \\ 0 & y_{32} & y_{33} \end{bmatrix} \middle \begin{array}{l} y_{32} \in F^\times, y_{12} \neq -y, \\ y_{23} = -y_{32}^{-1}(y_{22} - 1)^2 \end{array} \right\}$	1

If $m \in M_2$ is not one of the types mentioned above, we have $\chi_{\rho_2}(m) = 0$.

For $1 \leq i, j \leq 10$, we let

$$T(i, j) = \{k = (m_1, m_2) \in H_A \mid m_1 \in \text{Type } -i, m_2 \in \text{Type } -j\}.$$

Theorem 4.10. Let $\rho = \theta|_{F^\times} \otimes \rho_1 \otimes \rho_2$. Let χ_ρ be the character of ρ . For $m = (a, m_1, m_2) \in Z \times M_1 \times M_2$, we have

$$\chi_\rho(m) = \theta(a)\chi_{\rho_1}(m_1)\chi_{\rho_2}(m_2)$$

where $(m_1, m_2) \in T(i, j)$, $i, j \in \{1, \dots, 10\}$. Otherwise, $\chi_\rho(m) = 0$.

Proof. We summarize the results from Table (6) and Table (7). \square

TABLE 8. Character of ρ

	Type-1	Type-2	Type-3	Type-4	Type-5	Type-6	Type-7	Type-8	Type-9	Type-10
Type-1	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$
Type-2	$\theta(a)(1-q)$	$\theta(a)$	$\theta(a)(1-q)$	$\theta(a)(1-q)(1-q^2)$	$\theta(a)(1-q)$	$\theta(a)(1-q)$	$\theta(a)(1-q)$	$\theta(a)$	$\theta(a)(1-q)$	$\theta(a)$
Type-3	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$
Type-4	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)(1-q^2)$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2(1-q^2)^2$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)(1-q^2)$
Type-5	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$
Type-6	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$
Type-7	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$
Type-8	$\theta(a)(1-q)$	$\theta(a)$	$\theta(a)(1-q)$	$\theta(a)(1-q)(1-q^2)$	$\theta(a)(1-q)$	$\theta(a)(1-q)$	$\theta(a)(1-q)$	$\theta(a)$	$\theta(a)(1-q)$	$\theta(a)$
Type-9	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2(1-q^2)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$	$\theta(a)(1-q)^2$	$\theta(a)(1-q)$
Type-10	$\theta(a)(1-q)$	$\theta(a)$	$\theta(a)(1-q)$	$\theta(a)(1-q)(1-q^2)$	$\theta(a)(1-q)$	$\theta(a)(1-q)$	$\theta(a)$	$\theta(a)(1-q)$	$\theta(a)(1-q)$	$\theta(a)$

4.2. Character calculation for π_{N,ψ_A} .

Lemma 4.11. Let $m = ah \in M_{\psi_A}$, where $a \in Z$ and $h \in H_A$. Then,

$$\Theta_{N,\psi_A}(m) = \theta(a)\Theta_{N,\psi_A}(h).$$

Proof. We have

$$\begin{aligned} \Theta_{N,\psi_A}(m) &= \Theta_{N,\psi_A}(ah) \\ &= \frac{1}{|N|} \sum_{n \in N} \Theta_\theta(ahn) \overline{\psi_A(n)} \\ &= \frac{1}{|N|} \sum_{n \in N} \mathrm{Tr}(\pi(ahn)) \overline{\psi_A(n)} \\ &= \frac{1}{|N|} \sum_{n \in N} \mathrm{Tr}(\pi(a)\pi(hn)) \overline{\psi_A(n)} \\ &= \omega_\pi(a) \frac{1}{|N|} \sum_{n \in N} \mathrm{Tr}(\pi(hn)) \overline{\psi_A(n)} \\ &= \omega_\pi(a) \Theta_{N,\psi_A}(h) \end{aligned}$$

where ω_π is the central character of π . Explicitly, we have

$$\Theta_\theta(a) = \mathrm{Tr}(\pi(a)) = \mathrm{Tr}(\omega_\pi(a)) = \omega_\pi(a) \dim(\pi).$$

Using Theorem 2.2, it is easy to see that

$$\Theta_\theta(a) = \theta(a) \dim(\pi).$$

Thus, we have $\omega_\pi(a) = \theta(a)$ and the result follows. \square

Lemma 4.12. Let $\tau = \begin{bmatrix} 0 & w_0 \\ w_0 & 0 \end{bmatrix}$. For $1 \leq i, j \leq 10$, we have

$$T(j, i) = \tau T(i, j)^\top \tau^{-1}.$$

Proof. Trivial. □

Theorem 4.13. Let $m' \in T(j, i)$. Then there exists $m \in T(i, j)$ such that

$$\Theta_{N, \psi_A}(m) = \Theta_{N, \psi_A}(m').$$

Proof. Let $m' = \begin{bmatrix} m'_1 & 0 \\ 0 & m'_2 \end{bmatrix} \in T(j, i)$. Since $T(j, i) = \tau T(i, j)^\top \tau^{-1}$, it follows that,

$$m' = \tau m^\top \tau^{-1}$$

for some $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(i, j)$. Thus we have $m'_1 = w_0 m_2^\top w_0^{-1}$ and $m'_2 = w_0 m_1^\top w_0^{-1}$. Since $m'_1 \in M_1$, clearly $\psi_A(X) = \psi_A((w_0 m_2^\top w_0)^{-1} X)$. We have

$$\begin{aligned} \Theta_{N, \psi_A}(m') &= \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \left[\begin{bmatrix} w_0 m_2^\top w_0^{-1} & X \\ 0 & w_0 m_1^\top w_0^{-1} \end{bmatrix} \overline{\psi_A(X)} \right] \\ &= \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \left(\begin{bmatrix} w_0 & 0 \\ 0 & w_0 \end{bmatrix} \begin{bmatrix} m_2^\top & w_0^{-1} X w_0 \\ 0 & m_1^\top \end{bmatrix} \begin{bmatrix} w_0 & 0 \\ 0 & w_0 \end{bmatrix} \right) \overline{\psi_A(X)} \\ &= \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \left(\begin{bmatrix} m_2^\top & w_0^{-1} X w_0 \\ 0 & m_1^\top \end{bmatrix} \right) \overline{\psi_A(X)}. \end{aligned}$$

On the other hand, using $\text{Tr}(A(w_0^{-1} X^\top w_0)) = \text{Tr}(AX)$ we have

$$\begin{aligned} \Theta_{N, \psi_A}(m) &= \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \left[\begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(X)} \right] \\ &= \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \left[\begin{bmatrix} m_1 & X^\top \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(X^\top)} \right] \\ &= \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \left[\begin{bmatrix} m_1 & w_0^{-1} X^\top w_0 \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(w_0^{-1} X^\top w_0)} \right] \\ &= \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \left[\begin{bmatrix} m_1 & w_0^{-1} X^\top w_0 \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(X)} \right]. \end{aligned}$$

Since

$$\begin{aligned} \text{Rank} \left(\begin{bmatrix} m_2^\top - 1 & w_0^{-1} X w_0 \\ 0 & m_1^\top - 1 \end{bmatrix} \right) &= \text{Rank} \left(\begin{bmatrix} m_2 - 1 & 0 \\ w_0^{-1} X^\top w_0 & m_1 - 1 \end{bmatrix} \right) \\ &= \text{Rank} \left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} m_2 - 1 & 0 \\ w_0^{-1} X^\top w_0 & m_1 - 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right) \\ &= \text{Rank} \left(\begin{bmatrix} m_1 - 1 & w_0^{-1} X^\top w_0 \\ 0 & m_2 - 1 \end{bmatrix} \right) \end{aligned}$$

we have,

$$\dim(\ker \begin{bmatrix} m_2^\top - 1 & w_0^{-1} X w_0 \\ 0 & m_1^\top - 1 \end{bmatrix}) = \dim(\ker \begin{bmatrix} m_1 - 1 & w_0^{-1} X^\top w_0 \\ 0 & m_2 - 1 \end{bmatrix}).$$

Hence,

$$\Theta_\theta \begin{bmatrix} m_2^\top & w_0^{-1} X w_0 \\ 0 & m_1^\top \end{bmatrix} = \Theta_\theta \begin{bmatrix} m_1 & w_0^{-1} X^\top w_0 \\ 0 & m_2 \end{bmatrix}$$

and the result follows. \square

Remark 4.14. We have used the fact that $\mathrm{Rank}(M) = \mathrm{Rank}(M^\top)$ and $\mathrm{Rank}(NMP) = \mathrm{Rank}(M)$ if N and P are invertible matrices.

Let $m = (m_1, m_2) \in M_1 \times M_2 = H_A$. Suppose also that m_1, m_2 are unipotent. To calculate $\Theta_{N, \psi_A}(m)$, we need to compute $\Theta_\theta(h)$, where $h = \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix}$. Using Theorem 2.2, it suffices to compute $\dim \mathrm{Ker}(h - 1)$. We note that the following proposition is valid even when H_A is a subgroup of $\mathrm{GL}(2n, F)$.

Proposition 4.15. Let $h = \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \in \mathrm{GL}(2n, F)$, where $(m_1, m_2) \in H_A$.

Suppose that m_1 and m_2 are also unipotent. Let $W' = \mathrm{Ker}(m_2 - 1)$. Then, we have

$$\begin{aligned} \dim \mathrm{Ker}(h - 1) &= \dim \mathrm{Ker}(m_1 - 1) + \dim \mathrm{Ker}(m_2 - 1) \\ &\quad - \dim(XW') + \dim\{XW' \cap \mathrm{Im}(m_1 - 1)\}. \end{aligned}$$

Proof. Let V be an n -dimensional vector space over F and let m_1, m_2, X be linear operators on V . Suppose that $\{e_1, \dots, e_m\}$ is a basis for $\mathrm{Ker}(m_1 - 1)$ and $\{f_1, \dots, f_k\}$ is a basis for $\mathrm{Ker}(m_2 - 1)$. Extending the basis of $\mathrm{Ker}(m_1 - 1)$ and $\mathrm{Ker}(m_2 - 1)$ we get ordered bases $\beta = \{e_1, \dots, e_n\}$ and $\beta' = \{f_1, \dots, f_n\}$ of V . Consider the ordered basis $\tilde{\beta} = \{(e_1, 0), \dots, (e_n, 0), (0, f_1), \dots, (0, f_n)\}$ of $V \oplus V$. We let h to be the linear operator on $V \oplus V$ defined as follows. For $1 \leq i, j \leq n$,

$$h((e_i, 0)) = (m_1, 0)(e_i, 0) = (m_1(e_i), 0)$$

and

$$h((0, f_j)) = (X, m_2)(0, f_j) = (X(f_j), m_2(f_j)).$$

Then,

$$[h]_{\tilde{\beta}} = \begin{bmatrix} [m_1]_\beta & [X]_{\beta'}^\beta \\ 0 & [m_2]_{\beta'} \end{bmatrix}$$

where

$$[X]_{\beta'}^\beta = [X_1 \ X_2 \ \cdots \ X_n].$$

Let

$$W_1 = \mathrm{Span}\{(m_1 - 1, 0)(e_{m+1}, 0), \dots, (m_1 - 1, 0)(e_n, 0)\} = \mathrm{Im}(m_1 - 1),$$

$$W_2 = \mathrm{Span}\{(X, m_2 - 1)(0, f_1), \dots, (X, m_2 - 1)(0, f_k)\} = XW'$$

and

$$W_3 = \text{Span}\{(Xf_{k+1}, (m_2 - 1)f_{k+1}), \dots, (Xf_n, (m_2 - 1)f_n)\}.$$

Clearly,

$$\text{Im}(h - 1) = W_1 + W_2 + W_3.$$

It is easy to see that

$$W_2 \cap W_3 = \{0\} = W_1 \cap W_3.$$

Since $\dim(\text{Ker}(m_2 - 1)) = k$, we have that

$$\dim(W_3) = \dim(\text{Im}(m_2 - 1)).$$

Therefore,

$$\begin{aligned} \dim(\text{Im}(h - 1)) &= \dim(\text{Im}(m_1 - 1)) + \dim(\text{Im}(m_2 - 1)) \\ &\quad + \dim(XW') - \dim(XW' \cap \text{Im}(m_1 - 1)), \end{aligned}$$

hence the result. \square

Remark 4.16. Let h be as in Proposition 4.15. We note that

$$XW' = \text{Span}\{X_1, X_2, \dots, X_k\}.$$

We will continue to use this in our character calculations at several instances to follow.

Let $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in H_A$, where $m_1 \in M_1$, $m_2 \in M_2$. Throughout we write $W' = \text{Ker}(m_2 - 1)$. For $X \in M(3, F)$, we let $h = \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix}$. For $\beta \in F$, we define

$$S(\beta) = \{X \in M(3, F) \mid \text{Tr}(Am_1^{-1}X) = \text{Tr}(AX) = \beta\}.$$

Let

$$E = \bigcup_{\substack{i \leq j \\ i, j \in \{1, 2, 4\}}} T(i, j).$$

We call E to be the fundamental set. To determine $\Theta_{N, \psi_A}(m)$ for $m \in T(i, j)$, it is enough to compute $\Theta_{N, \psi_A}(m)$ for $m \in E$.

Theorem 4.17. *Let $m \in T(1, 1)$. Then, we have*

$$\Theta_{N, \psi_A}(m) = (1 - q)^2.$$

Proof. We have

$$\Theta_{N, \psi_A}(m) = \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \left[\begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \right] \overline{\psi_A(m_1^{-1}X)}.$$

Note that $\dim \text{Ker}(m_1 - 1) = 2$, $\text{Im}(m_1 - 1) = \text{Span}\{e_1\}$ and $\text{Ker}(m_2 - 1) = \text{Span}\{e_1, e_2\}$. To calculate the character value, we write

$$\Theta_{N, \psi_A}(m) = \frac{1}{q^9} (A_1 + A_2)$$

where

$$A_1 = \sum_{X \in S(0)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}$$

and

$$A_2 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}.$$

For simplicity, we let $t = \dim(\mathrm{Ker}(h - 1))$. To compute A_1 , we find a partition of $S(0)$ according to the value of t and compute the respective cardinalities. We record the details in the table below.

TABLE 9. A_1

	Partition of $S(0)$	$\dim(XW')$	$\dim(XW' \cap \mathrm{Im}(m_1 - 1))$	$t = \dim(\mathrm{Ker}(h - 1))$	Cardinality
1)(a)	$\left\{ \begin{bmatrix} 0 & 0 & e \\ 0 & 0 & f \\ 0 & 0 & l \end{bmatrix} \right\}$	0	0	4	q^3
1)(b)	$\left\{ \begin{bmatrix} 0 & 0 & e \\ 0 & 0 & f \\ 0 & k & l \end{bmatrix} \middle k \in F^\times \right\}$	1	0	3	$(q - 1)q^3$
2)(a)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ 0 & 0 & l \end{bmatrix} \middle ad - bc \neq 0 \right\}$	2	1	3	$(q^2 - 1)(q - 1)q^4$
2)(b)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ 0 & k & l \end{bmatrix} \middle c, k \in F^\times, ad - bc \neq 0 \right\}$	2	0	2	$(q - 1)^3 q^5$
2)(c)	$\left\{ \begin{bmatrix} a & b & e \\ 0 & d & f \\ 0 & k & l \end{bmatrix} \middle k \in F^\times, ad \neq 0 \right\}$	2	1	3	$(q - 1)^3 q^4$
3)(a)	$\left\{ \begin{bmatrix} a & ya & e \\ c & yc & f \\ 0 & 0 & l \end{bmatrix} \middle c \in F^\times, y \in F \right\}$	1	0	3	$(q - 1)q^3$
3)(b)	$\left\{ \begin{bmatrix} a & ya & e \\ 0 & 0 & f \\ 0 & 0 & l \end{bmatrix} \middle a \in F^\times, y \in F \right\}$	1	1	4	$(q - 1)q^4$
3)(c)	$\left\{ \begin{bmatrix} a & ya & e \\ c & yc & f \\ 0 & k & l \end{bmatrix} \middle c, k \in F^\times, y \in F \right\}$	2	0	2	$(q - 1)^2 q^5$
3)(d)	$\left\{ \begin{bmatrix} a & ya & e \\ 0 & 0 & f \\ 0 & k & l \end{bmatrix} \middle k \in F^\times, y \in F \right\}$	2	1	3	$(q - 1)^2 q^4$
4)(a)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ 0 & 0 & l \end{bmatrix} \middle d \in F^\times \right\}$	1	0	3	$(q - 1)q^4$
4)(b)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & 0 & f \\ 0 & 0 & l \end{bmatrix} \middle b \in F^\times \right\}$	1	1	4	$(q - 1)q^3$
4)(c)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ 0 & k & l \end{bmatrix} \middle k \in F^\times \right\}$	1	0	3	$(q^2 - 1)(q - 1)q^3$

Hence,

$$A_1 = K_1 + K_2 + K_3$$

where

- a) $K_1 = \sum_{\substack{X \in S(0) \\ t=4}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^5(1-q)(1-q^2)(1-q^3).$
- b) $K_2 = \sum_{\substack{X \in S(0) \\ t=3}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^5(2q+1)(q-1)^2(q^2-1).$
- c) $K_3 = \sum_{\substack{X \in S(0) \\ t=2}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^6(q-1)^3.$

It follows that

$$A_1 = \sum_{X \in S(0)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^8(q-1)^3. \quad (4.1)$$

Proceeding in a similar way, we find a partition of $S(\beta)$ to compute A_2 . We record the details in the table 10.

Hence, $A_2 = K_4 + K_5$, where

- a) $K_4 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^7(q-1)(q^2-1).$
- b) $K_5 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^7(q-1)^2.$

It follows that

$$A_2 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^8(q-1)^2. \quad (4.2)$$

From (4.1) and (4.2), it follows that

$$\Theta_{N,\psi_A}(m) = (1-q)^2.$$

TABLE 10. A_2

	Partition of $S(\beta)$	$\dim(XW')$	$\dim(XW' \cap \mathrm{Im}(m_1 - 1))$	$t = \dim(\mathrm{Ker}(h - 1))$	Cardinality
1)(a)	$\left\{ \begin{bmatrix} 0 & 0 & e \\ 0 & 0 & f \\ \beta & 0 & l \end{bmatrix} \right\}$	1	0	3	q^3
1)(b)	$\left\{ \begin{bmatrix} 0 & 0 & e \\ 0 & 0 & f \\ \beta & k & l \end{bmatrix} \middle k \in F^\times \right\}$	1	0	3	$(q - 1)q^3$
2)(a)	$\left\{ \begin{bmatrix} a & b & c \\ 0 & d & f \\ \beta & 0 & l \end{bmatrix} \middle d \in F^\times, ad - bc \neq 0 \right\}$	2	0	2	$(q - 1)^2 q^4$
2)(b)	$\left\{ \begin{bmatrix} a & b & e \\ c & 0 & f \\ \beta & 0 & l \end{bmatrix} \middle bc \neq 0 \right\}$	2	1	3	$(q - 1)^2 q^4$
2)(c)	$\left\{ \begin{bmatrix} a & b & e \\ 0 & d & f \\ \beta & k & l \end{bmatrix} \middle k \in F^\times, ad \neq 0 \right\}$	2	0	2	$(q - 1)^2(q^2 - q)q^3$
2)(d)	$\left\{ \begin{bmatrix} a & b & e \\ c & 0 & f \\ \beta & k & l \end{bmatrix} \middle k \in F^\times, bc \neq 0 \right\}$	2	0	2	$(q - 1)^2(q^2 - q)q^3$
2)(d)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ \beta & k & l \end{bmatrix} \middle c, d, k \in F^\times, ad - bc \neq 0, d = \beta^{-1}ck \right\}$	2	1	3	$(q - 1)^2(q^2 - q)q^3$
2)(e)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ \beta & k & l \end{bmatrix} \middle c, d, k \in F^\times, ad - bc \neq 0, d \neq \beta^{-1}ck \right\}$	2	0	2	$(q - 1)^3(q - 2)q^4$
3)(a)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ c & \gamma c & f \\ \beta & 0 & l \end{bmatrix} \middle c, \gamma \in F^\times \right\}$	2	0	2	$(q - 1)^2 q^4$
3)(b)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ 0 & 0 & f \\ \beta & 0 & l \end{bmatrix} \middle a, \gamma \in F^\times \right\}$	2	1	3	$(q - 1)^2 q^3$
3)(c)	$\left\{ \begin{bmatrix} a & 0 & e \\ c & 0 & f \\ \beta & 0 & l \end{bmatrix} \middle (a, c) \neq 0 \right\}$	1	0	3	$(q^2 - 1)q^3$
3)(d)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ c & \gamma c & f \\ \beta & k & l \end{bmatrix} \middle \gamma \in F^\times, k = \gamma\beta \right\}$	1	0	3	$(q^2 - 1)(q - 1)q^3$
3)(e)	$\left\{ \begin{bmatrix} a & 0 & e \\ c & 0 & f \\ \beta & k & l \end{bmatrix} \middle k, c \in F^\times \right\}$	2	0	2	$(q - 1)^2 q^4$
3)(f)	$\left\{ \begin{bmatrix} a & 0 & e \\ 0 & 0 & f \\ \beta & k & l \end{bmatrix} \middle k \in F^\times \right\}$	2	1	3	$(q - 1)^2 q^3$
3)(g)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ c & \gamma c & f \\ \beta & k & l \end{bmatrix} \middle c, k, \gamma \in F^\times, k \neq \gamma\beta \right\}$	2	0	2	$(q - 1)^2(q - 2)q^4$
3)(h)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ 0 & 0 & f \\ \beta & k & l \end{bmatrix} \middle a, k, \gamma \in F^\times, k \neq \gamma\beta \right\}$	2	1	3	$(q - 1)^2(q - 2)q^3$
4)(a)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ \beta & k & l \end{bmatrix} \middle d \in F^\times \right\}$	2	0	2	$(q - 1)q^5$
4)(b)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & 0 & f \\ \beta & k & l \end{bmatrix} \middle b \in F^\times \right\}$	2	1	3	$(q - 1)q^4$

□

Theorem 4.18. *Let $m \in T(1, 2)$. Then, we have*

$$\Theta_{N, \psi_A}(m) = (1 - q).$$

Proof. We have

$$\Theta_{N, \psi_A}(m) = \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}.$$

Note that $\dim \text{Ker}(m_1 - 1) = 2$, $\text{Im}(m_1 - 1) = \text{Span}\{e_1\}$ and $\text{Ker}(m_2 - 1) = \text{Span}\{e_1\}$. To calculate the character value, we write

$$\Theta_{N, \psi_A}(m) = \frac{1}{q^9} (B_1 + B_2)$$

where

$$B_1 = \sum_{X \in S(0)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}$$

and

$$B_2 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}.$$

For simplicity, we let $t = \dim(\text{Ker}(h - 1))$. To compute B_1 , we find a partition of $S(0)$ according to the value of t and compute the respective cardinalities. We record the details in the table below.

TABLE 11. B_1

	Partition of $S(0)$	$\dim(XW')$	$\dim(XW' \cap \text{Im}(m_1 - 1))$	$t = \dim(\text{Ker}(h - 1))$	Cardinality
1)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ 0 & k & l \end{bmatrix} \right\}$	0	0	3	q^6
2)(a)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ 0 & k & l \end{bmatrix} \middle c \in F^\times \right\}$	1	0	2	$(q - 1)q^7$
2)(b)	$\left\{ \begin{bmatrix} a & b & e \\ 0 & d & f \\ 0 & k & l \end{bmatrix} \middle a \in F^\times \right\}$	1	1	3	$(q - 1)q^6$

Hence,

$$B_1 = K_1 + K_2$$

where

$$\text{a) } K_1 = \sum_{\substack{X \in S(0) \\ t=3}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^7(1 - q)(1 - q^2).$$

$$\text{b) } K_2 = \sum_{\substack{X \in S(0) \\ t=2}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^7(1-q)^2.$$

It follows that

$$B_1 = \sum_{X \in S(0)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^8(q-1)^2. \quad (4.3)$$

Proceeding in a similar way, we find a partition of $S(\beta)$ to compute B_2 . We record the details in the table below.

TABLE 12. B_2

	Partition of $S(\beta)$	$\dim(XW')$	$\dim(XW' \cap \mathrm{Im}(m_1 - 1))$	$t = \dim(\mathrm{Ker}(h - 1))$	Cardinality
1)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ \beta & k & l \end{bmatrix} \right\}$	1	0	2	q^8

Hence,

$$B_2 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^8(1-q). \quad (4.4)$$

From (4.3) and (4.4), it follows that

$$\Theta_{N, \psi_A}(m) = (1-q).$$

□

Theorem 4.19. Let $m \in T(4, 1)$. Then, we have

$$\Theta_{N, \psi_A}(m) = (1-q)^2(1-q^2).$$

Proof. We have

$$\Theta_{N, \psi_A}(m) = \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}.$$

Note that $\dim \mathrm{Ker}(m_1 - 1) = 3$, $\mathrm{Im}(m_1 - 1) = \{0\}$ and $\mathrm{Ker}(m_2 - 1) = \mathrm{Span}\{e_1, e_2\}$. To calculate the character value, we write

$$\Theta_{N, \psi_A}(m) = \frac{1}{q^9}(C_1 + C_2)$$

where we have

$$C_1 = \sum_{X \in S(0)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}$$

and

$$C_2 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}.$$

For simplicity, we let $t = \dim(\text{Ker}(h - 1))$. To compute C_1 , we find a partition of $S(0)$ according to the value of t and compute the respective cardinalities. We record the details in the following table.

TABLE 13. C_1

	Partition of $S(0)$	$\dim(XW')$	$\dim(XW' \cap \text{Im}(m_1 - 1))$	$t = \dim(\text{Ker}(h - 1))$	Cardinality
1)(a)	$\left\{ \begin{bmatrix} 0 & 0 & e \\ 0 & 0 & f \\ 0 & 0 & t \end{bmatrix} \right\}$	0	0	5	q^3
1)(b)	$\left\{ \begin{bmatrix} 0 & 0 & e \\ 0 & 0 & f \\ 0 & k & l \end{bmatrix} \middle k \in F^\times \right\}$	1	0	4	$(q-1)q^3$
2)(a)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ 0 & 0 & l \end{bmatrix} \middle ad - bc \neq 0 \right\}$	2	0	3	$(q^2-1)(q^2-q)q^3$
2)(b)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ 0 & k & l \end{bmatrix} \middle k \in F^\times, ad - bc \neq 0 \right\}$	2	0	3	$(q^2-1)(q-1)^2q^4$
3)(a)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ c & \gamma c & f \\ 0 & 0 & l \end{bmatrix} \middle \gamma \in F \right\}$	1	0	4	$(q^2-1)q^4$
3)(b)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ c & \gamma c & f \\ 0 & k & l \end{bmatrix} \middle k \in F^\times, \gamma \in F \right\}$	2	0	3	$(q^2-1)(q-1)q^4$
4)(a)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ 0 & 0 & l \end{bmatrix} \right\}$	1	0	4	$(q^2-1)q^3$
4)(b)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ 0 & k & l \end{bmatrix} \middle k \in F^\times \right\}$	1	0	4	$(q^2-1)(q-1)q^3$

Hence,

$$C_1 = K_1 + K_2 + K_3$$

where

$$\text{a)} K_1 = \sum_{\substack{X \in S(0) \\ t=5}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^3(1-q)(1-q^2)(1-q^3)(1-q^4).$$

$$\text{b)} K_2 = \sum_{\substack{X \in S(0) \\ t=4}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^3(1-q)^2(1-q^2)(1-q^3)(2q^2+2q+1).$$

$$\text{c) } K_3 = \sum_{\substack{X \in S(0) \\ t=3}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^4(q^2-1)^2(1-q)(1-q^2).$$

It follows that

$$C_1 = \sum_{X \in S(0)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^8(1-q)^3(1-q^2). \quad (4.5)$$

Proceeding in a similar way, we find a partition of $S(\beta)$ to compute C_2 . We record the details in the table below.

TABLE 14. C_2

	Partition of $S(\beta)$	$\dim(XW')$	$\dim(XW' \cap \mathrm{Im}(m_1 - 1))$	$t = \dim(\mathrm{Ker}(h - 1))$	Cardinality
1)(a)	$\left\{ \begin{bmatrix} 0 & 0 & e \\ 0 & 0 & f \\ \beta & 0 & l \end{bmatrix} \right\}$	1	0	4	q^3
1)(b)	$\left\{ \begin{bmatrix} 0 & 0 & e \\ 0 & 0 & f \\ \beta & k & l \end{bmatrix} \middle k \in F^\times \right\}$	1	0	4	$(q-1)q^3$
2)(a)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ \beta & 0 & l \end{bmatrix} \middle ad - bc \neq 0 \right\}$	2	0	3	$(q^2-1)(q^2-q)q^3$
2)(b)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ \beta & k & l \end{bmatrix} \middle k \in F^\times, ad - bc \neq 0 \right\}$	2	0	3	$(q^2-1)(q-1)^2q^4$
3)(a)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ c & \gamma c & f \\ \beta & 0 & l \end{bmatrix} \middle \gamma \in F^\times \right\}$	2	0	3	$(q^2-1)(q-1)q^3$
3)(b)	$\left\{ \begin{bmatrix} a & 0 & e \\ c & 0 & f \\ \beta & 0 & l \end{bmatrix} \middle (a, c) \neq 0 \right\}$	1	0	4	$(q^2-1)q^3$
3)(c)	$\left\{ \begin{bmatrix} a & 0 & e \\ c & 0 & f \\ \beta & k & l \end{bmatrix} \middle (a, c) \neq 0, k \in F^\times \right\}$	2	0	3	$(q^2-1)(q-1)q^3$
3)(d)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ c & \gamma c & f \\ \beta & k & l \end{bmatrix} \middle \gamma \in F^\times, k = \gamma\beta \right\}$	1	0	4	$(q^2-1)(q-1)q^3$
3)(e)	$\left\{ \begin{bmatrix} a & \gamma a & e \\ c & \gamma c & f \\ \beta & k & l \end{bmatrix} \middle k, \gamma \in F^\times, k \neq \gamma\beta \right\}$	2	0	3	$(q^2-1)(q-1)(q-2)q^3$
4)(a)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ \beta & 0 & l \end{bmatrix} \middle (b, d) \neq 0 \right\}$	2	0	3	$(q^2-1)q^3$
4)(b)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ \beta & k & l \end{bmatrix} \middle (b, d) \neq 0, k \in F^\times \right\}$	2	0	3	$(q^2-1)(q-1)q^3$

We have

$$C_2 = K_4 + K_5$$

where

- a) $K_4 = \sum_{\beta \in F^\times} \sum_{\substack{X \in S(\beta) \\ t=4}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^6(1-q)(1-q^2)(1-q^3).$
- b) $K_5 = \sum_{\beta \in F^\times} \sum_{\substack{X \in S(\beta) \\ t=3}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^6(1-q)(1-q^2)^2.$

It follows that

$$C_2 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^8(1-q)^2(1-q^2). \quad (4.6)$$

From (4.5) and (4.6), we have

$$\Theta_{N,\psi_A}(m) = (1-q^2)(1-q)^2.$$

□

Remark 4.20. Let $m \in T(1, 4)$. Since

$$\Theta_{N,\psi_A}(m) = \Theta_{N,\psi_A}(m')$$

for some $m' \in T(4, 1)$, it is enough to compute $\Theta_{N,\psi_A}(m')$ for $m' \in T(4, 1)$ to obtain the character value $\Theta_{N,\psi_A}(m)$.

Theorem 4.21. Let $m \in T(2, 2)$. Then, we have

$$\Theta_{N,\psi_A}(m) = 1.$$

Proof. We have

$$\Theta_{N,\psi_A}(m) = \frac{1}{|N|} \sum_{X \in M(3,F)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}.$$

Note that $\dim \text{Ker}(m_1 - 1) = 1$, $\text{Im}(m_1 - 1) = \text{Span}\{e_1, e_2\}$ and $\text{Ker}(m_2 - 1) = \text{Span}\{e_1\}$. To calculate the character value, we write

$$\Theta_{N,\psi_A}(m) = \frac{1}{q^9} (D_1 + D_2)$$

where

$$D_1 = \sum_{X \in S(0)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}$$

and

$$D_2 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}.$$

Let $t = \dim(\text{Ker}(h - 1))$. To compute D_1 we find a partition of $S(0)$ according to the value of t and compute the respective cardinalities. We record the details in the table below.

TABLE 15. D_1

	Partition of $S(0)$	$\dim(XW')$	$\dim(XW' \cap \mathrm{Im}(m_1 - 1))$	$t = \dim(\mathrm{Ker}(h - 1))$	Cardinality
1)(a)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ 0 & k & l \end{bmatrix} \mid (a, c) \neq 0 \right\}$	1	1	2	$(q^2 - 1)q^6$
1)(b)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ 0 & k & l \end{bmatrix} \right\}$	0	0	2	q^6

Hence,

$$D_1 = \sum_{\substack{X \in S(0) \\ t=2}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^8(1 - q). \quad (4.7)$$

Proceeding in a similar way, we find a partition of $S(\beta)$ to compute D_2 . We record the details in the following table.

TABLE 16. D_2

	Partition of $S(\beta)$	$\dim(XW')$	$\dim(XW' \cap \mathrm{Im}(m_1 - 1))$	$t = \dim(\mathrm{Ker}(h - 1))$	Cardinality
1)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ \beta & k & l \end{bmatrix} \right\}$	1	0	1	q^8

Thus, we have

$$D_2 = \sum_{\beta \in F^\times} \sum_{\substack{X \in S(\beta) \\ t=1}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^8. \quad (4.8)$$

From (4.7) and (4.8), it follows that

$$\Theta_{N, \psi_A}(m) = 1.$$

□

Theorem 4.22. *Let $m \in T(4, 2)$. Then, we have*

$$\Theta_{N, \psi_A}(m) = (1 - q)(1 - q^2).$$

Proof. We have

$$\Theta_{N, \psi_A}(m) = \frac{1}{|N|} \sum_{X \in M(3, F)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}.$$

Note that $\dim \text{Ker}(m_1 - 1) = 3$, $\text{Im}(m_1 - 1) = \{0\}$ and $\text{Ker}(m_2 - 1) = \text{Span}\{e_1\}$. To calculate the character value, we write

$$\Theta_{N,\psi_A}(m) = \frac{1}{q^9}(H_1 + H_2)$$

where

$$H_1 = \sum_{X \in S(0)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}$$

and

$$H_2 = \sum_{\beta \in F^\times} \sum_{X \in S(\beta)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)}.$$

Let $t = \dim(\text{Ker}(h - 1))$. To compute H_1 , we find a partition of $S(0)$ according to the value of t and compute the respective cardinalities. We record the details in the table below.

TABLE 17. H_1

	Partition of $S(0)$	$\dim(XW')$	$\dim(XW' \cap \text{Im}(m_1 - 1))$	$t = \dim(\text{Ker}(h - 1))$	Cardinality
1)(a)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ 0 & k & l \end{bmatrix} \mid (a, c) \neq 0 \right\}$	1	0	3	$(q^2 - 1)q^6$
1)(b)	$\left\{ \begin{bmatrix} 0 & b & e \\ 0 & d & f \\ 0 & k & l \end{bmatrix} \right\}$	0	0	4	q^6

Hence,

$$H_1 = K_1 + K_2$$

where

- a) $K_1 = \sum_{\substack{X \in S(0) \\ t=4}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^6(1-q)(1-q^2)(1-q^3).$
- b) $K_2 = \sum_{\substack{X \in S(0) \\ t=3}} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^6(1-q)(1-q^2)^2.$

It follows that

$$H_1 = \sum_{X \in S(0)} \Theta_\theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = -q^8(1-q)^2(1-q^2). \quad (4.9)$$

Proceeding in a similar way, we find a partition of $S(\beta)$ to compute H_2 . We record the details in the following table.

TABLE 18. H_2

	Partition of $S(\beta)$	$\dim(XW')$	$\dim(XW' \cap \mathrm{Im}(m_1 - 1))$	$t = \dim(\mathrm{Ker}(h - 1))$	Cardinality
1)	$\left\{ \begin{bmatrix} a & b & e \\ c & d & f \\ \beta & k & l \end{bmatrix} \right\}$	1	0	3	q^8

Thus, we have

$$H_2 = \sum_{\beta \in F^\times} \sum_{\substack{X \in S(\beta) \\ t=3}} \Theta \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix} \overline{\psi_A(m_1^{-1}X)} = q^8(1-q)(1-q^2). \quad (4.10)$$

From (4.9) and (4.10), it follows that

$$\Theta_{N, \psi_A}(m) = (1-q)(1-q^2).$$

□

Remark 4.23. Let $m \in T(2, 4)$. Since

$$\Theta_{N, \psi_A}(m) = \Theta_{N, \psi_A}(m')$$

for some $m' \in T(4, 2)$, it is enough to compute $\Theta_{N, \psi_A}(m')$ for $m' \in T(4, 2)$ to obtain $\Theta_{N, \psi_A}(m)$.

Theorem 4.24. Let $m \in T(4, 4)$. Then, we have

$$\Theta_{N, \psi_A}(m) = (1-q)^2(1-q^2)^2.$$

Proof. Since $m \in T(4, 4)$, we have $m = 1$, and the result follows from Theorem 3.3. To be precise, we have

$$\Theta_{N, \psi_A}(m) = \dim_{\mathbb{C}}(\pi_{N, \psi_A}) = (1-q)^2(1-q^2)^2.$$

□

Theorem 4.25. Let $1 \leq i \leq 10$. Suppose that $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(i, 5)$ and

$$m' = \begin{bmatrix} m_1 & 0 \\ 0 & m'_2 \end{bmatrix} \in T(i, 1). \text{ Then, we have}$$

$$\Theta_{N, \psi_A}(m) = \Theta_{N, \psi_A}(m').$$

Proof. Let $h = \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix}$ and $h' = \begin{bmatrix} m_1 & X \\ 0 & m'_2 \end{bmatrix}$ for $X \in M(3, F)$. Let

$$M_{m_1, m_2}^{d, \beta} = \{X \in S(\beta) \mid \dim(\mathrm{Ker}(h - 1)) = d\}$$

for $\beta \in F$. Clearly,

$$\mathrm{Ker}(m_2 - 1) = \mathrm{Ker}(m'_2 - 1).$$

Hence for any $X \in M(3, F)$,

$$X \mathrm{Ker}(m_2 - 1) = X \mathrm{Ker}(m'_2 - 1)$$

and

$$X \operatorname{Ker}(m_2 - 1) \cap \operatorname{Im}(m_1 - 1) = X \operatorname{Ker}(m'_2 - 1) \cap \operatorname{Im}(m_1 - 1).$$

In particular, for any $\beta \in F$, we have that

$$M_{m_1, m_2}^{d, \beta} = M_{m_1, m'_2}^{d, \beta}. \quad (4.11)$$

We have,

$$\Theta_{N, \psi_A}(m) = \frac{1}{q^9}(R_1 + R_2)$$

where

$$\begin{aligned} R_1 &= \sum_{d=1}^6 \sum_{X \in M_{m_1, m_2}^{d, 0}} \Theta_\theta(h) \overline{\psi_A(m_1^{-1}X)} \\ &= \sum_{d=1}^6 (-1)^{6-d}(1-q) \cdots (1-q^{d-1})(\#M_{m_1, m_2}^{d, 0}) \overline{\psi_0(0)} \end{aligned}$$

and

$$\begin{aligned} R_2 &= \sum_{\beta \in F^\times} \sum_{d=1}^6 \sum_{X \in M_{m_1, m_2}^{d, \beta}} \Theta_\theta(h) \overline{\psi_A(m_1^{-1}X)} \\ &= \sum_{\beta \in F^\times} \sum_{d=1}^6 (-1)^{6-d}(1-q) \cdots (1-q^{d-1})(\#M_{m_1, m_2}^{d, \beta}) \overline{\psi_0(\beta)} \end{aligned}$$

Thus,

$$\Theta_{N, \psi_A}(m) = \frac{1}{q^9} \sum_{d=1}^6 (-1)^{6-d}(1-q) \cdots (1-q^{d-1})(\#M_{m_1, m_2}^{d, 0} + \sum_{\beta \in F^\times} \#M_{m_1, m_2}^{d, \beta} \overline{\psi_0(\beta)}).$$

Similarly,

$$\Theta_{N, \psi_A}(m') = \frac{1}{q^9} \sum_{d=1}^6 (-1)^{6-d}(1-q) \cdots (1-q^{d-1})(\#M_{m_1, m'_2}^{d, 0} + \sum_{\beta \in F^\times} \#M_{m_1, m'_2}^{d, \beta} \overline{\psi_0(\beta)}).$$

Hence, it follows from equation (4.11) that

$$\Theta_{N, \psi_A}(m) = \Theta_{N, \psi_A}(m').$$

□

Proposition 4.26. *Let $1 \leq i \leq 10$ and $m \in T(i, 3)$. Then, there exists some $m' \in T(i, 5)$ such that*

$$\Theta_{N, \psi_A}(m) = \Theta_{N, \psi_A}(m').$$

Proof. Let $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(i, 3)$, and $w = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Let $m'_2 = wm_2w^{-1}$ and $m' = \begin{bmatrix} m_1 & 0 \\ 0 & m'_2 \end{bmatrix}$. Clearly, we have $m = \begin{bmatrix} 1 & 0 \\ 0 & w \end{bmatrix} m' \begin{bmatrix} 1 & 0 \\ 0 & w \end{bmatrix}^{-1}$ and $m' \in T(i, 5)$, hence the result. \square

Corollary 4.27. Let $1 \leq i \leq 10$ and $m \in T(i, 3)$. Then,

$$\Theta_{N, \psi_A}(m) = \Theta_{N, \psi_A}(m')$$

for some $m' \in T(i, 1)$.

Proof. Using Proposition 4.26 and Theorem 4.25, the result follows. \square

Proposition 4.28. Let $1 \leq i \leq 10$ and $m \in T(i, 7)$. Then, there exists some $m' \in T(i, 1)$ such that

$$\Theta_{N, \psi_A}(m) = \Theta_{N, \psi_A}(m').$$

Proof. Let $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(i, 7)$, and $w = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Let $m'_2 = wm_2w^{-1}$ and $m' = \begin{bmatrix} m_1 & 0 \\ 0 & m'_2 \end{bmatrix}$. Clearly, we have $m = \begin{bmatrix} 1 & 0 \\ 0 & w \end{bmatrix} m' \begin{bmatrix} 1 & 0 \\ 0 & w \end{bmatrix}^{-1}$ and $m' \in T(i, 1)$, hence the result. \square

Proposition 4.29. Let $1 \leq i \leq 10$ and $m \in T(i, 8)$. Then, there exists some $m' \in T(i, 2)$ such that

$$\Theta_{N, \psi_A}(m) = \Theta_{N, \psi_A}(m').$$

Proof. Let $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(i, 8)$, and $w = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. Let $m'_2 = wm_2w^{-1}$ and $m' = \begin{bmatrix} m_1 & 0 \\ 0 & m'_2 \end{bmatrix}$. Clearly, we have $m = \begin{bmatrix} 1 & 0 \\ 0 & w \end{bmatrix} m' \begin{bmatrix} 1 & 0 \\ 0 & w \end{bmatrix}^{-1}$ and $m' \in T(i, 2)$, hence the result. \square

Theorem 4.30. Let $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(1, 6)$ or $T(1, 9)$. Then, we have

$$\Theta_{N, \psi_A}(m) = (1 - q)^2.$$

Proof. Note that $\dim \mathrm{Ker}(m_1 - 1) = 2$, $\mathrm{Im}(m_1 - 1) = \mathrm{Span}\{e_1\}$, $\dim \mathrm{Ker}(m_2 - 1) = 2$. From Remark 4.16, it follows that whenever

$$h = \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix},$$

we have

$$XW' = \mathrm{Span}\{X_1, X_2\}.$$

Thus, proceeding in a similar fashion as in Theorem 4.17, we get that

$$\Theta_{N,\psi_A}(m) = (1 - q)^2.$$

□

Theorem 4.31. Let $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(2, 6)$ or $T(2, 9)$. Then, we have

$$\Theta_{N,\psi_A}(m) = (1 - q).$$

Proof. Note that $\dim \text{Ker}(m_1 - 1) = 1$, $\text{Im}(m_1 - 1) = \{e_1, e_2\}$, $\dim \text{Ker}(m_2 - 1) = 2$. From Remark 4.16, it follows that whenever

$$h = \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix},$$

we have

$$XW' = \text{Span}\{X_1, X_2\}.$$

Thus, the computations are similar to the case where $m \in T(2, 1)$. The result follows from Theorem 4.13 and Theorem 4.18. □

Theorem 4.32. Let $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(4, 6)$ or $T(4, 9)$. Then, we have

$$\Theta_{N,\psi_A}(m) = (1 - q)^2(1 - q^2).$$

Proof. Note that $\dim \text{Ker}(m_1 - 1) = 3$, $\text{Im}(m_1 - 1) = \{0\}$, $\dim \text{Ker}(m_2 - 1) = 2$. From Remark 4.16, it follows that whenever

$$h = \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix},$$

we have

$$XW' = \text{Span}\{X_1, X_2\}.$$

Thus, proceeding in a similar fashion as in Theorem 4.19, we get that

$$\Theta_{N,\psi_A}(m) = (1 - q)^2(1 - q^2).$$

□

Theorem 4.33. Let $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(6, 6)$ or $T(6, 9)$. Then, we have

$$\Theta_{N,\psi_A}(m) = (1 - q)^2.$$

Proof. Note that $\dim \text{Ker}(m_1 - 1) = 2$, $\dim \text{Ker}(m_2 - 1) = 2$, $\text{Im}(m_1 - 1) = \text{Span}\{\eta e_1 + e_2\}$ for $\eta \in F^\times$. From Remark 4.16, it follows that computing $\Theta_{N,\psi_A}(m)$ for $m \in T(6, 6)$ or $m \in T(6, 9)$ is the same as computing $\Theta_{N,\psi_A}(m')$ for $m' \in T(6, 1)$. Using Theorem 4.13 and Theorem 4.30, it follows that

$$\Theta_{N,\psi_A}(m) = (1 - q)^2.$$

□

Theorem 4.34. Let $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(9, 9)$. Then, we have

$$\Theta_{N, \psi_A}(m) = (1 - q)^2.$$

Proof. The proof is similar to Theorem 4.33. \square

Theorem 4.35. Let $1 \leq i \leq 10$. Suppose $m = \begin{bmatrix} m_1 & 0 \\ 0 & m_2 \end{bmatrix} \in T(i, 10)$ and $m' = \begin{bmatrix} m_1 & 0 \\ 0 & m'_2 \end{bmatrix} \in T(i, 2)$. Then, we have

$$\Theta_{N, \psi_A}(m) = \Theta_{N, \psi_A}(m').$$

Proof. Let $h = \begin{bmatrix} m_1 & X \\ 0 & m_2 \end{bmatrix}$ and $h' = \begin{bmatrix} m_1 & X \\ 0 & m'_2 \end{bmatrix}$ for $X \in M(3, F)$. Let

$$M_{m_1, m_2}^{d, \beta} = \{X \in S(\beta) \mid \dim(\mathrm{Ker}(h - 1)) = d\}$$

for $\beta \in F$. Clearly,

$$\mathrm{Ker}(m_2 - 1) = \mathrm{Ker}(m'_2 - 1).$$

Hence for any $X \in M(3, F)$,

$$X \mathrm{Ker}(m_2 - 1) = X \mathrm{Ker}(m'_2 - 1)$$

and

$$X \mathrm{Ker}(m_2 - 1) \cap \mathrm{Im}(m_1 - 1) = X \mathrm{Ker}(m'_2 - 1) \cap \mathrm{Im}(m_1 - 1).$$

In particular, for any $\beta \in F$, we have that

$$M_{m_1, m_2}^{d, \beta} = M_{m_1, m'_2}^{d, \beta}. \quad (4.12)$$

Therefore,

$$\Theta_{N, \psi_A}(m) = \frac{1}{q^9}(R_1 + R_2)$$

where

$$\begin{aligned} R_1 &= \sum_{d=1}^6 \sum_{X \in M_{m_1, m_2}^{d, 0}} \Theta_\theta(h) \overline{\psi_A(m_1^{-1}X)} \\ &= \sum_{d=1}^6 (-1)^{6-d}(1 - q) \cdots (1 - q^{d-1}) (\#M_{m_1, m_2}^{d, 0}) \overline{\psi_0(0)} \end{aligned}$$

and

$$\begin{aligned} R_2 &= \sum_{\beta \in F^\times} \sum_{d=1}^6 \sum_{X \in M_{m_1, m_2}^{d, \beta}} \Theta_\theta(h) \overline{\psi_A(m_1^{-1}X)} \\ &= \sum_{\beta \in F^\times} \sum_{d=1}^6 (-1)^{6-d}(1 - q) \cdots (1 - q^{d-1}) (\#M_{m_1, m_2}^{d, \beta}) \overline{\psi_0(\beta)} \end{aligned}$$

Thus,

$$\Theta_{N,\psi_A}(m) = \frac{1}{q^9} \sum_{d=1}^6 (-1)^{6-d} (1-q) \cdots (1-q^{d-1}) (\#M_{m_1, m_2}^{d,0} + \sum_{\beta \in F^\times} \#M_{m_1, m_2}^{d,\beta} \overline{\psi_0(\beta)}).$$

Similarly,

$$\Theta_{N,\psi_A}(m') = \frac{1}{q^9} \sum_{d=1}^6 (-1)^{6-d} (1-q) \cdots (1-q^{d-1}) (\#M_{m_1, m'_2}^{d,0} + \sum_{\beta \in F^\times} \#M_{m_1, m'_2}^{d,\beta} \overline{\psi_0(\beta)}).$$

Hence, it follows from equation (4.12) that

$$\Theta_{N,\psi_A}(m) = \Theta_{N,\psi_A}(m').$$

□

Remark 4.36. Let $1 \leq i \leq j \leq 10$. To determine $\Theta_{N,\psi_A}(m)$ for $m \in T(i, j)$, it is enough to compute $\Theta_{N,\psi_A}(m)$ for $m \in E$. We illustrate this by an example.

Suppose that we want to compute the character value $\Theta_{N,\psi_A}(m)$ for $m \in T(3, 7)$. From Proposition 4.28, it follows that $\Theta_{N,\psi_A}(m) = \Theta_{N,\psi_A}(k)$ for some $k \in T(3, 1)$. By Theorem 4.13, we have $\Theta_{N,\psi_A}(k) = \Theta_{N,\psi_A}(x)$ for some $x \in T(1, 3)$. Using Theorem 4.26, we have, $\Theta_{N,\psi_A}(x) = \Theta_{N,\psi_A}(y)$ for some $y \in T(1, 1)$. Thus, using Theorem 4.17 we have

$$\Theta_{N,\psi_A}(m) = (1-q)^2.$$

For clarity, we give an example to illustrate the chain of computations used to determine the character value of an element in $T(i, j)$ using the character value of an element in the fundamental set E .

$$(3, 7) \rightarrow (3, 1) \rightarrow (1, 3) \rightarrow (1, 1).$$

We summarize the sequence of computations used to calculate $\Theta_{N,\psi_A}(m)$ for $m \in T(i, j)$, $1 \leq i \leq j \leq 5$ in Table 19.

TABLE 19. Sequence of computations for $\Theta_{N,\psi_A}(m)$

	Type-1	Type-2	Type-3	Type-4	Type-5
Type-1	(1, 1)	(1, 2)	(1, 3) \rightarrow (1, 1)	(1, 4)	(1, 5) \rightarrow (1, 1)
Type-2	-	(2, 2)	(2, 3) \rightarrow (2, 1) \rightarrow (1, 2)	(2, 4)	(2, 5) \rightarrow (2, 1) \rightarrow (1, 2)
Type-3	-	-	(3, 3) \rightarrow (3, 1) \rightarrow (1, 3) \rightarrow (1, 1)	(4, 3) \rightarrow (4, 1) \rightarrow (1, 4)	(3, 5) \rightarrow (3, 1) \rightarrow (1, 3) \rightarrow (1, 1)
Type-4	-	-	-	(4, 4)	(4, 5) \rightarrow (4, 1) \rightarrow (1, 4)
Type-5	-	-	-	-	(5, 5) \rightarrow (5, 1) \rightarrow (1, 5) \rightarrow (1, 1)

Table 20 summarizes the sequence of computations used to calculate $\Theta_{N,\psi_A}(m)$ for $m \in T(i, j)$, $1 \leq i \leq 5, 6 \leq j \leq 10, i \leq j$.

TABLE 20. Sequence of computations for $\Theta_{N,\psi_A}(m)$

	Type-6	Type-7	Type-8	Type-9	Type-10
Type-1	$(1,6) \rightarrow (1,1)$	$(1,7) \rightarrow (1,1)$	$(1,8) \rightarrow (1,2)$	$(1,9) \rightarrow (1,1)$	$(1,10) \rightarrow (1,2)$
Type-2	$(2,6) \rightarrow (2,1) \rightarrow (1,2)$	$(2,7) \rightarrow (2,1) \rightarrow (1,2)$	$(2,8) \rightarrow (2,2)$	$(2,9) \rightarrow (2,1) \rightarrow (1,2)$	$(2,10) \rightarrow (2,2)$
Type-3	$(3,6) \rightarrow (6,3) \rightarrow (6,1) \rightarrow (1,6) \rightarrow (1,1)$	$(3,7) \rightarrow (3,1) \rightarrow (1,3) \rightarrow (1,1)$	$(3,8) \rightarrow (3,2) \rightarrow (2,3) \rightarrow (2,1) \rightarrow (1,2)$	$(3,9) \rightarrow (9,3) \rightarrow (9,1) \rightarrow (1,9) \rightarrow (1,1)$	$(3,10) \rightarrow (3,2) \rightarrow (2,3) \rightarrow (2,1) \rightarrow (1,2)$
Type-4	$(4,6) \rightarrow (4,1) \rightarrow (1,4)$	$(4,7) \rightarrow (4,1) \rightarrow (1,4)$	$(4,8) \rightarrow (4,2) \rightarrow (2,4)$	$(4,9) \rightarrow (4,1) \rightarrow (1,4)$	$(4,10) \rightarrow (4,2) \rightarrow (2,4)$
Type-5	$(5,6) \rightarrow (6,5) \rightarrow (6,1) \rightarrow (1,6) \rightarrow (1,1)$	$(5,7) \rightarrow (5,1) \rightarrow (1,5) \rightarrow (1,1)$	$(5,8) \rightarrow (5,2) \rightarrow (2,5) \rightarrow (2,1) \rightarrow (1,2)$	$(5,9) \rightarrow (9,5) \rightarrow (9,1) \rightarrow (1,9) \rightarrow (1,1)$	$(5,10) \rightarrow (5,2) \rightarrow (2,5) \rightarrow (2,1) \rightarrow (1,2)$

Table 21 summarizes the sequence of computations used to calculate $\Theta_{N,\psi_A}(m)$ for $m \in T(i, j)$, $6 \leq i \leq 10$, $6 \leq j \leq 10$, $i \leq j$.

TABLE 21. Sequence of computations for $\Theta_{N,\psi_A}(m)$

	Type-6	Type-7	Type-8	Type-9	Type-10
Type-6	$(6,6) \rightarrow (6,1) \rightarrow (1,6) \rightarrow (1,1)$	$(6,7) \rightarrow (6,1) \rightarrow (1,6) \rightarrow (1,1)$	$(6,8) \rightarrow (6,2) \rightarrow (2,6) \rightarrow (2,1) \rightarrow (1,2)$	$(6,9) \rightarrow (6,1) \rightarrow (1,6) \rightarrow (1,1)$	$(6,10) \rightarrow (6,2) \rightarrow (2,6) \rightarrow (2,1) \rightarrow (1,2)$
Type-7	-	$(7,7) \rightarrow (7,1) \rightarrow (1,7) \rightarrow (1,1)$	$(7,8) \rightarrow (7,2) \rightarrow (2,7) \rightarrow (2,1) \rightarrow (1,2)$	$(7,9) \rightarrow (9,7) \rightarrow (9,1) \rightarrow (1,9) \rightarrow (1,1)$	$(7,10) \rightarrow (7,2) \rightarrow (2,7) \rightarrow (2,1) \rightarrow (1,2)$
Type-8	-	-	$(8,8) \rightarrow (8,2) \rightarrow (2,8) \rightarrow (2,2)$	$(8,9) \rightarrow (9,8) \rightarrow (9,2) \rightarrow (2,9) \rightarrow (2,1) \rightarrow (1,2)$	$(8,10) \rightarrow (8,2) \rightarrow (2,8) \rightarrow (2,2)$
Type-9	-	-	-	$(9,9) \rightarrow (9,1) \rightarrow (1,9) \rightarrow (1,1)$	$(9,10) \rightarrow (9,2) \rightarrow (2,9) \rightarrow (2,1) \rightarrow (1,2)$
Type-10	-	-	-	-	$(10,10) \rightarrow (10,2) \rightarrow (2,10) \rightarrow (2,2)$

Acknowledgements

We thank Professor Dipendra Prasad for suggesting this problem and for some helpful discussions. Research of Kumar Balasubramanian is supported by the SERB grant: MTR/2019/000358.

References

- [1] BALASUBRAMANIAN, KUMAR; KHURANA, HIMANSI. A certain twisted Jacquet module of $\mathrm{GL}(4)$ over a finite field. *J. Pure Appl. Algebra* **226** (2022), no. 5, Paper No. 106932, 16 pp. MR4328653, Zbl 7455909, doi: 10.1016/j.jpaa.2021.106932. 874, 875, 877
- [2] BERNŠTEĬN, JOSEPH N.; ZELEVINSKIĬ, ANDREY V. Representations of the group $\mathrm{GL}(n, F)$, where F is a local non-Archimedean field. *Uspehi Mat. Nauk* **31** (1976), no. 3(189), 5–70. MR0425030, Zbl 0342.43017, doi: 10.1070/RM1976v03n03ABEH001532. 875
- [3] GELFAND, SERGEI I. Representations of the full linear group over a finite field, *Mat. Sb. (N. S.)* **83 (125)** (1970), 15–41. MR0272916, Zbl 0252.20034, doi: 10.1070/SM1970v012n01ABEH000907. 876

- [4] PRASAD, DIPENDRA. The space of degenerate Whittaker models for general linear groups over a finite field. *Internat. Math. Res. Notices* (2000), no. 11, 579–595. MR1763857, Zbl 0983.22014, doi: <https://doi.org/10.1155/S107379280000313>. 874, 876
- [5] SERRE, JEAN-PIERRE. Linear representations of finite groups. Graduate Texts in Mathematics, 42. Springer-Verlag, New York-Heidelberg, 1977. x+170 pp. ISBN:0-387-90190-6. MR0450380, Zbl 0355.20006, doi: 10.1007/978-1-4684-9458-7. 877

(Balasubramanian) DEPARTMENT OF MATHEMATICS, IISER BHOPAL, BHOPAL, MADHYA PRADESH 462066, INDIA
bkumar@iiserb.ac.in

(Khurana) DEPARTMENT OF MATHEMATICS, IISER BHOPAL, BHOPAL, MADHYA PRADESH 462066, INDIA
himanshi18@iiserb.ac.in

This paper is available via <http://nyjm.albany.edu/j/2023/29-34.html>.