

Power integral bases in a family of octic fields

István Gaál

ABSTRACT. Several recent results prove the monogeneity of some polynomials. In these cases the root of the polynomial generates a power integral basis in the number field generated by the root. A straightforward question is whether such a number field admits other generators of power integral bases? We have investigated this problem in some previous papers and here we extend this research to a family of octic polynomials, following a recent result of L. Jones [11].

CONTENTS

1. Introduction	619
2. The octic polynomial	620
3. Integral basis	622
4. A quartic relative Thue equations	623
5. Solving the quartic relative Thue equation	625
6. Reduction	626
7. Determining P and Q	627
8. Determining generators of power integral bases of K	628
9. Results of our calculations	629
10. Another solution	630
References	632

1. Introduction

A number field K of degree n with ring of integers \mathbb{Z}_K is called *monogenic* (cf. [2]) if there exists $\xi \in \mathbb{Z}_K$ such that $(1, \xi, \dots, \xi^{n-1})$ is an integral basis, called a power integral basis. We call ξ the generator of this power integral basis. $\alpha, \beta \in \mathbb{Z}_K$ are called *equivalent*, if $\alpha + \beta \in \mathbb{Z}$ or $\alpha - \beta \in \mathbb{Z}$. Obviously, α generates a power integral basis in K if and only if any β , equivalent to α ,

Received January 23, 2025.

2010 *Mathematics Subject Classification.* 11Y50, 11R04, 11D25.

Key words and phrases. monogeneity; power integral basis; octic fields; relative quartic fields; relative Thue equations.

does. As it is known, any algebraic number field admits up to equivalence only finitely many generators of power integral bases.

A monic irreducible polynomial $f(x) \in \mathbb{Z}[x]$ is called *monogenic*, if a root ξ of $f(x)$ generates a power integral basis in $K = \mathbb{Q}(\xi)$. If $f(x)$ is monogenic, then K is monogenic, but the converse is not true.

For $\alpha \in \mathbb{Z}_K$ (generating K over \mathbb{Q}) we call the module index

$$I(\alpha) = (\mathbb{Z}_K : \mathbb{Z}[\alpha])$$

the *index* of α . α generates a power integral basis in K if and only if $I(\alpha) = 1$. If $\alpha^{(i)}$ ($1 \leq i \leq n$) are the conjugates of α in K of degree n with absolute discriminant D_K , then

$$I(\alpha) = \frac{1}{\sqrt{|D_K|}} \prod_{1 \leq i < j \leq n} |\alpha^{(i)} - \alpha^{(j)}|.$$

For more details concerning monogeneity and power integral bases cf. [2].

In some recent papers we investigated number fields generated by a root of a monogenic polynomial and made calculations to figure out, whether these fields admit any additional generators of power integral bases. We refer to [3] for some sextic trinomials, [4] for pure sextic fields, [5] for pure octic fields, [6] for certain quartic trinomials and [7] for some quartic polynomials with given Galois groups.

L. Jones [10] (see also [11]) gave conditions for the monogeneity of certain even octic polynomials of type $x^8 + ax^6 + bx^4 + ax^2 + 1$. In the present paper we extend our calculations to this type of polynomials. Among others we prove the existence of a non-trivial generator of power integral basis.

The octic field, generated by a root of the above polynomial is a quadratic extension of a quartic field. It is an interesting point of our arguments, that this octic field can also be considered as a quartic extension of a quadratic field, which makes it much easier to deal with.

All tools used in our calculations are optimized to this special case, in order to make our calculations more efficient.

2. The octic polynomial

Let

$$f(x) = x^8 + ax^6 + bx^4 + ax^2 + 1 \quad (1)$$

with $a, b \in \mathbb{Z}$. Set $W_1 = b + 2 - 2a$, $W_2 = b + 2 + 2a$, $W_3 = a^2 - 4b + 8$. L. Jones [11] proved:

Theorem 2.1. *If $W_1 W_2 W_3$ is square free and*

$$(a \pmod 4, b \pmod 4) \in \{(1, 3), (3, 1), (3, 3)\} \quad (2)$$

then the polynomial $f(x)$ in (1) is monogenic.

Assume $f(x)$ is monogenic, not necessarily satisfying (2). We wonder how many generators of power integral bases the number field K has, generated by a root α of $f(x)$.

Set

$$g(x) = x^4 + ax^3 + bx^2 + ax + 1. \quad (3)$$

Obviously, $\beta = \alpha^2$ is a root of $g(x)$. Therefore, the octic number field $K = \mathbb{Q}(\alpha)$ is a quadratic extension of the quartic field $L = \mathbb{Q}(\beta)$:

$$\mathbb{Q} \subset L = \mathbb{Q}(\beta) \subset K = \mathbb{Q}(\alpha).$$

Unfortunately, there exist no feasible algorithms for solving index form equations, that is, for determining generators of power integral bases, only for a restricted class of number fields. Apart from low degree fields, like cubic and quartic fields, there exist such algorithms only for some higher degree fields with special structure. These are, e.g. sextic fields with a quadratic subfield and octic fields with a quadratic subfield (cf. [2]). Above we have an octic field with a quartic subfield, but using the reciprocal structure of $f(x)$ and $g(x)$ we can help this problem.

If

$$\beta^4 + a\beta^3 + b\beta^2 + a\beta + 1 = 0,$$

then

$$\beta^2 + a\beta + b + \frac{a}{\beta} + \frac{1}{\beta^2} = 0,$$

hence

$$\begin{aligned} \left(\beta^2 + \frac{1}{\beta^2}\right) + a\left(\beta + \frac{1}{\beta}\right) + b &= 0, \\ \left(\beta + \frac{1}{\beta}\right)^2 + a\left(\beta + \frac{1}{\beta}\right) + b - 2 &= 0. \end{aligned}$$

This yields, that

$$\delta = \beta + \frac{1}{\beta} \quad (4)$$

satisfies the quadratic equation

$$\delta^2 + a\delta + (b - 2) = 0. \quad (5)$$

Consequently, the number field $M = \mathbb{Q}(\delta)$ is a quadratic subfield of K :

$$\mathbb{Q} \subset M = \mathbb{Q}(\delta) \subset K = \mathbb{Q}(\alpha).$$

By (4) we have

$$\beta^2 - \delta\beta + 1 = 0,$$

therefore

$$\alpha^4 - \delta\alpha^2 + 1 = 0.$$

This means that

$$h(x) = x^4 - \delta x^2 + 1 \quad (6)$$

is the relative defining polynomial of α over M , and this is what we need for our procedure.

Note that in some previous papers (cf. [9]) we have developed an algorithm for the complete resolution of index form equations in octic fields with a quadratic subfield. This takes quite a long CPU time, since one has to solve a unit equation in the octic field. Also, it can take long to calculate the fundamental units of that field, which is a necessary input data for the calculations.

Therefore, if we would like to have an overall picture about the generators of power integral bases of our octic fields, we have to restrict ourselves to the calculation of the so called "small solutions", that means we calculate all generators of power integral bases having coefficients, say $\leq 10^{200}$ in absolute value in a given integral basis. Since the generators of power integral bases usually have very small coefficients, such an algorithm determines all generators of power integral bases with a very high probability. Moreover, it certainly indicates, if a number field, generated by a root of a monogenic polynomial, has also other generators of power integral bases, in addition to the root of the polynomial.

As we shall see in the following, a crucial point in this algorithm is the resolution of a relative quartic Thue equation over the quadratic subfield. The fast algorithm [1] for determining "small" solutions of quartic relative Thue equations over quadratic fields is only efficient if the quadratic subfield is complex. Therefore in our calculations we assume

$$a^2 - 4b + 8 < 0, \quad (7)$$

which guarantees by (5), that M is a complex quadratic subfield.

On the other hand, we shall not restrict ourselves to those monogenic polynomials $f(x)$, satisfying all conditions of Theorem 2.1. We shall run the parameters a, b in certain regions and consider all irreducible polynomials $f(x)$ that are monogenic. The only condition we keep is that $W_3 = a^2 - 4b + 8$ is square-free, in order to fix the basis element of M and to make our arguments simpler. Note, that we made calculations also for non-squarefree W_3 , and had completely the same experiences, including also the non-trivial generator of power integral bases (cf. Theorem 10.1).

3. Integral basis

We have $M = \mathbb{Q}(\delta)$, with

$$\delta = \frac{-a + \sqrt{a^2 - 4b + 8}}{2}. \quad (8)$$

According to the above arguments, we assume $W_3 = a^2 - 4b + 8 < 0$ is square-free. To keep usual notation we set $m = W_3$. This number can only be square-free if $a \equiv \pm 1 \pmod{4}$, whence $m \equiv 1 \pmod{4}$, therefore the integral basis of the complex quadratic field $M = \mathbb{Q}(\sqrt{m})$ is $(1, \omega)$, where

$$\omega = \frac{1 + \sqrt{m}}{2}.$$

We shall make calculations for monogenic polynomials $f(x)$. In this case a root α of $f(x)$ generates a power integral basis in $K = \mathbb{Q}(\alpha)$.

We shall use the following statements of [8] which are certainly well known:

Theorem 3.1.

A. If K is monogenic, then K is also relative monogenic over the subfield M .

B. All generators of power integral bases of K are of the form

$$\gamma = X_0 + \varepsilon\gamma_0,$$

where $X_0 \in \mathbb{Z}_M$, ε is a unit in M and γ_0 generates a relative power integral basis of K over M .

If

$$(1, \alpha, \alpha^2, \alpha^3, \alpha^4, \alpha^5, \alpha^6, \alpha^7)$$

is an integral basis of K , then by the first part of the theorem

$$(1, \alpha, \alpha^2, \alpha^3)$$

is a relative integral basis of K over M , that is any $\gamma \in \mathbb{Z}_K$ can be written in the form

$$\gamma = C + X\alpha + Y\alpha^2 + Z\alpha^3, \quad (9)$$

where

$$C = c_1 + \omega c_2, X = x_1 + \omega x_2, Y = y_1 + \omega y_2, Z = z_1 + \omega z_2 \in \mathbb{Z}_M$$

with $c_1, c_2, x_1, x_2, y_1, y_2, z_1, z_2 \in \mathbb{Z}$.

4. A quartic relative Thue equations

A consequence of [9] is the following

Lemma 4.1. Let $x^4 + a_1x^3 + a_2x^2 + a_3x + a_4 \in \mathbb{Z}_M[x]$ be the relative defining polynomial of α over M . Let

$$\begin{aligned} F(u, v) &= u^3 - a_2u^2v + (a_1a_3 - 4a_4)uv^2 + (4a_2a_4 - a_3^2 - a_1^2a_4)v^3, \\ Q_1(x, y, z) &= x^2 - xy a_1 + y^2 a_2 + xz(a_1^2 - 2a_2) + yz(a_3 - a_1a_2) \\ &\quad + z^2(-a_1a_3 + a_2^2 + a_4) = u, \\ Q_2(x, y, z) &= y^2 - xz - a_1yz + z^2a_2 = v. \end{aligned}$$

If γ_0 , represented in the form (9) generates a relative power integral basis of K over M (that is the relative index of $I_{K/M}(\gamma_0) = (\mathbb{Z}_K : \mathbb{Z}_M[\gamma_0])$ is equal to 1), then there exist $U, V \in \mathbb{Z}_M$ such that

$$N_{M/\mathbb{Q}}(F(U, V)) = \pm 1, \quad (10)$$

$$Q_1(X, Y, Z) = U, \quad (11)$$

$$Q_2(X, Y, Z) = V. \quad (12)$$

In our case by (6) we have $a_1 = 0, a_2 = -\delta, a_3 = 0, a_4 = 1$, hence

$$F(u, v) = (u - 2v)(u + 2v)(u + \delta v).$$

If γ_0 generates a relative power integral basis of K over M , then in view of the above Lemma, together with the $X, Y, Z \in \mathbb{Z}_M$ appearing in its representation (9) there exist $U, V \in \mathbb{Z}_M$ with

$$N_{M/\mathbb{Q}}(F(U, V)) = \pm 1.$$

If $F(U, V)$ is a unit in M , then $U - 2V, U + 2V, U + \delta V$ are also units in M . Therefore

$$U - 2V = \varepsilon_1, U + 2V = \varepsilon_2$$

and

$$4V = \varepsilon_2 - \varepsilon_1.$$

In a complex quadratic field each unit is of absolute value 1, and $V \in \mathbb{Z}_M$ is of absolute value 0 or ≥ 1 . Since the right side is of absolute value ≤ 2 , the above equation implies $V = 0$. As a consequence, U is a unit in M . Then we have

$$Q_1(X, Y, Z) = \varepsilon, Q_2(X, Y, Z) = 0$$

with a unit $\varepsilon \in M$. Following the arguments of [9] we construct

$$Q_0(X, Y, Z) = UQ_2(X, Y, Z) - VQ_1(X, Y, Z) = 0,$$

whence

$$Q_2(X, Y, Z) = Y^2 - XZ - \delta Z^2 = 0.$$

$X_0 = 1, Y_0 = 0, Z_0 = 0$ is a non-trivial solution of $Q_2(X, Y, Z) = 0$. Using an argument of L. J. Mordell [12] we parametrize X, Y, Z with $R, P, Q \in M$:

$$\begin{aligned} X &= RX_0 \\ Y &= RY_0 + P \\ Z &= RZ_0 + Q \end{aligned} \tag{13}$$

Substituting this representation of X, Y, Z into $Q_2(X, Y, Z) = 0$ we obtain

$$RQ = P^2 - \delta Q^2.$$

We multiply by Q the equations in (13) and replace RQ by $P^2 - \delta Q^2$, then

$$\begin{aligned} kX &= P^2 && -\delta Q^2, \\ kY &= && PQ, \\ kZ &= && Q^2, \end{aligned} \tag{14}$$

with a $k \in M$. Further, applying the arguments of [9], in (14) we can replace the parameters $k, P, Q \in M$ by integer parameters in \mathbb{Z}_M , and it follows from the form of the above coefficient matrix of P^2, PQ, Q^2 (and the property of γ_0 being a generator of a relative power integral basis) that k is a unit in M . Finally, we substitute the representation (14) into $Q_1(X, Y, Z) = U$ and then we obtain

$$F(P, Q) = P^4 - \delta P^2 Q^2 + Q^4 = \varepsilon, \tag{15}$$

with a unit ε . This is a quartic relative Thue equation over the quadratic subfield M . As $F(x, 1)$ is just the relative defining polynomial of α over M , the equation can be written in the form

$$N_{K/M}(P - \alpha Q) = \varepsilon. \tag{16}$$

5. Solving the quartic relative Thue equation

M is a complex quadratic field, therefore the conjugate of any $\nu \in M$ is its complex conjugate $\bar{\nu}$. Denote by $\alpha^{(1)}, \alpha^{(2)}, \alpha^{(3)}, \alpha^{(4)}$ the relative conjugates of $\alpha \in K$ over M , corresponding to ω (these are the roots of $h(x)$ in (6)), then $\overline{\alpha^{(1)}}, \overline{\alpha^{(2)}}, \overline{\alpha^{(3)}}, \overline{\alpha^{(4)}}$ are the relative conjugates of α over M corresponding to $\bar{\omega}$. Set $P = p_1 + \omega p_2, Q = q_1 + \omega q_2$ with $p_1, p_2, q_1, q_2 \in \mathbb{Z}$.

Let $P, Q \in \mathbb{Z}_M$ be an arbitrary but fixed solution of (16). The unit ε in (16) is of absolute value 1, hence using $\beta = P - \alpha Q$ (16) implies

$$|\beta^{(1)}\beta^{(2)}\beta^{(3)}\beta^{(4)}| = 1. \tag{17}$$

Denote by i_0 the conjugate with

$$|\beta^{(i_0)}| = \min_{1 \leq j \leq 4} |\beta^{(j)}|.$$

(We have to perform all calculations for all possible values of i_0 .) Then by (17) we have $|\beta^{(i_0)}| \leq 1$, whence

$$|P| \leq |\beta^{(i_0)}| + |\overline{\alpha}| |Q| \leq 1 + |\overline{\alpha}| |Q|, \tag{18}$$

where we denote by $|\overline{\alpha}|$ the size of α , that is the maximum absolute value of its conjugates.

Our purpose is to determine $c_2, x_1, x_2, y_1, y_2, z_1, z_2$ in (9) with absolute value $\leq S = 10^{200}$. This implies $|X| = |x_1 + \omega x_2| \leq (1 + |\omega|)S$ and similarly $|Z| \leq (1 + |\omega|)S$. The representation (14) of Z implies

$$|Q| \leq \sqrt{|Z|} \leq \sqrt{(1 + |\omega|)S}.$$

The representation of X implies

$$|P|^2 \leq |X| + |\delta| |Q|^2 \leq |X| + |\delta| |Z|$$

whence

$$|P| \leq \sqrt{(1 + |\omega|)(1 + |\delta|)S}.$$

therefore

$$\max(|P|, |Q|) \leq \max(|X|, |Y|) \leq \sqrt{(1 + |\omega|)(1 + |\delta|)S}. \tag{19}$$

We have

$$|p_1| = \frac{|\overline{\omega}P - \omega\overline{P}|}{|\overline{\omega} - \omega|} \leq \frac{2|\omega||P|}{|\overline{\omega} - \omega|}, \quad |p_2| = \frac{|P - \overline{P}|}{|\omega - \overline{\omega}|} \leq \frac{2|P|}{|\omega - \overline{\omega}|} \tag{20}$$

and similarly

$$|q_1| \leq \frac{2|\omega||Q|}{|\overline{\omega} - \omega|}, \quad |q_2| \leq \frac{2|Q|}{|\omega - \overline{\omega}|}.$$

These imply

$$\begin{aligned} A &= \max(|p_1|, |p_2|, |q_1|, |q_2|) \leq \\ &\leq \frac{2|\omega|}{|\omega - \bar{\omega}|} \max(|P|, |Q|) \leq \frac{2|\omega|}{|\omega - \bar{\omega}|} \sqrt{(1 + |\omega|)(1 + |\delta|)S}, \end{aligned} \quad (21)$$

that is we have to determine the solutions of (16) until this bound. Note that for $S = 10^{200}$ this bound is of magnitude 10^{100} .

Further, together with (18) we have

$$A \leq \frac{2|\omega|}{|\omega - \bar{\omega}|} \max(|P|, |Q|) \leq c_1|Q|, \quad (22)$$

with

$$c_1 = \frac{2|\omega|}{|\omega - \bar{\omega}|} (1 + |\alpha|).$$

If $|Q| \geq 10$ then for $1 \leq j \leq 4, j \neq i_0$ this yields

$$\begin{aligned} |\beta^{(j)}| &\geq |\beta^{(j)} - \beta^{(i_0)}| - |\beta^{(i_0)}| \geq \\ &\geq |\alpha^{(j)} - \alpha^{(i_0)}||Q| - 1 \geq (|\alpha^{(j)} - \alpha^{(i_0)}| - 0.1)|Q|. \end{aligned} \quad (23)$$

In our calculations we have to check all possible q_1, q_2 with $|Q| < 10$ separately. (17) and (23) imply

$$|\beta^{(i_0)}| = \frac{1}{\prod_{\substack{1 \leq j \leq 4 \\ j \neq i_0}} |\beta^{(j)}|} \geq \frac{1}{\prod_{\substack{1 \leq j \leq 4 \\ j \neq i_0}} (|\alpha^{(j)} - \alpha^{(i_0)}| - 0.1)} |Q|^{-3} \leq c_{2,i_0} A^{-3}, \quad (24)$$

with

$$c_{2,i_0} = \frac{c_1^3}{\prod_{\substack{1 \leq j \leq 4 \\ j \neq i_0}} (|\alpha^{(j)} - \alpha^{(i_0)}| - 0.1)}$$

(depending on i_0).

6. Reduction

We apply a reduction procedure to reduce the bound in (22), using inequality (24), that is

$$|p_1 + \omega p_2 - \alpha^{(i_0)} q_1 - \omega \alpha^{(i_0)} q_2| \leq c_2 A^{-3}. \quad (25)$$

We follow the arguments of [1]. Let H be a large constant to be determined appropriately (for a practical choice of H see later). Consider the lattice generated by the columns of the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ H & H\Re(\omega) & H\Re(-\alpha^{(i_0)}) & H\Re(-\alpha^{(i_0)}\omega) \\ 0 & H\Im(\omega) & H\Im(-\alpha^{(i_0)}) & H\Im(-\alpha^{(i_0)}\omega) \end{pmatrix}.$$

Lemma 6.1. (cf. [1], or Lemma 5.3 of [2]) Denote by ℓ_1 the first vector of the LLL reduced basis of this lattice. If $A \leq A_0$ and H is large enough to have

$$|\ell_1| \geq \sqrt{40} \cdot A_0, \tag{26}$$

then

$$A \leq \left(\frac{c_{2,i_0} \cdot H}{A_0} \right)^{1/3}. \tag{27}$$

Note that this procedure must be performed for all possible values of i_0 .

We start with the upper bound A_0 in (21). For a certain A_0 usually A_0^2 , $10 \cdot A_0^2$ or $100 \cdot A_0^2$ is a suitable choice for H . We have to make H so large that (26) is satisfied. In view of (27) the new bound for A will be of magnitude $A_0^{1/3}$ in the first reduction steps. The following steps of the reduction is not so fast anymore, but in about 8-10 steps the original bound of magnitude 10^{100} is reduced to about 10. A typical sequence is the following:

step	A_0	H	new A_0
1	10^{100}	10^{202}	$9.1198 \cdot 10^{33}$
2	$9.1198 \cdot 10^{33}$	$8.3172 \cdot 10^{69}$	$8.8440 \cdot 10^{11}$
3	$8.8440 \cdot 10^{11}$	$7.8217 \cdot 10^{25}$	87540.0136
4	87540.0136	$7.6632 \cdot 10^{11}$	187.9568
5	187.9568	$3.5327 \cdot 10^6$	24.2479
6	24.2479	58796.3577	12.2522
7	12.2522	15011.6532	9.7587

The reduction process is very fast, it usually only takes a few seconds. For a constant H of magnitude 10^{200} we have to use multiply precision arithmetic with about 250 digits.

7. Determining P and Q

We return to the quartic relative Thue equation (15), that is

$$P^4 - \delta P^2 Q^2 + Q^4 - \varepsilon = 0. \tag{28}$$

Let A_R be the reduced bound for A obtained in the previous section.

By $m \equiv 1 \pmod{4}$, $|Q| < 10$ yields $|q_1 + \frac{1+\sqrt{m}}{2}q_2| < 10$ whence $|q_2| < 20/\sqrt{|m|}$ and $|q_1| < 10 + |q_2|/2 < 10 + 10/\sqrt{|m|}$.

Set $S_1 = 10 + 10/\sqrt{|m|}$, $S_2 = 20/\sqrt{|m|}$. Let $A_1 = \max(A_R, S_1)$, $A_2 = \max(A_R, S_2)$.

We let q_1 run up to $|q_1| \leq A_1$ and q_2 run up to $|q_2| \leq A_2$. For each pair (q_1, q_2) we calculate $Q = q_1 + \omega q_2$, substitute it into (28), and for all possible unit $\varepsilon \in M$ we solve the quartic polynomial equation (28) for the complex number P . Having the real and complex parts of P we can determine p_1, p_2 with $P = p_1 + \omega p_2$ (similarly as in (20)) and check if these values of p_1, p_2 are integers.

Having P and Q we can determine X, Y, Z from (14). Recall that k in (14) is a unit of M . Therefore all generators of relative power integral bases of K over M are of the form $C + \varepsilon(\alpha X + \alpha^2 Y + \alpha^3 Z)$ with arbitrary $C \in \mathbb{Z}_M$ and arbitrary unit $\varepsilon \in M$.

8. Determining generators of power integral bases of K

For all possible X, Y, Z as calculated above, we set $\gamma_0 = \alpha X + \alpha^2 Y + \alpha^3 Z$. In view of Theorem 3.1 all generators of power integral bases are of the form

$$\gamma = c_1 + \omega c_2 + \varepsilon \gamma_0, \quad (29)$$

with $c_1, c_2 \in \mathbb{Z}_M$, ε is a unit in M . In order to determine all non-equivalent generators of power integral bases of M we have to determine ε and c_2 so that $I(\gamma) = 1$. For this purpose we shall use the following consequence of Proposition 1 of [8]. Here we denote by $\gamma^{(1,j)}$ the conjugates of γ corresponding to $\alpha^{(j)}$ and by $\gamma^{(2,j)}$ the conjugates of γ corresponding to $\overline{\alpha^{(j)}}$ for $1 \leq j \leq 4$.

Lemma 8.1.

$$I(\gamma) = I_{K/M}(\gamma) \cdot J(\gamma)$$

where

$$I_{K/M}(\gamma) = \frac{1}{\sqrt{|N_{M/\mathbb{Q}}(D_{K/M})|}} \prod_{i=1}^2 \prod_{1 \leq j_1 \leq j_2 \leq 4} |\gamma^{(i,j_1)} - \gamma^{(i,j_2)}|$$

is the relative index of α and

$$J(\gamma) = \frac{1}{|D_M|^2} \prod_{j_1=1}^4 \prod_{j_2=1}^4 |\gamma^{(1,j_1)} - \gamma^{(2,j_2)}|.$$

In view of Lemma 4.1 we calculated γ_0 to have relative index 1. Any γ of type (29) is relative equivalent to γ_0 , that is their relative indices are equal. Therefore, we have to determine ε and c_2 using $J(\gamma) = 1$. For all of the few possible units ε of M , we calculate $J(\gamma)$. The equation

$$\prod_{j_1=1}^4 \prod_{j_2=1}^4 (\gamma^{(1,j_1)} - \gamma^{(2,j_2)}) \pm D_M^2 = 0$$

is a polynomial equation with rational integer coefficients of degree 16. To determine the possible values (if any) of $c_2 \in \mathbb{Z}$, corresponding to ε , we have to determine the integer roots in c_2 of this polynomial. Note that $D_M = a^2 - 4b + 8$.

9. Results of our calculations

Our routines were written in Maple. We made calculations for several pairs (a, b) , such that the polynomial $f(x)$ is irreducible, monogenic and m is square-free. These pairs seldom satisfied the conditions of Theorem 2.1. Note that we also made calculations in cases when m is not square-free (then $m = m_0 \cdot m_1^2$ with square-free m_1 and ω can be either $(1 + \sqrt{m_1})/2$ or $\sqrt{m_1}$) and we had similar experiences.

The table below summarizes generators of power integral bases of K , represented in the form

$$\gamma = (c_1 + \omega c_2) + (x_1 + \omega x_2)\alpha + (y_1 + \omega y_2)\alpha^2 + (z_1 + \omega z_2)\alpha^3.$$

We let (a, b) run in $-25 \leq a \leq 25, 2 \leq b \leq 25$ and took those pairs (a, b) for which $f(x)$ is irreducible, monogenic and m is square-free. In these 51 examples, it took 526 seconds (using an average PC) to calculate all generators of power integral bases with coefficients $\leq 10^{200}$ in absolute value. We list (a, b, m) and then the coefficients $[c_2, x_1, x_2, y_1, y_2, z_1, z_2]$ of generators of power integral bases. We omit the trivial $[0, 1, 0, 0, 0, 0, 0]$.

$(-9, 23, -3)$,	$[0, 4, 1, 0, 0, -1, 0]$		
$(-7, 15, -3)$,	$[0, 3, 1, 0, 0, -1, 0]$		
$(-7, 19, -19)$,	$[0, 3, 1, 0, 0, -1, 0]$		
$(-7, 23, -35)$,	$[0, 3, 1, 0, 0, -1, 0]$		
$(-5, 9, -3)$,	$[0, 2, 1, 0, 0, -1, 0]$,	$[1, -2, 1, 1, -1, 1, -1]$,	$[1, 2, -1, 1, -1, -1, 1]$
$(-5, 10, -7)$,	$[0, 2, 1, 0, 0, -1, 0]$		
$(-5, 11, -11)$,	$[0, 2, 1, 0, 0, -1, 0]$		
$(-5, 14, -23)$,	$[0, 2, 1, 0, 0, -1, 0]$		
$(-5, 18, -39)$,	$[0, 2, 1, 0, 0, -1, 0]$		
$(-5, 19, -43)$,	$[0, 2, 1, 0, 0, -1, 0]$		
$(-5, 21, -51)$,	$[0, 2, 1, 0, 0, -1, 0]$		
$(-5, 22, -55)$,	$[0, 2, 1, 0, 0, -1, 0]$		
$(-5, 23, -59)$,	$[0, 2, 1, 0, 0, -1, 0]$		
$(-5, 25, -67)$,	$[0, 2, 1, 0, 0, -1, 0]$		
$(-3, 7, -11)$,	$[0, 1, 1, 0, 0, -1, 0]$		
$(-3, 15, -43)$,	$[0, 1, 1, 0, 0, -1, 0]$		
$(-1, 3, -3)$,	$[0, 0, 1, 0, 0, -1, 0]$,	$[0, 1, -1, 0, 0, 0, 1]$,	$[0, 1, 0, 0, 0, -1, 1]$,
		$[0, 1, -1, 0, 0, 0, 0]$,	$[0, 0, 1, 0, 0, 0, 0]$
$(-1, 6, -15)$,	$[0, 0, 1, 0, 0, -1, 0]$		
$(-1, 7, -19)$,	$[0, 0, 1, 0, 0, -1, 0]$		
$(-1, 10, -31)$,	$[0, 0, 1, 0, 0, -1, 0]$		
$(-1, 11, -35)$,	$[0, 0, 1, 0, 0, -1, 0]$		
$(-1, 13, -43)$,	$[0, 0, 1, 0, 0, -1, 0]$		
$(-1, 15, -51)$,	$[0, 0, 1, 0, 0, -1, 0]$		
$(-1, 17, -59)$,	$[0, 0, 1, 0, 0, -1, 0]$		
$(-1, 19, -67)$,	$[0, 0, 1, 0, 0, -1, 0]$		

$(-1, 22, -79),$	$[0, 0, 1, 0, 0, -1, 0]$		
$(1, 3, -3),$	$[0, -1, 1, 0, 0, -1, 0],$	$[0, 0, 0, 1, -1, 0, 1],$	$[0, 0, 1, 0, 0, -1, 1],$
		$[0, 1, 0, 0, 0, 0, 1],$	$[0, 0, 0, 1, -1, 0, -1],$
		$[1, -1, 2, -1, 1, -1, 0],$	$[1, 1, -2, -1, 1, 1, 0],$
		$[0, 1, -1, 0, 0, 0, 0],$	$[0, 0, 1, 0, 0, 0, 0]$
$(1, 7, -19),$	$[0, -1, 1, 0, 0, -1, 0]$		
$(1, 11, -35),$	$[0, -1, 1, 0, 0, -1, 0]$		
$(1, 15, -51),$	$[0, -1, 1, 0, 0, -1, 0]$		
$(1, 19, -67),$	$[0, -1, 1, 0, 0, -1, 0]$		
$(3, 5, -3),$	$[0, -2, 1, 0, 0, -1, 0],$	$[0, 1, -2, 0, 0, 1, -1],$	$[0, -1, -1, 0, 0, 0, -1],$
		$[-1, 1, 0, 0, -1, 1, 0],$	$[1, 1, 0, 0, 1, 1, 0],$
		$[0, 0, -2, 0, -1, 1, -1],$	$[0, 0, -2, 0, 1, 1, -1],$
		$[0, 1, -1, 0, 0, 0, 0],$	$[0, 0, 1, 0, 0, 0, 0]$
$(3, 6, -7),$	$[0, -2, 1, 0, 0, -1, 0]$		
$(3, 7, -11),$	$[0, -2, 1, 0, 0, -1, 0]$		
$(3, 9, -19),$	$[0, -2, 1, 0, 0, -1, 0]$		
$(3, 14, -39),$	$[0, -2, 1, 0, 0, -1, 0]$		
$(3, 15, -43),$	$[0, -2, 1, 0, 0, -1, 0]$		
$(3, 18, -55),$	$[0, -2, 1, 0, 0, -1, 0]$		
$(3, 21, -67),$	$[0, -2, 1, 0, 0, -1, 0]$		
$(3, 25, -83),$	$[0, -2, 1, 0, 0, -1, 0]$		
$(5, 11, -11),$	$[0, -3, 1, 0, 0, -1, 0]$		
$(5, 19, -43),$	$[0, -3, 1, 0, 0, -1, 0]$		
$(5, 23, -59),$	$[0, -3, 1, 0, 0, -1, 0]$		
$(7, 15, -3),$	$[0, -4, 1, 0, 0, -1, 0],$		
$(7, 17, -11),$	$[0, -4, 1, 0, 0, -1, 0],$		
$(7, 18, -15),$	$[0, -4, 1, 0, 0, -1, 0],$		
$(7, 19, -19),$	$[0, -4, 1, 0, 0, -1, 0],$		
$(7, 22, -31),$	$[0, -4, 1, 0, 0, -1, 0],$		
$(7, 23, -35),$	$[0, -4, 1, 0, 0, -1, 0],$		
$(7, 25, -43),$	$[0, -4, 1, 0, 0, -1, 0],$		
$(9, 23, -3),$	$[0, -5, 1, 0, 0, -1, 0],$		

10. Another solution

In addition to $[0, 1, 0, 0, 0, 0, 0]$ in all cases considered, there appeared another solution with $[0, k, 1, 0, 0, -1, 0]$ where k seems to be related to a . The above table was constructed to indicate this relation clearly. The vector $[0, k, 1, 0, 0, -1, 0]$ yields the element

$$\alpha(k + \omega) - \alpha^3.$$

If we try find the corresponding X, Y, Z in the form (14), we get $X = k + \omega = P^2 - \delta Q^2, Y = 0 = PQ, Z = -1 = Q^2$, which is not possible. But if we consider the negative of this element, that is

$$\alpha(-k - \omega) - \alpha^3,$$

then we have $X = -k - \omega = P^2 - \delta Q^2$, $Y = 0 = PQ$, $Z = 1 = Q^2$ which has the solution $P = 0$, $Q = 1$, whence

$$-k - \omega = -\delta.$$

This implies

$$k + \frac{1 + \sqrt{m}}{2} = \frac{-a + \sqrt{m}}{2},$$

implying

$$k = -\frac{a + 1}{2}.$$

Indeed, this is shown by the examples. There remained to prove it formally.

Theorem 10.1. *If $m = a^2 - 4b + 8$ is square free, then*

$$\gamma = \left(\frac{a + 1}{2} - \omega \right) \alpha - \alpha^3 \quad (30)$$

generates a power integral basis in K .

Proof. We have

$$\frac{a + 1}{2} - \omega = \frac{a - \sqrt{m}}{2}.$$

Using the notation of Lemma 8.1 we have

$$\gamma^{(1,j)} = \frac{a - \sqrt{m}}{2} \cdot \alpha^{(j)} + (\alpha^{(j)})^3,$$

$$\gamma^{(2,j)} = \frac{a + \sqrt{m}}{2} \cdot \overline{\alpha^{(j)}} + (\overline{\alpha^{(j)}})^3,$$

for $1 \leq j \leq 4$. As we have seen above, the element (30) satisfies $I_{K/M}(\gamma) = 1$ (it comes from a valid representation of X, Y, Z by suitable P, Q in (14)), therefore we only have to check $J(\gamma) = 1$. Using symmetric polynomials ($\alpha^{(j)}$ satisfies $x^4 - \delta x^2 + 1 = 0$ and $\overline{\alpha^{(j)}}$ satisfies $x^4 - \delta x^2 + 1 = 0$) we calculated

$$\prod_{j_1=1}^4 \prod_{j_2=1}^4 |\gamma^{(1,j_1)} - \gamma^{(2,j_2)}|,$$

and, making all possible simplifications, we found that it is equal to m^2 . This calculation was also performed by Maple, and after several optimizations it took a negligible time. Note that this calculation of the above degree 16 polynomial must be made very carefully, otherwise it results unusable formulas. \square

References

- [1] GAÁL. I. Calculating "small" solutions of relative Thue equations, *Exp. Math.* **24** (2015), 142-149. [MR3350521](#), [Zbl 0985.11070](#). [622](#), [626](#), [627](#)
- [2] GAÁL. I. Diophantine equations and power integral bases. Theory and algorithms, 2nd edition, *Birkhäuser, Boston*, 2019. [MR3970246](#), [Zbl 1016.11059](#). [619](#), [620](#), [621](#), [627](#)
- [3] GAÁL. I. An experiment on the monogeneity of a family of trinomials, *JP J. Algebra Number Theory Appl.*, **51** (2021), 97–111. [Zbl 1499.11318](#). [620](#)
- [4] GAÁL. I. On the monogeneity of totally complex pure sextic fields, *JP J. Algebra Number Theory Appl.*, **60** (2023), 85–96. [Zbl 1538.11191](#). [620](#)
- [5] GAÁL. I. On the monogeneity of totally complex pure octic fields, [arXiv:2402.09293](#), 2024. [620](#)
- [6] GAÁL. I. A note on the monogeneity of some trinomials of type $x^4 + ax^2 + b$, *JP J. Algebra Number Theory Appl.*, **63** (2024), 265–279. [Zbl 1549.11164](#). [620](#)
- [7] GAÁL. I. Calculating power integral bases in some quartic fields corresponding to monogenic families of polynomials, *JP J. Algebra Number Theory Appl.*, **64** (2025), 99–115. [Zbl 07988331](#). [620](#)
- [8] GAÁL, I.; REMETE, L.; SZABÓ, T. Calculating power integral bases by using relative power integral bases, *Functiones Appr.*, **54** (2016), 141.149. [MR3513574](#), [Zbl 1395.11121](#). [623](#), [628](#)
- [9] GAÁL, I.; POHST, M. Computing power integral bases in quartic relative extensions, *J. Number Theory*, **85** (2000), 201-219. [MR1802712](#) , [Zbl 0993.11055](#). [622](#), [623](#), [624](#)
- [10] JONES, L. Monogenic even octic polynomials and their Galois groups, *New York J. Math.*, **31** (2025), 91–125. [MR4853788](#), [Zbl 07977803](#). [620](#)
- [11] JONES, L. The monogenicity and Galois groups of certain reciprocal quintinomials, [arXiv:2411.00523](#), 2024. [619](#), [620](#)
- [12] MORDELL, L. J. Diophantine Equations, *Academic Press, New York–London*, 1969. [MR0249355](#), [Zbl 0188.34503](#). [624](#)

(István Gaál) UNIVERSITY OF DEBRECEN, MATHEMATICAL INSTITUTE, H-4002 DEBRECEN PF.400., HUNGARY
gaal.istvan@unideb.hu

This paper is available via <http://nyjm.albany.edu/j/2025/31-21.html>.