

Transfer of quantum game strategies

Gage Hoefer

ABSTRACT. We develop a method for the transfer of perfect strategies between various classes of two-player, one round cooperative non-local games with quantum inputs and outputs using the simulation paradigm in quantum information theory. We show that such a transfer is possible when canonically associated operator spaces for each game are quantum homomorphic or isomorphic, as defined in [21]. We examine a new class of QNS correlations, needed for the transfer of strategies between games, and characterize them in terms of states on tensor products of canonical operator systems. We define jointly tracial correlations and show they correspond to traces acting on tensor products of canonical C^* -algebras associated with individual game parties. We then make an inquiry into the initial application of such results to the study of concurrent quantum games.

CONTENTS

1. Introduction	1607
2. Preliminaries	1611
3. Strongly stochastic operator matrices	1614
4. Strategy transport	1625
5. Perfect strategies for various quantum games	1629
6. Characterization of SQNS correlations and applications to concurrent games	1634
References	1653

1. Introduction

In the past few decades, non-local games have been studied under a variety of names (such as Bell inequalities) and across the disciplines of physics, mathematics, and computer science; they have significant connections to areas as diverse as noncommutative geometry, quantum complexity theory, entanglement theory, and operator algebras. The latter provides a particularly fruitful framework for approaching questions of non-locality in quantum systems, as the input-output behavior of measurements on bipartite quantum systems can

Received June 29, 2025.

2020 *Mathematics Subject Classification.* 47L07, 47L25, 81P16, 81R50.

Key words and phrases. Non-local game, quantum no-signalling correlations, operator systems, game algebra.

be encoded through noncommutative operator algebras and their state spaces. The study of such bipartite systems can therefore be translated into the study of associated operator algebras, and so numerous tools from functional analysis can be applied to help obtain answers for more physically motivated questions.

A non-local game is (formally speaking) a tuple $\mathbb{G} = (X, Y, A, B, \lambda)$ of finite sets X, Y, A, B and a function $\lambda : X \times Y \times A \times B \rightarrow \{0, 1\}$. The game is played cooperatively by two players, against a referee. Our two players— call them Alice and Bob— are separated spatially, and are not allowed to communicate during the game. For our purposes, the game takes place over a single round; during a round, the referee samples question pairs $(x, y) \in X \times Y$, and sends question x to Alice and question y to Bob. Alice and Bob must respond with answers $a \in A$ and $b \in B$, respectively. The two win the round if and only if λ evaluates to 1 on their question-answer pairs; that is, they win if and only if $\lambda(x, y, a, b) = 1$.

While our two players are not allowed to communicate during the game, to improve their chances of success they can coordinate their answers according to a predetermined strategy. Our players may have access to a shared quantum entangled state, and measurements on this state by each player can improve their chance of winning by coordinating their answers to the referee in such a way that would not be possible classically (see [2]: the CHSH inequality and its use in the proof of the celebrated “Bell’s Theorem” during the 1960’s, or [8]). Specific classes of strategies— known as correlations— based on the use (or non-use) of shared classical and quantum resources between the two players are of particular interest in describing their behavior; such correlations are formalized using classical information channels, when considered as conditional probability distributions. Different mathematical models corresponding to each strategy type describe the outcomes of these experiments: local (corresponding to the use of classical resources), quantum (corresponding to finite-dimensional entangled resources), quantum approximate (corresponding to liminal entangled resources), quantum commuting (which arises from the commuting model of quantum mechanics), and general no-signalling (which does not necessarily rely on the use of a shared resource, but as a probabilistic strategy must still satisfy the basic constraints of the game), are the main correlation classes of interest. These are denoted as $\mathcal{C}_{\text{loc}}, \mathcal{C}_{\text{q}}, \mathcal{C}_{\text{qa}}, \mathcal{C}_{\text{qc}}$ and $\mathcal{C}_{\text{ns}}$, respectively.

One particular connection between the study of quantum information theory, non-locality, and operator algebras driving much of the recent development in these areas is the equivalence of Tsirelson’s problem in quantum physics, and Connes’ embedding problem (or CEP) in von Neumann algebra theory; this equivalence was established in [18, 23, 31]. The subsequent investigation of this equivalence led to the resolution of many other important questions including a refutation to the strong Tsirelson problem in [35] (see also [16]), and a negative answer to the CEP in [22]. Non-local games lay at the base of all of

these approaches, and questions involving non-local games are the main motivation for the current work.

While classical non-local games are fruitful objects of study, as combinatorial objects with finite sets of inputs and outputs they ultimately have inherent restrictions; in an attempt to surpass some of these limitations, more attention has been given to *quantum games*. These are non-local games where the inputs and outputs are allowed to be quantum states, or sometimes mixtures of classical and quantum states. In this setting, question and answer sets X, Y, A and B are replaced by spaces $\mathbb{C}^{|X|}, \mathbb{C}^{|Y|}, \mathbb{C}^{|A|}$ and $\mathbb{C}^{|B|}$, and strategies are implemented using quantum channels $\Phi : M_X \otimes M_Y \rightarrow M_A \otimes M_B$ instead of classical channels $\mathcal{N} : X \times Y \rightarrow A \times B$. The lack of communication between players is enforced by strictly requiring the use of *quantum no-signalling (QNS)* channels, as introduced in [15]. Furthermore, the hierarchy of classical correlations is replaced by their quantum analogues, first introduced in [39] (see also [4]). The rules of the game can be generalized to the quantum context by replacing λ with a 0-preserving and join-preserving map between the projection lattices $\mathcal{P}_{XY}$ and $\mathcal{P}_{AB}$ of $M_X \otimes M_Y$ and $M_A \otimes M_B$, respectively. Such games have been increasingly studied over the past few years (see [6, 7, 11, 34, 39] for a non-exhaustive list). As legitimate generalizations of classical non-local games, the hope is that by “enlarging” the space of possible inputs and outputs (when compared to finite sets), a wealth of new examples are provided that might help shed more light on some of the previously mentioned questions in all areas.

The purpose of the present paper is to generalize the results of [20] to the context of quantum non-local games; in that work, generalized homomorphisms and isomorphisms between classical non-local games were introduced. The existence of a homomorphism or isomorphism of type t from game $\mathbb{G}_1$ to game $\mathbb{G}_2$ lead to a relation between optimal game values, when playing with strategies of type t ; specifically, an inequality of values in the former case, and equality in the latter case. We wanted to obtain similar results for quantum games, and identify necessary conditions for when two quantum games are similar in this sense. In order to identify when two quantum games are homomorphic (respectively, isomorphic) of some type t , we looked at several classes of quantum games which have canonically associated *quantum hypergraphs*. These are subspaces of linear operators acting between finite-dimensional spaces which in some sense encode the properties or rules of the game. Using the notion of t -homomorphism and t -isomorphism of quantum hypergraphs introduced in [21], we thus had a way to characterize when our quantum games were similar.

Quantum game homomorphisms of a given type t were defined using QNS correlations of the same type, subject to additional constraints. Such conditions were added in order to allow the transport of perfect strategies of type t from the first game to perfect strategies of type t for the second. As in [20, 21], t -isomorphisms require the use of QNS bicorrelations, which were first defined in [7]. This process of strategy transfer employs the simulation paradigm for quantum channels (see [15]). According to the paradigm, if we start with a

quantum channel $\mathcal{E} : M_{X_1} \rightarrow M_{A_1}$ from alphabet X_1 to alphabet A_1 , using assistance from no-signalling resources over (X_2, A_1, X_1, A_2) (i.e., a QNS channel $\Gamma : M_{X_2} \otimes M_{A_1} \rightarrow M_{X_1} \otimes M_{A_2}$) we can construct a new quantum channel $\Gamma[\mathcal{E}] : M_{X_2} \rightarrow M_{A_2}$ dependent on $\mathcal{E}$ and Γ from alphabet X_2 to A_2 . A similar approach for classical channels was used in [20], and one of the main focuses of this work is to show how it extends to the quantum case.

We now describe the organization of the paper in more detail. In Section 2, we set notation and recall the definition of the main no-signalling correlation types for both classical and quantum channels, and introduce the simulation setup for quantum channels. Section 3 contains the definitions of the different types of a class of stochastic (and bistochastic) operator matrices, and how these operator matrices will be used to define necessary subclasses of QNS correlations (which we call strongly quantum no-signalling) over the quadruple $(X_2 \times Y_2) \times (A_1 \times B_1) \times (X_1 \times Y_1) \times (A_2 \times B_2)$. In Section 4, the strongly quantum no-signalling correlations are used in the simulation paradigm for quantum channels to establish strategy transport for quantum games; this is achieved in Theorem 4.1 and Theorem 4.2 for the QNS correlation and bicorrelation cases, respectively. We note that the definitions of Sections 3 and 3.1 generalize those from [20, Section 4, 5]; thus, we recover many of the results (in the classical case) from [20] in Sections 3-4.

Section 5 contains some of the main results of the paper, where we restrict our attention to the transfer of perfect strategies for various types of quantum games. This application employs SQNS correlations as strategies for the t-homomorphism and isomorphism games between quantum non-local games, when utilizing the framework of generalized homomorphisms of quantum hypergraphs as introduced in [21, Section 3]. A characterization of when perfect strategy transfer is possible between games when the first leg is classical while the second is quantum, along with when both are implication games are included (see [39] for relevant introductions).

The last section features the other main focus of the paper, wherein we investigate our new QNS correlations leading up to an operator system characterization for each subclass (similar to those obtained in [29, 39] and [20, Section 7]). Using these results, we also focus on the transfer of strategies between *concurrent* quantum games—a class of quantum game first introduced in [6] and further developed in [7, 39]. As a proposed quantization of the class of synchronous games, and analogous to the adaptation of the synchronicity condition for classical game homomorphisms which necessitated the definition of *jointly synchronous* correlations in [20, Section 8], we define *jointly tracial* SQNS correlations in this section. A tracial characterization of jointly tracial correlations is contained in Theorem 6.17 (in the same vein as the characterizations established in [6], [7] and [19]), and Theorem 6.18 shows that these are the right subclass of SQNS correlations to use for transferring tracial correlations. Thus, the transfer of strategies for quantum games developed within specializes to

the concurrent case. Finally, initial connections between the transfer of concurrent/tracial strategies and traces on canonically associated $*$ -algebras and C^* -algebras for concurrent games are discussed.

Acknowledgements. I would like to thank my advisor Ivan G. Todorov for the many fruitful discussions on earlier drafts of the paper. I would also like to thank Lyudmila Turowska for helpful comments on some results in the last section, and Alexandros Chatzinikolaou for identifying the gap in the proof of [20, Lemma 5.8], necessitating the change in definition. Finally, I would like to thank the referee(s) for their detailed reading and remarks.

2. Preliminaries

In this section we set notation, and include the necessary preliminaries on quantum no-signalling correlations to be used throughout the rest of the paper. For a finite set X , we let $\mathbb{C}^X = \bigoplus_{x \in X} \mathbb{C}$ and write $(e_x)_{x \in X}$ for the canonical orthonormal basis of $\mathbb{C}^X$. Similarly, if H is a Hilbert space we set $H^X = \bigoplus_{x \in X} H$. If X is countable (not necessarily finite), let ℓ_2^X denote the Hilbert space of square summable sequences over $\mathbb{C}$; in the specific case that X is finite, we will sometimes write $\ell_2^X = \mathbb{C}^X$. For finite X, Y , we will often abbreviate the Cartesian product $X \times Y$ as XY for ease of use in notation. We denote by M_X the algebra of all complex matrices of size $X \times X$, and by $\mathcal{D}_X$ its subalgebra of all diagonal matrices. We write $\epsilon_{xx'}, x, x' \in X$ for the canonical matrix units in M_X , denote by Tr the trace functional on M_X , and set $\langle S, T \rangle = \text{Tr}(ST^*)$, where the adjoint is with respect to the canonical orthonormal basis. We let $\Delta_X : M_X \rightarrow \mathcal{D}_X$ denote the canonical conditional expectation on the full matrix algebra. Set

$$J_X := \sum_{x, x'} \epsilon_{xx'} \otimes \epsilon_{xx'}, \quad J_X^{\text{cl}} := \sum_{x \in X} \epsilon_{xx} \otimes \epsilon_{xx},$$

and $\tilde{J}_X := \frac{1}{|X|} J_X$. If $\mathbf{m}_X = \frac{1}{\sqrt{|X|}} \sum_{x \in X} e_x \otimes e_x$ is the maximally entangled unit vector in $\mathbb{C}^X \otimes \mathbb{C}^X$, then $\tilde{J}_X = \mathbf{m}_X \mathbf{m}_X^*$ is the corresponding rank-one projection. Note that $J_X^{\text{cl}} = \Delta_{XX}(J_X)$, and thus we may think of J_X^{cl} as the “classical” part of the (unnormalized) state J_X .

For a Hilbert space H , let $\mathcal{B}(H)$ be the C^* -algebra of all bounded linear operators on H , and denote by I_H the identity operator on H . An *operator system* in $\mathcal{B}(H)$ is a selfadjoint linear subspace $\mathcal{S} \subseteq \mathcal{B}(H)$ such that $I_H \in \mathcal{S}$. If $\mathcal{A}$ is a C^* -algebra, we denote by $\mathcal{A}^{\text{op}}$ its *opposite* C^* -algebra. As a set, $\mathcal{A}^{\text{op}}$ can be identified with $\mathcal{A}$, with $\mathcal{A}^{\text{op}} = \{a^{\text{op}} : a \in \mathcal{A}\}$; they both have the same additive, norm, and involutive structure—their only difference is in their multiplicative structure, as we set $a^{\text{op}} b^{\text{op}} = (ba)^{\text{op}}$, for $a^{\text{op}}, b^{\text{op}} \in \mathcal{A}^{\text{op}}$ in the opposite algebra.

Let X and A be finite sets. A *classical information channel* from X to A is a positive trace preserving linear map $\mathcal{N} : \mathcal{D}_X \rightarrow \mathcal{D}_A$. If $\mathcal{N}$ is an information channel, setting $p(\cdot|x) = \mathcal{N}(\epsilon_{xx})$ for each $x \in X$ it is easy to see that $\mathcal{N}$ is completely determined by its corresponding family of conditional probability distributions $\{(p(a|x))_{a \in A} : x \in X\}$.

A *quantum information channel* from X to A is a completely positive trace preserving linear map $\Phi : M_X \rightarrow M_A$. A quantum channel will be called (X, A) -classical if $\Phi = \Delta_A \circ \Phi \circ \Delta_X$. Any classical channel $\mathcal{N} : \mathcal{D}_X \rightarrow \mathcal{D}_A$, has a corresponding (X, A) -classical (quantum) channel $\Phi_{\mathcal{N}} : M_X \rightarrow M_A$ given by $\Phi_{\mathcal{N}} = \mathcal{N} \circ \Delta_X$. Conversely, any quantum channel $\Phi : M_X \rightarrow M_A$ induces a classical channel $\mathcal{N}_{\Phi} : \mathcal{D}_X \rightarrow \mathcal{D}_A$ given by $\Delta_A \circ \Phi|_{\mathcal{D}_X}$. Finally, if $\mathcal{E} : \mathcal{D}_X \rightarrow M_A$ is a (classical-to-quantum) channel, set $\Gamma_{\mathcal{E}} = \mathcal{E} \circ \Delta_X$, so $\Gamma_{\mathcal{E}}$ is a quantum channel from M_X to M_A .

In the remainder of this section, we recall the basic types of quantum and classical no-signalling correlations that will be used throughout the paper, along with establishing the simulation paradigm arising from quantum information theory. Let X, Y, A and B be finite sets. A *quantum no-signalling (QNS) correlation* [15] is a quantum channel $\Gamma : M_{XY} \rightarrow M_{AB}$ such that

$$\text{Tr}_A \Gamma(\rho_X \otimes \rho_Y) = 0 \text{ whenever } \rho_X \in M_X \text{ and } \text{Tr}(\rho_X) = 0, \quad (1)$$

and

$$\text{Tr}_B \Gamma(\rho_X \otimes \rho_Y) = 0 \text{ whenever } \rho_Y \in M_Y \text{ and } \text{Tr}(\rho_Y) = 0. \quad (2)$$

We set

$$\Gamma(aa', bb' | xx', yy') = \langle \Gamma(\epsilon_{x,x'} \otimes \epsilon_{y,y'}), \epsilon_{a,a'} \otimes \epsilon_{b,b'} \rangle;$$

thus, $(\Gamma(aa', bb' | xx', yy'))_{x,x',a,a'}^{y,y',b,b'}$ is the Choi matrix of Γ (see e.g. [32]).

A *stochastic operator matrix* acting on a Hilbert space H is a positive block operator matrix $E = (E_{x,x',a,a'})_{x,x',a,a'} \in M_{XA}(\mathcal{B}(H))$ such that $\text{Tr}_A(E) = I_X \otimes I_H$. Stochastic operator matrix E is *bistochastic* [7] if $X = A$ and we have $\text{Tr}_X(E) = I_A \otimes I_H$. For a stochastic operator matrix E over (X, A) , set

$$E_{a,a'} = (E_{x,x',a,a'})_{x,x' \in X} \in M_X \otimes \mathcal{B}(H).$$

As discussed in [39, Section 3], stochastic operator matrices E are the Choi matrices of unital completely positive maps $\Phi_E : M_A \rightarrow M_X \otimes \mathcal{B}(H)$ given by

$$\Phi_E(\epsilon_{a,a'}) = E_{a,a'}, \quad a, a' \in A. \quad (3)$$

If we let $\Phi = \Phi_E$ as in (3), for any state $\sigma \in \mathcal{T}(H)$ we let $\Gamma_{E,\sigma} : M_X \rightarrow M_A$ be the quantum channel defined via

$$\Gamma_{E,\sigma}(\rho_X) = \Phi_*(\rho_X \otimes \sigma), \quad \rho_X \in M_X. \quad (4)$$

Note here that $\Phi_* : M_X \otimes \mathcal{T}(H) \rightarrow M_A$ is the predual of the unital completely positive map Φ . If $E = (E_{x,x',a,a'})_{x,x',a,a'}$ and $F = (F_{y,y',b,b'})_{y,y',b,b'}$ are stochastic operator matrices in $M_{XA} \otimes \mathcal{B}(H)$ and $M_{YB} \otimes \mathcal{B}(H)$, respectively, such that

$$E_{x,x',a,a'} F_{y,y',b,b'} = F_{y,y',b,b'} E_{x,x',a,a'}$$

for all $x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B$, we let $E \cdot F$ be the (unique) stochastic operator matrix over (XY, AB) (see [39, Proposition 4.1]) defined by

$$(E_{x,x',a,a'} F_{y,y',b,b'})_{x,x',y,y'}^{a,a',b,b'} \in M_{XYAB} \otimes \mathcal{B}(H).$$

If ξ is a unit vector in Hilbert space H , we let $\Gamma_{E,F,\xi} = \Gamma_{E \cdot F, \xi \xi^*}$ where the latter is the quantum channel from M_{XY} to M_{AB} defined as in (4). If $E \in M_{XA} \otimes \mathcal{B}(H_A)$ and $F \in M_{YB} \otimes \mathcal{B}(H_B)$ are stochastic operator matrices, we let $E \odot F$ denote the stochastic operator matrix $E \otimes F$, considered as an element of $M_{XY} \otimes M_{AB} \otimes \mathcal{B}(H_A \otimes H_B)$.

A QNS correlation $\Gamma : M_{XY} \rightarrow M_{AB}$ is called *quantum commuting* if there exists a Hilbert space H , a unit vector $\xi \in H$ and stochastic operator matrices $E = (E_{x,x',a,a'})_{x,x',a,a'}$ and $F = (F_{y,y',b,b'})_{y,y',b,b'}$ on H such that E and F are mutually commuting, with $\Gamma = \Gamma_{E,F,\xi}$. *Quantum* QNS correlations are those for which there exist finite dimensional Hilbert spaces H_A, H_B , stochastic operator matrices $E \in M_{XA} \otimes \mathcal{B}(H_A)$ and $F \in M_{YB} \otimes \mathcal{B}(H_B)$, and a pure state $\sigma \in \mathcal{T}(H_A \otimes H_B)$ such that $\Gamma = \Gamma_{E \odot F, \sigma}$. *Approximately quantum* QNS correlations are the limits of quantum QNS correlations, while *local* QNS correlations are the convex combinations of the form $\Gamma = \sum_{i=1}^k \lambda_i \Phi_i \otimes \Psi_i$, where $\Phi_i : M_X \rightarrow M_A$ and $\Psi_i : M_Y \rightarrow M_B$ are quantum channels, $i = 1, \dots, k$. We write $\mathcal{Q}_{qc}$ (resp. $\mathcal{Q}_{qa}$, $\mathcal{Q}_q$, $\mathcal{Q}_{loc}$) for the (convex) set of all quantum commuting (resp. approximately quantum, quantum, local) QNS correlations, and note the (strict, see [39]) inclusions

$$\mathcal{Q}_{loc} \subseteq \mathcal{Q}_q \subseteq \mathcal{Q}_{qa} \subseteq \mathcal{Q}_{qc} \subseteq \mathcal{Q}_{ns}. \quad (5)$$

Let

$$\mathcal{L}_{X,A} = \left\{ (\lambda_{x,x',a,a'}) \in M_{XA} : \exists c \in \mathbb{C} \text{ s.t. } \sum_{a \in A} \lambda_{x,x',a,a} = \delta_{x,x'} c, \ x, x' \in X \right\},$$

and consider it as an operator subsystem of M_{XA} . Similarly,

$$\begin{aligned} \mathcal{L}_X &= \left\{ (\lambda_{x,x',a,a'}) \in M_{XX} : \exists c \in \mathbb{C} \text{ s.t. } \sum_{a \in X} \lambda_{x,x',a,a} = \delta_{x,x'} c \right. \\ &\quad \left. \text{and } \sum_{x \in X} \lambda_{x,x,a,a'} = \delta_{a,a'} c, \ x, x', a, a' \in X \right\} \end{aligned}$$

may be considered as an operator subsystem of M_{XX} . By [39, Proposition 5.5, Theorem 6.2], the elements Γ of $\mathcal{Q}_{ns}$ correspond canonically to elements of the tensor product $\mathcal{L}_{X,A} \otimes_{\min} \mathcal{L}_{Y,B}$ (viewed as an operator subsystem of $M_{XA} \otimes M_{YB}$), and elements in $\mathcal{Q}_{ns}^{\text{bi}}$ corresponding to elements of $\mathcal{L}_X \otimes_{\min} \mathcal{L}_Y$ [7, Proposition 3.6, Theorem 5.4].

A *classical correlation* over (X, Y, A, B) is a collection

$$p = \left\{ (p(a, b|x, y))_{a \in A, b \in B} : (x, y) \in X \times Y \right\},$$

where $(p(a, b|x, y))_{a \in A, b \in B}$ is a probability distribution for each $(x, y) \in X \times Y$. Given a classical correlation p , let $\mathcal{N}_p : \mathcal{D}_{XY} \rightarrow \mathcal{D}_{AB}$ be the classical channel given by

$$\mathcal{N}_p(\rho) = \sum_{x \in X, y \in Y} \sum_{a \in A, b \in B} p(a, b|x, y) \langle \rho(e_x \otimes e_y), e_x \otimes e_y \rangle \epsilon_{aa} \otimes \epsilon_{bb}. \quad (6)$$

If $t \in \{\text{loc}, q, \text{qa}, \text{qc}, \text{ns}\}$, let $\mathcal{C}_t$ denote the collection of all classical correlations p for which $\Gamma_{\mathcal{N}_p} \in \mathcal{Q}_t$.

Suppose X_i and Y_i ($i = 1, 2$) are finite sets. Let Γ be a QNS correlation over (X_2, Y_1, X_1, Y_2) and $\mathcal{E} : M_{X_1} \rightarrow M_{Y_1}$ be a quantum channel. If we write $\Gamma = \sum_{i=1}^k \Phi_i \otimes \Psi_i$, where $\Phi_i : M_{X_2} \rightarrow M_{X_1}$ and $\Psi_i : M_{Y_1} \rightarrow M_{Y_2}$ are linear maps for $i = 1, \dots, k$, we let $\Gamma[\mathcal{E}] : M_{X_2} \rightarrow M_{Y_2}$ be the linear map defined by setting

$$\Gamma[\mathcal{E}] = \sum_{i=1}^k \Psi_i \circ \mathcal{E} \circ \Phi_i. \quad (7)$$

It was shown in [15] that $\Gamma[\mathcal{E}]$ is a quantum channel, called therein the *simulated channel* from $\mathcal{E}$ assisted by *simulator* Γ . We note that (see [20], [21]) the Choi matrix of $\Gamma[\mathcal{E}]$ coincides with

$$\left(\sum_{x_1, x'_1} \sum_{y_1, y'_1} \Gamma(x_1 x'_1, y_2 y'_2 | x_2 x'_2, y_1 y'_1) \mathcal{E}(y_1 y'_1 | x_1 x'_1) \right)_{x_2, x'_2, y_2, y'_2}, \quad (8)$$

where the internal sums range over X_1 and Y_1 , for all $x_2, x'_2 \in X_2, y_2, y'_2 \in Y_2$. Thus, channel simulation is a process for constructing new quantum channels from given ones, with the assistance of correlations; moreover, it is natural to want the simulated channel to depend on a shared resource between two parties. If Alice and Bob have access to some shared resource (for instance: shared randomness, or entanglement, etc.), their interaction with the resource yields the no-signalling correlation Γ , and simulated channel $\Gamma[\mathcal{E}]$ is dependent on their local operations (see [12, Section II], and [15]).

3. Strongly stochastic operator matrices

Let X, Y, A and B be finite sets, and H be a Hilbert space. In the sequel, to simplify notation we will abbreviate an ordered pair $(x, y) \in X \times Y$ to xy . A stochastic operator matrix

$$P = (P_{xx', yy'}^{aa', bb'})_{xx', yy', aa', bb'} \in M_{XYAB} \otimes \mathcal{B}(H)$$

over (XY, AB) will be called a *strongly stochastic operator matrix* over (X, Y, A, B) if $\text{Tr}_B(L_{\sigma_Y}(P))$ (resp. $\text{Tr}_A(L_{\sigma_X}(P))$) is a well-defined stochastic operator matrix over (X, A) (resp. (Y, B)) and $\text{Tr}_B(L_{\sigma_Y}(P)) = \text{Tr}_B(L_{\sigma'_Y}(P))$ (resp. $\text{Tr}_A(L_{\sigma_X}(P)) = \text{Tr}_A(L_{\sigma'_X}(P))$) for each pure state $\sigma_X, \sigma'_X \in M_X$ and $\sigma_Y, \sigma'_Y \in M_Y$.

Remark 3.1. In fact, by convexity and linearity of the slice map we may assume $\sigma_X, \sigma_{X'} \in M_X$ and $\sigma_Y, \sigma_{Y'} \in M_Y$ are arbitrary states.

Remark 3.2. A positive operator $P = (P_{xy, ab})_{x, y, a, b} \in \mathcal{D}_{XYAB} \otimes \mathcal{B}(H)$ is a *no-signalling (NS) operator matrix* [20] if marginal operators

$$P_{x, a} := \sum_{b \in B} P_{xy, ab}, \quad P_{y, b} := \sum_{a \in A} P_{xy, ab}$$

are well-defined, and $(P_{x,a})_{a \in A}, (P_{y,b})_{b \in B}$ are POVM's for every $x \in X$ and $y \in Y$. If we start with a NS operator matrix P , for $x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B$ set

$$P_{xx',yy'}^{aa',bb'} = \delta_{xx'} \delta_{yy'} \delta_{aa'} \delta_{bb'} P_{xy,ab}.$$

Setting $\tilde{P} = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'} \in M_{XYAB} \otimes \mathcal{B}(H)$, one may easily check that $\tilde{P}$ is strongly stochastic over (XY, AB) with $\text{Tr}_B(L_{\sigma_Y}(\tilde{P}))$ and $\text{Tr}_A(L_{\sigma_X}(\tilde{P}))$ classical stochastic operator matrices (as introduced in [39, Section 3]) for each (pure) state $\sigma_X \in M_X$ and $\sigma_Y \in M_Y$. Thus, we may think of strongly stochastic operator matrices over (XY, AB) as generalizations of NS operator matrices, whose “marginal” stochastic operators are no longer necessarily classical.

A strongly stochastic operator matrix $P = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'}$ is called *dilatable* if there exists a Hilbert space K , an isometry $V : H \rightarrow K$ and stochastic operator matrices $(E_{xx',aa'})_{x,x' \in X, a, a' \in A}$ (resp. $(F_{yy',bb'})_{y,y' \in Y, b, b' \in B}$) in $M_{XA} \otimes \mathcal{B}(K)$ (resp. $M_{YB} \otimes \mathcal{B}(K)$) such that $E_{xx',aa'} F_{yy',bb'} = F_{yy',bb'} E_{xx',aa'}$ and

$$P_{xx',yy'}^{aa',bb'} = V^* E_{xx',aa'} F_{yy',bb'} V, \quad x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B. \quad (9)$$

We will call a strongly stochastic operator matrix $P = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'}$ acting on a Hilbert space H *locally dilatable* if there exists a dilation of the form (9), with the additional stipulation that the family

$$\{E_{xx',aa'}, F_{yy',bb'} : x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B\}$$

is commutative. We will call a strongly stochastic operator matrix

$$P = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'}$$

acting on a Hilbert space H *quantum dilatable* if there exist stochastic operator matrices $(E_{xx',aa'})_{x,x' \in X, a, a' \in A}$ and $(F_{yy',bb'})_{y,y' \in Y, b, b' \in B}$ acting on finite dimensional Hilbert spaces H_A and H_B respectively, and an isometry $V : H \rightarrow H_A \otimes H_B$, such that

$$P_{xx',yy'}^{aa',bb'} = V^* (E_{xx',aa'} \otimes F_{yy',bb'}) V, \quad (10)$$

for $x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B$.

Proposition 3.3. *A QNS correlation Γ over (X, Y, A, B) belongs to $\mathcal{Q}_{\text{loc}}$ if and only if there exists a Hilbert space H , a locally dilatable strongly stochastic operator matrix $P = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'}$ acting on H and a unit vector $\xi \in H$ such that*

$$\Gamma(aa', bb' | xx', yy') = \langle P_{xx',yy'}^{aa',bb'} \xi, \xi \rangle, \quad (11)$$

for $x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B$.

Proof. First, assume that $\Gamma \in \mathcal{Q}_{\text{loc}}$ is a convex combination $\Gamma = \sum_{j=1}^k \lambda_j \Phi^{(j)} \otimes \Psi^{(j)}$, where $\Phi^{(j)} : M_X \rightarrow M_A$ and $\Psi^{(j)} : M_Y \rightarrow M_B$ are quantum channels, $j = 1, \dots, k$. By the comment before [39, Remark 3.2], $\Phi^{(j)}$ (resp. $\Psi^{(j)}$) is of the form

$\Gamma_{E^{(j)},1}$ (resp. $\Gamma_{F^{(j)},1}$) for some stochastic operator matrix $E^{(j)} \in M_X \otimes M_A \otimes \mathcal{B}(\mathbb{C})$ (resp. $F^{(j)} \in M_Y \otimes M_B \otimes \mathcal{B}(\mathbb{C})$) for $j = 1, \dots, k$. For each $x, x' \in X, a, a' \in A$ define the matrix $E_{x,x'}^{a,a'} = (E_{x,x',a,a'}^{(j)})_{j=1}^k \in \mathcal{D}_k$. Define the corresponding matrices $F_{y,y'}^{b,b'}$ in $\mathcal{D}_k$ for each $y, y' \in Y, b, b' \in B$. We note that the family

$$\{E_{x,x'}^{a,a'}, F_{y,y'}^{b,b'} : x, x' \in X, a, a' \in A, y, y' \in Y, b, b' \in B\},$$

is commutative. Then, let

$$E := (E_{x,x'}^{a,a'})_{x,x',a,a'}, \quad F := (F_{y,y'}^{b,b'})_{y,y',b,b'}.$$

Note that E (resp. F) is stochastic over (X, A) (resp. (Y, B)). Indeed: since $E^{(j)}$ is a stochastic operator matrix in $M_X \otimes M_A \otimes \mathcal{B}(\mathbb{C})$ for each $j = 1, \dots, k$, we know that $\sum_a E_{x,x',a,a}^{(j)} = \delta_{x,x'}$ for each $x, x' \in X$ and $j = 1, \dots, k$. Thus,

$$\begin{aligned} \text{Tr}_A(E) &= \sum_a \sum_{x,x'} \sum_{j=1}^k E_{x,x',a,a}^{(j)} \epsilon_{x,x'} \otimes \epsilon_{jj} \\ &= \sum_{j=1}^k \sum_{x,x'} \sum_a E_{x,x',a,a}^{(j)} \epsilon_{x,x'} \otimes \epsilon_{jj} \\ &= \sum_{j=1}^k \sum_{x,x'} \delta_{x,x'} \epsilon_{x,x'} \otimes \epsilon_{jj} \\ &= I_X \otimes I_k. \end{aligned}$$

Set $\xi = \sum_{j=1}^k \sqrt{\lambda_j} e_j$, so ξ is a unit vector in $\mathbb{C}^k$. For $x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B$ we see

$$\begin{aligned} \langle E_{x,x'}^{a,a'} F_{y,y'}^{b,b'} \xi, \xi \rangle &= \sum_{j=1}^k \lambda_j E_{x,x',a,a'}^{(j)} F_{y,y',b,b'}^{(j)} \\ &= \sum_{j=1}^k \lambda_j \langle E_{x,x',a,a'}^{(j)} \otimes F_{y,y',b,b'}^{(j)}, I \rangle \\ &= \langle \Gamma(\epsilon_{x,x'} \otimes \epsilon_{y,y'}), \epsilon_{a,a'} \otimes \epsilon_{b,b'} \rangle \\ &= \Gamma(aa', bb' | xx', yy'). \end{aligned}$$

Setting $P_{xx',yy'}^{aa',bb'} = E_{x,x'}^{a,a'} F_{y,y'}^{b,b'}$ for each $x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B$ and letting $P = (P_{xx',yy'}^{aa',bb'})$ (where the latter matrix entries range over X, Y, A, B) gives us our locally dilatable strongly stochastic operator matrix satisfying (11).

Now, assume $P = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'}$ is a locally dilatable strongly stochastic operator matrix acting on H with unit vector $\xi \in H$ satisfying (11). If we replace H with the Hilbert space K arising from the dilation (9) of P

and the vector ξ with $V\xi$, we may without loss of generality directly assume $P_{xx',yy'}^{aa',bb'} = E_{x,x',a,a'} F_{y,y',b,b'}$, $x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B$, where the family

$$\{E_{x,x',a,a'}, F_{y,y',b,b'} : x, x' \in X, a, a' \in A, y, y' \in Y, b, b' \in B\}$$

is commutative.

Let $\mathcal{A}_X^A$ (resp. $\mathcal{A}_Y^B$) be the abelian C^* -algebra generated by $\{E_{x,x',a,a'} : x, x' \in X, a, a' \in A\}$ (resp. $\{F_{y,y',b,b'} : y, y' \in Y, b, b' \in B\}$). Let $s : \mathcal{A}_X^A \otimes_{\max} \mathcal{A}_Y^B \rightarrow \mathbb{C}$ be the state given by $s(S \otimes T) = \langle ST\xi, \xi \rangle$. By nuclearity of abelian C^* -algebras, we may consider s as a state on $\mathcal{A}_X^A \otimes_{\min} \mathcal{A}_Y^B$. Using the identification $\mathcal{A}_X^A = C(\Omega_1), \mathcal{A}_Y^B = C(\Omega_2)$ for compact Hausdorff spaces Ω_1, Ω_2 , we view s as a Borel probability measure μ on the product topological space $\Omega_1 \times \Omega_2$. For each $\omega_1 \in \Omega_1$ (resp. $\omega_2 \in \Omega_2$), let $\Phi(\omega_1) : M_X \rightarrow M_A$ (resp. $\Psi(\omega_2) : M_Y \rightarrow M_B$) be the quantum channel given by $\Phi(\omega_1)(\epsilon_{x,x'}) = (E_{x,x',a,a'}(\omega_1))_{a,a'}$ (resp. $\Psi(\omega_2)(\epsilon_{y,y'}) = (F_{y,y',b,b'}(\omega_2))_{b,b'}$). We then have

$$\begin{aligned} & \langle \Gamma(\epsilon_{x,x'} \otimes \epsilon_{y,y'}), \epsilon_{a,a'} \otimes \epsilon_{b,b'} \rangle \\ &= \langle E_{x,x',a,a'} F_{y,y',b,b'} \xi, \xi \rangle \\ &= \int_{\Omega_1 \times \Omega_2} E_{x,x',a,a'}(\omega_1) F_{y,y',b,b'}(\omega_2) d\mu(\omega_1, \omega_2) \\ &= \left\langle \int_{\Omega_1 \times \Omega_2} \Phi(\omega_1)(\epsilon_{x,x'}) \otimes \Psi(\omega_2)(\epsilon_{y,y'}) d\mu(\omega_1, \omega_2), \epsilon_{a,a'} \otimes \epsilon_{b,b'} \right\rangle. \end{aligned}$$

If we approximate measure μ using convex combinations of product measures $\mu_1 \times \mu_2$ (where $\mu_1 \in M(\Omega_1), \mu_2 \in M(\Omega_2)$) we see we can approximate Γ using channels which may be written as convex combinations $\tilde{\Gamma} = \sum_{j=1}^k \lambda_j \Phi_j \otimes \Psi_j$, where $\Phi_j : M_X \rightarrow M_A$ and $\Psi_j : M_Y \rightarrow M_B$ are quantum channels, $j = 1, \dots, k$. By the Carathéodory Theorem and compactness, use a similar argument as in [39, Remark 4.10] to conclude that Γ itself is of that form. Thus, $\Gamma \in \mathcal{Q}_{\text{loc}}$. $\square$

For certain classes of quantum games (for instance, those concerned with the behavior of quantum symmetries of quantum objects as in [7, 13]) we will draw our questions and answer states from the same space; thus, a particular notion of strongly stochastic operator matrix will be required in this context. We assume that $X = A$ and $Y = B$. For notational ease, we will continue to refer to X and A (resp. Y and B) as distinct entities, even though they are copies of the same set.

A positive operator $P = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'} \in M_{XYAB} \otimes \mathcal{B}(H)$ will be called a *strongly bistochastic operator matrix* if it is a strongly stochastic operator matrix over (XY, AB) , with $\text{Tr}_B(L_{\sigma_Y}(P))$ and $\text{Tr}_A(L_{\sigma_X}(P))$ bistochastic operator matrices for each state $\sigma_X \in M_X, \sigma_Y \in M_Y$. A strongly bistochastic operator matrix $P = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'}$ is called *dilatable* if there exists

a Hilbert space K , an isometry $V : H \rightarrow K$, and bistochastic operator matrices $(E_{xx',aa'})_{x,x' \in X, a,a' \in A}$ (resp. $(F_{yy',bb'})_{y,y' \in Y, b,b' \in B}$) in $M_{XA} \otimes \mathcal{B}(K)$ (resp. $M_{YB} \otimes \mathcal{B}(K)$) such that $E_{xx',aa'}F_{yy',bb'} = F_{yy',bb'}E_{xx',aa'}$ and which satisfy relations (9) for all $x, x' \in X, a, a' \in A, y, y' \in Y, b, b' \in B$. *Locally dilatable* and *quantum dilatable* strongly bistochastic operator matrices are defined exactly as with strongly stochastic operator matrices, but using bistochastic operator matrices in place of stochastic operator matrices.

3.1. SQNS correlations. One goal for this work is to develop a method for transferring perfect strategies of one quantum input-output game to another. The framework we use for this purpose is by embedding both quantum games into a single “game-of-games”, whose winning QNS strategies encode the winning information for both simultaneously. The winning QNS strategies are used as simulators in the simulation paradigm, allowing us to take a perfect strategy for the first game and—in conjunction with a winning strategy for the “game-of-games”—construct the desired strategy for the second. In order to adhere to the no-signalling condition for both games, we have to modify the classes of QNS correlations we use as simulators; this leads us to the introduction of a particular sub-class of QNS correlations, which is the focus of this subsection.

Let $X_i, Y_i, A_i, B_i, i = 1, 2$ be finite sets. A quantum channel

$$\Gamma : M_{X_2 Y_2 \times A_1 B_1} \rightarrow M_{X_1 Y_1 \times A_2 B_2}$$

will be called a *strongly quantum no-signalling (SQNS) correlation* if

$$\text{Tr}_{X_1} \Gamma(\rho_{X_2 Y_2} \otimes \rho_{A_1 B_1}) = 0 \text{ if } \rho_{X_2 Y_2} \in M_{X_2 Y_2} \text{ and } \text{Tr}_{X_2}(\rho_{X_2 Y_2}) = 0, \quad (12)$$

$$\text{Tr}_{Y_1} \Gamma(\rho_{X_2 Y_2} \otimes \rho_{A_1 B_1}) = 0 \text{ if } \rho_{X_2 Y_2} \in M_{X_2 Y_2} \text{ and } \text{Tr}_{Y_2}(\rho_{X_2 Y_2}) = 0, \quad (13)$$

$$\text{Tr}_{A_2} \Gamma(\rho_{X_2 Y_2} \otimes \rho_{A_1 B_1}) = 0 \text{ if } \rho_{A_1 B_1} \in M_{A_1 B_1} \text{ and } \text{Tr}_{A_1}(\rho_{A_1 B_1}) = 0, \quad (14)$$

and

$$\text{Tr}_{B_2} \Gamma(\rho_{X_2 Y_2} \otimes \rho_{A_1 B_1}) = 0 \text{ if } \rho_{A_1 B_1} \in M_{A_1 B_1} \text{ and } \text{Tr}_{B_1}(\rho_{A_1 B_1}) = 0. \quad (15)$$

We denote by $\mathcal{Q}_{\text{sns}}$ the (convex) set of all SQNS correlations; it is clear that $\mathcal{Q}_{\text{sns}} \subseteq \mathcal{Q}_{\text{ns}}$.

A *classical strongly no – signalling (SNS) correlation* [20] is a correlation

$$p = (p(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b_1))_{x_1 y_1, a_1 b_1, x_2 y_2, a_2 b_2}$$

that satisfies the conditions

$$\begin{aligned} \sum_{x_1 \in X_1} p(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b_1) &= \sum_{x_1 \in X_1} p(x_1 y_1, a_2 b_2 | x'_2 y_2, a_1 b_1), \quad x_2, x'_2 \in X_2, \\ \sum_{y_1 \in Y_1} p(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b_1) &= \sum_{y_1 \in Y_1} p(x_1 y_1, a_2 b_2 | x_2 y'_2, a_1 b_1), \quad y_2, y'_2 \in Y_2, \\ \sum_{a_2 \in A_2} p(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b_1) &= \sum_{a_2 \in A_2} p(x_1 y_1, a_2 b_2 | x_2 y_2, a'_1 b_1), \quad a_1, a'_1 \in A_1, \end{aligned}$$

and

$$\sum_{b_2 \in B_2} p(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b_1) = \sum_{b_2 \in B_2} p(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b'_1), \quad b_1, b'_1 \in B_1.$$

The (convex) collection of all SNS correlations is denoted by $\mathcal{C}_{\text{sns}}$. If $p \in \mathcal{C}_{\text{sns}}$, then marginal conditional probability distributions

$$p(x_1 y_1, a_2 | x_2 y_2, a_1), \quad p(x_1 y_1, b_2 | x_2 y_2, b_1), \\ p(x_1, a_2 b_2 | x_2, a_1 b_1) \text{ and } p(y_1, a_2 b_2 | y_2, a_1 b_1)$$

are all well-defined.

Remark 3.4. If p is a classical correlation over $(X_2 Y_2, A_1 B_1, X_1 Y_1, A_2 B_2)$, then p is an SNS correlation precisely when Γ_p is a SQNS correlation. To see why, first assume that $\rho_{X_2 Y_2} \in M_{X_2 Y_2}$ with $\text{Tr}_{X_2}(\rho_{X_2 Y_2}) = 0$. Writing

$$\rho_{X_2 Y_2} = \sum \tilde{\rho}_{x_2, x'_2}^{y_2, y'_2} \epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2},$$

where the sum is over all $x_2, x'_2 \in X_2$ and $y_2, y'_2 \in Y_2$. The partial trace condition on $\rho_{X_2 Y_2}$ implies

$$\sum_{x_2 \in X_2} \sum_{y_2, y'_2 \in Y_2} \tilde{\rho}_{x_2, x'_2}^{y_2, y'_2} \epsilon_{y_2 y'_2} = 0. \quad (16)$$

Specifically, $\sum_{x_2} \tilde{\rho}_{x_2, x'_2}^{y_2, y'_2} = 0$ for any $y_2 \in Y_2$. If we then take $\rho_{A_1 B_1} \in M_{A_1 B_1}$, we see

$$\begin{aligned} & \text{Tr}_{X_1} \Gamma_p(\rho_{X_2 Y_2} \otimes \rho_{A_1 B_1}) \\ &= \sum_{x_2, y_2} \sum_{a_1, b_1} \sum_{x_1, y_1} \sum_{a_2, b_2} \tilde{\rho}_{x_2, x'_2}^{y_2, y'_2} p(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b_1) \langle \rho_{A_1 B_1}(e_{a_1} \otimes e_{b_1}), e_{a_1} \otimes e_{b_1} \rangle \\ & \quad \times \epsilon_{y_1 y_1} \otimes \epsilon_{a_2 a_2} \otimes \epsilon_{b_2 b_2} \\ &= \sum_{a_1, b_1} \sum_{a_2, b_2} \sum_{y_1} \left(\sum_{y_2} \sum_{x_2} \tilde{\rho}_{x_2, x'_2}^{y_2, y'_2} p(y_1, a_2 b_2 | y_2, a_1 b_1) \right) \langle \rho_{A_1 B_1}(e_{a_1} \otimes e_{b_1}), e_{a_1} \otimes e_{b_1} \rangle \\ & \quad \times \epsilon_{y_1 y_1} \otimes \epsilon_{a_2 a_2} \otimes \epsilon_{b_2 b_2} \\ &= 0. \end{aligned}$$

Using a similar argument, we can check that conditions (13), (14), and (15) are satisfied. Conversely, assuming Γ_p satisfies (12)-(15), if we let

$$\rho = \epsilon_{x_2 x_2} \otimes \epsilon_{y_2 y_2} \otimes \epsilon_{a_1 a_1} \otimes \epsilon_{b_1 b_1} - \epsilon_{x'_2 x'_2} \otimes \epsilon_{y_2 y_2} \otimes \epsilon_{a_1 a_1} \otimes \epsilon_{b_1 b_1},$$

for $x_2, x'_2 \in X_2, y_2 \in Y_2, a_1 \in A_1$ and $b_1 \in B_1$ we have that $\text{Tr}_{X_2}(\rho) = 0$ with

$$\begin{aligned} \Gamma_p(\rho) &= \sum_{x_1, y_1} \sum_{a_2, b_2} (p(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b_1) - p(x_1 y_1, a_2 b_2 | x'_2 y_2, a_1 b_1)) \\ & \quad \times \epsilon_{x_1 x_1} \otimes \epsilon_{y_1 y_1} \otimes \epsilon_{a_2 a_2} \otimes \epsilon_{b_2 b_2}. \end{aligned}$$

By (12), we see

$$\sum_{x_1} p(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b_1) = \sum_{x_1} p(x_1 y_1, a_2 b_2 | x'_2 y_2, a_1 b_1), \quad x_2, x'_2 \in X_2.$$

The other SNS conditions can be verified by replacing ρ with

$$\begin{aligned} & \epsilon_{x_2 x_2} \otimes \epsilon_{y_2 y_2} \otimes \epsilon_{a_1 a_1} \otimes \epsilon_{b_1 b_1} - \epsilon_{x_2 x_2} \otimes \epsilon_{y'_2 y'_2} \otimes \epsilon_{a_1 a_1} \otimes \epsilon_{b_1 b_1}, \\ & \epsilon_{x_2 x_2} \otimes \epsilon_{y_2 y_2} \otimes \epsilon_{a_1 a_1} \otimes \epsilon_{b_1 b_1} - \epsilon_{x_2 x_2} \otimes \epsilon_{y_2 y_2} \otimes \epsilon_{a'_1 a'_1} \otimes \epsilon_{b_1 b_1}, \end{aligned}$$

and

$$\epsilon_{x_2 x_2} \otimes \epsilon_{y_2 y_2} \otimes \epsilon_{a_1 a_1} \otimes \epsilon_{b_1 b_1} - \epsilon_{x_2 x_2} \otimes \epsilon_{y_2 y_2} \otimes \epsilon_{a_1 a_1} \otimes \epsilon_{b'_1 b'_1},$$

respectively.

Thus, we see that if p is an SNS correlation over $(X_2 Y_2, A_1 B_1, X_1 Y_1, A_2 B_2)$, then Γ_p is SQNS. Also, if Γ is a $(X_2 Y_2 \times A_1 B_1, X_1 Y_1 \times A_2 B_2)$ classical SQNS correlation then $\Gamma = \Gamma_p$ for some SNS correlation p .

Remark 3.5. An easily verified alternative characterization of QNS correlations can be stated as follows: $\Phi : M_{XY} \rightarrow M_{AB}$ is QNS if and only if the unital completely positive map $\Phi^* : M_{AB} \rightarrow M_{XY}$ preserves subalgebras, in the sense that

$$\Phi^*(M_A \otimes 1) \subseteq M_X \otimes 1, \quad \Phi^*(1 \otimes M_B) \subseteq 1 \otimes M_Y.$$

The strengthening of partial trace conditions in the definition of an SQNS correlation leads to the following result.

Proposition 3.6. *A QNS correlation $\Phi : M_{X_2 Y_2, A_1 B_1} \rightarrow M_{X_1 Y_1, A_2 B_2}$ is SQNS if and only if the unital completely positive map $\Phi^* : M_{X_1 Y_1, A_2 B_2} \rightarrow M_{X_2 Y_2, A_1 B_1}$ preserves subalgebras in the sense that*

$$\Phi^*(1 \otimes M_{Y_1} \otimes M_{A_2} \otimes M_{B_2}) \subseteq 1 \otimes M_{Y_2} \otimes M_{A_1} \otimes M_{B_1}, \quad (17)$$

$$\Phi^*(M_{X_1} \otimes 1 \otimes M_{A_2} \otimes M_{B_2}) \subseteq M_{X_2} \otimes 1 \otimes M_{A_1} \otimes M_{B_1}, \quad (18)$$

$$\Phi^*(M_{X_1} \otimes M_{Y_1} \otimes 1 \otimes M_{B_2}) \subseteq M_{X_2} \otimes M_{Y_2} \otimes 1 \otimes M_{B_1}, \quad (19)$$

and

$$\Phi^*(M_{X_1} \otimes M_{Y_1} \otimes M_{A_2} \otimes 1) \subseteq M_{X_2} \otimes M_{Y_2} \otimes M_{A_1} \otimes 1. \quad (20)$$

Proof. First, let Φ be an SQNS correlation and take an arbitrary $\rho \in 1 \otimes M_{Y_1} \otimes M_{A_2} \otimes M_{B_2}$. Write

$$\rho = \sum_{y_1, y'_1} \sum_{a_2, a'_2} \sum_{b_2, b'_2} \lambda_{y_1 y'_1}^{a_2 a'_2, b_2 b'_2} 1 \otimes \epsilon_{y_1 y'_1} \otimes \epsilon_{a_2 a'_2} \otimes \epsilon_{b_2 b'_2}.$$

If we now fix $x_2, x'_2 \in X_2$ such that $x_2 \neq x'_2$, and for any $y_2, y'_2 \in Y_2, a_1, a'_1 \in A_1, b_1, b'_1 \in B_1$ we see

$$\begin{aligned}
& \langle \Phi^*(\rho), \epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2} \otimes \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1} \rangle \\
&= \sum_{y_1 y'_1} \lambda_{y_1 y'_1}^{a_2 a'_2, b_2 b'_2} \langle \Phi^*(1 \otimes \epsilon_{y_1 y'_1} \otimes \epsilon_{a_2 a'_2} \otimes \epsilon_{b_2 b'_2}), \epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2} \otimes \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1} \rangle \\
&= \sum_{y_1 y'_1} \lambda_{y_1 y'_1}^{a_2 a'_2, b_2 b'_2} \langle 1 \otimes \epsilon_{y_1 y'_1} \otimes \epsilon_{a_2 a'_2} \otimes \epsilon_{b_2 b'_2}, \Phi(\epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2} \otimes \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1}) \rangle \\
&= \sum_{y_1 y'_1} \lambda_{y_1 y'_1}^{a_2 a'_2, b_2 b'_2} \langle \epsilon_{y_1 y'_1} \otimes \epsilon_{a_2 a'_2} \otimes \epsilon_{b_2 b'_2}, \text{Tr}_{X_1} \Phi(\epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2} \otimes \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1}) \rangle \\
&= 0,
\end{aligned}$$

where the latter sums are over all $y_1, y'_1 \in Y_1, a_2, a'_2 \in A_2, b_2, b'_2 \in B_2$ as $\text{Tr}_{X_2}(\epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2}) = 0$ by our choice of $x_2, x'_2 \in X_2$. As this holds for all $y_2, y'_2 \in Y_2, a_1, a'_1 \in A_1$ and $b_1, b'_1 \in B_1$, and our choice of $\rho \in 1 \otimes M_{Y_1} \otimes M_{A_2} \otimes M_{B_2}$ was arbitrary, this implies

$$\Phi^*(1 \otimes M_{Y_1} \otimes M_{A_2} \otimes M_{B_2}) \subseteq 1 \otimes M_{Y_2} \otimes M_{A_1} \otimes M_{B_1}.$$

Using (13)-(15), we may then show that conditions (18)-(20) also hold.

Conversely, assume that Φ^* preserves subalgebras as in (17)-(20). We wish to show that Φ is an SQNS correlation; to that end, let $\rho_{X_2 Y_2} \in M_{X_2 Y_2}$ and $\rho_{A_1 B_1} \in M_{A_1 B_1}$ such that $\text{Tr}_{X_2}(\rho_{X_2 Y_2}) = 0$. Write

$$\rho_{X_2 Y_2} = \sum_{x_2, x'_2} \sum_{y_2, y'_2} \rho_{x_2 x'_2}^{y_2 y'_2} \epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2}, \quad \rho_{A_1 B_1} = \sum_{a_1, a'_1} \sum_{b_1, b'_1} \rho_{a_1 a'_1}^{b_1 b'_1} \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1}.$$

For $y_1, y'_1 \in Y_1, a_2, a'_2 \in A_2, b_2, b'_2 \in B_2$ we see

$$\begin{aligned}
& \langle \text{Tr}_{X_1} \Phi(\rho_{X_2 Y_2} \otimes \rho_{A_1 B_1}), \epsilon_{y_1 y'_1} \otimes \epsilon_{a_2 a'_2} \otimes \epsilon_{b_2 b'_2} \rangle \\
&= \sum_{x_2, x'_2} \sum_{y_2, y'_2} \rho_{x_2 x'_2}^{y_2 y'_2} \rho_{a_1 a'_1}^{b_1 b'_1} \langle \text{Tr}_{X_1} \Phi(\epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2} \otimes \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1}), \epsilon_{y_1 y'_1} \otimes \epsilon_{a_2 a'_2} \otimes \epsilon_{b_2 b'_2} \rangle \\
&= \sum_{x_2, x'_2} \sum_{y_2, y'_2} \rho_{x_2 x'_2}^{y_2 y'_2} \rho_{a_1 a'_1}^{b_1 b'_1} \langle \epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2} \otimes \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1}, \Phi^*(1 \otimes \epsilon_{y_1 y'_1} \otimes \epsilon_{a_2 a'_2} \otimes \epsilon_{b_2 b'_2}) \rangle \\
&= \sum_{x_2} \sum_{y_2, y'_2} \rho_{x_2 x_2}^{y_2 y'_2} \rho_{a_1 a'_1}^{b_1 b'_1} (\Phi^*)_{y'_2 y_2}^{a'_1 a_1, b'_1 b_1} \\
&= 0,
\end{aligned}$$

where the second summand ranges over all $y_2, y'_2 \in Y_2, a_1, a'_1 \in A_1, b_1, b'_1 \in B_1$ as $\sum_{x_2} \rho_{x_2 x_2}^{y_2 y'_2} = 0$ for any $y_2, y'_2 \in Y_2$. Note here that we are using (17) and writing

$$\Phi^*(1 \otimes \epsilon_{y_1 y'_1} \otimes \epsilon_{a_2 a'_2} \otimes \epsilon_{b_2 b'_2}) = \sum_{y_2 y'_2} (\Phi^*)_{y_2 y'_2}^{a_1 a'_1, b_1 b'_1} (1 \otimes \epsilon_{y_2 y'_2} \otimes \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1}),$$

where the latter sum is over all of Y_2, A_1 and B_1 . As our choice of $y_1, y'_1 \in Y_1, a_2, a'_2 \in A_2$ and $b_2, b'_2 \in B_2$ were arbitrary, we conclude that (12) holds. Conditions (13)-(15) are verified similarly. $\square$

Definition 3.7. An SQNS correlation Γ over $(X_2Y_2, A_1B_1, X_1Y_1, A_2B_2)$ is called

- (i) quantum commuting if there exists a Hilbert space H , stochastic operator matrices

$$E_X = (E_{x_2x'_2, x_1x'_1})_{x_2x'_2, x_1x'_1}, \quad E_Y = (E^{y_2y'_2, y_1y'_1})_{y_2y'_2, y_1y'_1},$$

$$F_A = (F_{a_1a'_1, a_2a'_2})_{a_1a'_1, a_2a'_2}, \quad F_B = (F^{b_1b'_1, b_2b'_2})_{b_1b'_1, b_2b'_2}$$

acting on H with mutually commuting entries, and a unit vector $\xi \in H$ such that $\Gamma = \Gamma_{E,F,\xi}$, where $E = E_X \cdot E_Y, F = F_A \cdot F_B$.

- (ii) quantum if there exists finite dimensional Hilbert spaces H and K , quantum dilatable strongly stochastic operator matrices

$$M = (M^{x_1x'_1, a_2a'_2})_{x_2x'_2, a_1a'_1} \text{ on } H,$$

$$N = (N^{y_1y'_1, b_2b'_2})_{y_2y'_2, b_1b'_1} \text{ on } K,$$

and a unit vector $\xi \in H \otimes K$, such that $\Gamma = \Gamma_{M \otimes N, \xi}$.

- (iii) approximately quantum if it is the limit of quantum SQNS correlations.
 (iv) local if it is quantum, and the matrices M and N from (ii) can be chosen to be locally dilatable.

We denote by $\mathcal{Q}_{\text{sqc}}$ (resp. $\mathcal{Q}_{\text{sqqa}}, \mathcal{Q}_{\text{sq}}, \mathcal{Q}_{\text{sloc}}$) the classes of quantum commuting (resp. approximately quantum, quantum, local) SQNS correlations. We note that, by definition, $\mathcal{Q}_{\text{st}} \subseteq \mathcal{Q}_{\text{t}}$, for $\text{t} \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$. In the sequel, for $\text{t} \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$ let $\mathcal{C}_{\text{st}}$ denote the class of all SNS correlations of type t , as defined in [20, Section 5].

Remark 3.8. We pause here for a correction to a previous work, and some necessary clarifying remarks. In [20], we defined a classical correlation Γ to belong to the class $\mathcal{C}_{\text{sqc}}$ if there existed a Hilbert space H , dilatable NS operator matrices $P = (P_{x_2y_2, x_1y_1})_{x_2y_2, x_1y_1}$ and $Q = (Q_{a_1b_1, a_2b_2})_{a_1b_1, a_2b_2}$ on H with mutually commuting entries, and a unit vector $\xi \in H$ such that $\Gamma = \Gamma_{P,Q,\xi}$. In [20, Lemma 5.8] we claimed that this was a necessary and sufficient condition for the existence of a Hilbert space K , PVM's $(P_{x_2, x_1})_{x_1 \in X_1}, (P^{y_2, y_1})_{y_1 \in Y_1}, (Q_{a_1, a_2})_{a_2 \in A_2}$ and $(Q^{b_1, b_2})_{b_2 \in B_2}$ on K with mutually commuting entries, and a unit vector $\eta \in K$ such that

$$\Gamma(x_1y_1, a_2b_2 | x_2y_2, a_1b_1) = \langle P_{x_2, x_1} P^{y_2, y_1} Q_{a_1, a_2} Q^{b_1, b_2} \eta, \eta \rangle$$

for all $x_i \in X_i, y_i \in Y_i, a_i \in A_i, b_i \in B_i, i = 1, 2$. However, we have since discovered a gap in the proof of said lemma resulting from the incorrect use of associativity of the commuting operator system tensor product (see Section 6).

In generalizing to the case of what we call SQNS correlations, as the natural generalization of NS operator matrices are strongly stochastic operator matrices if we were to follow the path laid out in [20, Definition 5.6 (i)] one would expect that we say $\Gamma \in \mathcal{Q}_{\text{sqc}}$ if there exists a Hilbert space H , dilatable strongly stochastic operator matrices $P = (P^{x_1x'_1, y_1y'_1})_{x_2x'_2, y_2y'_2}$ and

$Q = (Q_{a_1 a'_1, b_1 b'_1}^{a_2 a'_2, b_2 b'_2})_{a_1 a'_1, b_1 b'_1, a_2 a'_2, b_2 b'_2}$ with mutually commuting entries acting on H , and a unit vector $\xi \in H$ such that $\Gamma = \Gamma_{P, Q, \xi}$. However, in order to get the decomposition of strongly stochastic operator matrices that we will need for the results in Section 6 due to the failure of [20, Lemma 5.8] (and its extension) we must instead require it in the definition (as seen in Definition 3.7 (i)). Furthermore, we will need to amend the definition for $\Gamma \in \mathcal{C}_{\text{sqc}}$, which we do below. We point out that working with Definition 3.9 (see below) instead of [20, Definition 5.6(i)] recovers all subsequent results in [20] which were instead obtained using [20, Lemma 5.8].

Definition 3.9. An SNS correlation $\Gamma \in \mathcal{C}_{\text{sns}}$ is now quantum commuting if there exists a Hilbert space K , PVM's $(P_{x_2, x_1})_{x_1 \in X_1}, (P^{y_2, y_1})_{y_1 \in Y_1}, (Q_{a_1, a_2})_{a_2 \in A_2}$ and $(Q^{b_1, b_2})_{b_2 \in B_2}$ on K with mutually commuting entries, and a unit vector $\eta \in K$ such that

$$\Gamma(x_1 y_1, a_2 b_2 | x_2 y_2, a_1 b_1) = \langle P_{x_2, x_1} P^{y_2, y_1} Q_{a_1, a_2} Q^{b_1, b_2} \eta, \eta \rangle$$

for all $x_i \in X_i, y_i \in Y_i, a_i \in A_i, b_i \in B_i, i = 1, 2$.

Proposition 3.10. Let $t \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$ and p an SNS correlation. Then $p \in \mathcal{C}_{\text{st}}$ if and only if $\Gamma_p \in \mathcal{Q}_{\text{st}}$.

Proof. In the case when $t = \text{ns}$, this follows directly by Remark 3.4. For $t = \text{loc}, \text{q}, \text{qa}$ or qc , this follows from Remark 3.2 and [39, Lemma 7.2]. $\square$

Remark 3.11. Let Γ be a local SQNS correlation over $(X_2 Y_2, A_1 B_1, X_1 Y_1, A_2 B_2)$. If we choose dilations of the matrices M and N from (3.7) with mutually commuting entries, we may write the values of Γ in the form

$$\Gamma(x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2 | x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1) = \langle E_{x_2, x'_2}^{x_1, x'_1} E_{a_1, a'_1}^{a_2, a'_2} F_{y_2, y'_2}^{y_1, y'_1} F_{b_1, b'_1}^{b_2, b'_2} \xi, \xi \rangle$$

where the stochastic operator matrices

$$(E_{x_2, x'_2}^{x_1, x'_1})_{x_2, x'_2, x_1, x'_1}, \quad (E_{a_1, a'_1}^{a_2, a'_2})_{a_1, a'_1, a_2, a'_2}, \quad (F_{y_2, y'_2}^{y_1, y'_1})_{y_2, y'_2, y_1, y'_1}, \quad (F_{b_1, b'_1}^{b_2, b'_2})_{b_1, b'_1, b_2, b'_2}$$

have mutually commuting entries. Using the fact that the tensor product of convex combinations of channels remains a convex combination, along with the arguments from Proposition 3.3 we may conclude that Γ is a convex combination of the form

$$\Gamma = \sum_{j=1}^k \lambda_j \Phi_X^{(j)} \otimes \Phi_Y^{(j)} \otimes \Phi_A^{(j)} \otimes \Phi_B^{(j)}, \quad (21)$$

where $\Phi_X^{(j)} : M_{X_2} \rightarrow M_{X_1}, \Phi_Y^{(j)} : M_{Y_2} \rightarrow M_{Y_1}, \Phi_A^{(j)} : M_{A_1} \rightarrow M_{A_2}$, and $\Phi_B^{(j)} : M_{B_1} \rightarrow M_{B_2}$ are quantum channels, $j = 1, \dots, k$. It is also easy to verify that any SQNS correlation of the form (21) is in $\mathcal{Q}_{\text{loc}}$.

Remark 3.12. One may easily see that the operator matrices E, F arising from Definition 3.7 (i) of any $\Gamma \in \mathcal{Q}_{\text{sqc}}$ will be dilatable; as discussed in Remark 3.8

it is unknown if all mutually commuting dilatable strongly stochastic operator matrices can be made jointly dilatable.

In the remainder of this section, assume that $X_1 = X_2 := X$, $Y_1 = Y_2 := Y$, $A_1 = A_2 := A$, and $B_1 = B_2 := B$. Furthermore, assume that $X = A$ and $Y = B$. An SQNS correlation Γ will be called an *SQNS bicorrelation* if Γ is unital and Γ^* is an SQNS correlation. The collection of all SQNS correlations will be denoted $\mathcal{Q}_{\text{sns}}^{\text{bi}}$. An SQNS bicorrelation Γ over (XY, XY, XY, XY) is called *quantum commuting* if there exists a Hilbert space H , strongly bistochastic operator matrices

$$\begin{aligned} E_X &= (E_{x_2 x'_2, x_1 x'_1})_{x_2 x'_2, x_1 x'_1}, & E_Y &= (E_{y_2 y'_2, y_1 y'_1})_{y_2 y'_2, y_1 y'_1}, \\ F_A &= (F_{a_1 a'_1, a_2 a'_2})_{a_1 a'_1, a_2 a'_2}, & F_B &= (F_{b_1 b'_1, b_2 b'_2})_{b_1 b'_1, b_2 b'_2}, \end{aligned}$$

with mutually commuting entries acting on H , and a unit vector $\xi \in H$ such that $\Gamma = \Gamma_{E, F, \xi}$ where $E = E_X \cdot E_Y$, $F = F_A \cdot F_B$; we denote this class by $\mathcal{Q}_{\text{sqc}}^{\text{bi}}$. The classes of quantum SQNS bicorrelations (denoted $\mathcal{Q}_{\text{sq}}^{\text{bi}}$), approximately quantum SQNS bicorrelations (denoted $\mathcal{Q}_{\text{sqa}}^{\text{bi}}$) and local SQNS bicorrelations (denoted $\mathcal{Q}_{\text{sloc}}^{\text{bi}}$) are defined similarly to their SQNS correlation counterparts, with dilatable strongly bistochastic operator matrices of the appropriate type in place of the strongly stochastic operator matrices of said type. In the sequel, we let $\mathcal{C}_{\text{st}}^{\text{bi}}$ denote the class of all SQNS bicorrelations of type t , as defined in [20, Section 6] and Remark 3.8.

Remark 3.13. Arguing as in Remark 3.11, we may identify $\mathcal{Q}_{\text{sloc}}^{\text{bi}}$ with SQNS correlations Γ of the form

$$\Gamma = \sum_{j=1}^k \lambda_j \Phi_X^{(j)} \otimes \Phi_Y^{(j)} \otimes \Phi_A^{(j)} \otimes \Phi_B^{(j)},$$

where $\Phi_X^{(j)} : M_X \rightarrow M_X$, $\Phi_Y^{(j)} : M_Y \rightarrow M_Y$, $\Phi_A^{(j)} : M_A \rightarrow M_A$, $\Phi_B^{(j)} : M_B \rightarrow M_B$ are unital quantum channels, $\lambda_j \geq 0$, $j = 1, \dots, k$ with $\sum_{j=1}^k \lambda_j = 1$.

Remark 3.14. For a correlation type $t \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$, it is clear that $\mathcal{Q}_{\text{st}}^{\text{bi}} \subseteq \mathcal{Q}_{\text{st}}$. Additionally, if $\Gamma \in \mathcal{Q}_{\text{st}}^{\text{bi}}$, then $\Gamma^* \in \mathcal{Q}_{\text{st}}^{\text{bi}}$. Indeed: it is part of the definition when $t = \text{ns}$, and easily verified (using Remark 3.13) when $t = \text{loc}$. The case when $t = \text{qc}, \text{q}$ or qa follow using a modification of the argument given in [7, Remark 5.2].

Proposition 3.15. *Let $t \in \{\text{loc}, \text{q}, \text{qc}, \text{ns}\}$ and p an SNS bicorrelation. Then $p \in \mathcal{C}_{\text{st}}^{\text{bi}}$ if and only if $\Gamma_p \in \mathcal{Q}_{\text{st}}^{\text{bi}}$.*

Proof. This holds essentially by using the same arguments as in Proposition 3.10 and [7, Proposition 5.9]. $\square$

4. Strategy transport

Let X_i, Y_i, A_i, B_i be finite sets, for $i = 1, 2$. Recall that if $\mathcal{E}$ is a QNS correlation over (X_1, Y_1, A_1, B_1) , it acts as a quantum channel $\mathcal{E} : M_{X_1 Y_1} \rightarrow M_{A_1 B_1}$; thus, if Γ is SQNS over $(X_2 Y_2, A_1 B_1, X_1 Y_1, A_2 B_2)$, then $\Gamma[\mathcal{E}] : M_{X_2 Y_2} \rightarrow M_{A_2 B_2}$. The two theorems in this section are crucial to our goal of transferring strategies between quantum games; briefly, they say that if $\mathcal{E}$ is viewed as a strategy of some type for one game, under mild conditions on Γ the simulated channel $\Gamma[\mathcal{E}]$ can be viewed as a strategy of the same type for another quantum game.

Theorem 4.1. *Let Γ be an SQNS correlation over $(X_2 Y_2, A_1 B_1, X_1 Y_1, A_2 B_2)$ and $\mathcal{E}$ be a QNS correlation over (X_1, Y_1, A_1, B_1) . The following hold:*

- (i) $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{ns}}$;
- (ii) if $\Gamma \in \mathcal{Q}_{\text{sqc}}$ and $\mathcal{E} \in \mathcal{Q}_{\text{qc}}$, then $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{qc}}$;
- (iii) if $\Gamma \in \mathcal{Q}_{\text{sqa}}$ and $\mathcal{E} \in \mathcal{Q}_{\text{qa}}$, then $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{qa}}$;
- (iv) if $\Gamma \in \mathcal{Q}_{\text{sq}}$ and $\mathcal{E} \in \mathcal{Q}_{\text{q}}$, then $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{q}}$;
- (v) if $\Gamma \in \mathcal{Q}_{\text{sloc}}$ and $\mathcal{E} \in \mathcal{Q}_{\text{loc}}$, then $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{loc}}$.

Proof. (i) Let C denote the Choi matrix of quantum channel $\Gamma[\mathcal{E}]$. If

$$(\Gamma(x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2 | x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1))$$

is the Choi matrix of Γ and $(\mathcal{E}(a_1 a'_1, b_1 b'_1 | x_1 x'_1, y_1 y'_1))$ is the Choi matrix of $\mathcal{E}$ (where the former matrix ranges over $X_1, X_2, Y_1, Y_2, A_1, A_2, B_1, B_2$ and the latter ranges over X_1, Y_1, A_1 and B_1), as $\Gamma \in \mathcal{Q}_{\text{sns}}$ and $\mathcal{E} \in \mathcal{Q}_{\text{ns}}$ there exists $c_{x_1 x'_1, y_1 y'_1, x_2 x'_2, y_2 y'_2}^{b_2 b'_2, b_1 b'_1}, d_{y_1 y'_1}^{b_1, b'_1} \in \mathbb{C}$ such that

$$\begin{aligned} \sum_{a_2 \in A_2} \Gamma(x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2 | x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1) &= \delta_{a_1, a'_1} c_{x_1 x'_1, y_1 y'_1, x_2 x'_2, y_2 y'_2}^{b_2 b'_2, b_1 b'_1}, \\ \sum_{a_1 \in A_1} \mathcal{E}(a_1 a'_1, b_1 b'_1 | x_1 x'_1, y_1 y'_1) &= \delta_{x_1, x'_1} d_{y_1, y'_1}^{b_1, b'_1}, \end{aligned}$$

for $x_i, x'_i \in X_i, y_i, y'_i \in Y_i, b_i, b'_i \in B_i, i = 1, 2$ (see e.g. [15]). Additionally, we may use (12) to guarantee the existence of $c_{y_1 y'_1, y_2 y'_2}^{b_2 b'_2, b_1 b'_1} \in \mathbb{C}$ such that

$$\sum_{x_1 \in X_1} c_{x_1 x'_1, y_1 y'_1, x_2 x'_2, y_2 y'_2}^{b_2 b'_2, b_1 b'_1} = \delta_{x_2, x'_2} c_{y_1 y'_1, y_2 y'_2}^{b_2 b'_2, b_1 b'_1},$$

for $y_i, y'_i \in Y_i, b_i, b'_i \in B_i, i = 1, 2$. For $y_2, y'_2 \in Y_2, b_2, b'_2 \in B_2$, set

$$\tilde{c}_{y_2, y'_2}^{b_2, b'_2} := \sum_{y_1, y'_1 \in Y_1} \sum_{b_1, b'_1 \in B_1} c_{y_1 y'_1, y_2 y'_2}^{b_2 b'_2, b_1 b'_1} d_{y_1, y'_1}^{b_1, b'_1}.$$

If we fix $y_2, y'_2 \in Y_2, b_2, b'_2 \in B_2$, then

$$\begin{aligned}
& \sum_{a_2} \sum_{\substack{x_1 x'_1, y_1 y'_1, \\ a_1 a'_1, b_1 b'_1}} \Gamma(x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2 | x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1) \mathcal{E}(a_1 a'_1, b_1 b'_1 | x_1 x'_1, y_1 y'_1) \\
&= \sum_{\substack{x_1 x'_1, y_1 y'_1 \\ a_1 a'_1, b_1 b'_1}} \delta_{a_1, a'_1} c_{x_1 x'_1, y_1 y'_1, x_2 x'_2, y_2 y'_2}^{b_2 b'_2, b_1 b'_1} \mathcal{E}(a_1 a'_1, b_1 b'_1 | x_1 x'_1, y_1 y'_1) \\
&= \sum_{a_1} \sum_{\substack{x_1 x'_1, y_1 y'_1 \\ b_1 b'_1}} c_{x_1 x'_1, y_1 y'_1, x_2 x'_2, y_2 y'_2}^{b_2 b'_2, b_1 b'_1} \mathcal{E}(a_1 a'_1, b_1 b'_1 | x_1 x'_1, y_1 y'_1) \\
&= \sum_{\substack{x_1 x'_1, y_1 y'_1, \\ b_1 b'_1}} \delta_{x_1, x'_1} c_{x_1 x'_1, y_1 y'_1, x_2 x'_2, y_2 y'_2}^{b_2 b'_2, b_1 b'_1} d_{y_1, y'_1}^{b_1, b'_1} \\
&= \sum_{x_1} \sum_{\substack{y_1 y'_1, \\ b_1 b'_1}} c_{x_1 x_1, y_1 y'_1, x_2 x'_2, y_2 y'_2}^{b_2 b'_2, b_1 b'_1} d_{y_1, y'_1}^{b_1, b'_1} \\
&= \delta_{x_2, x'_2} \left(\sum_{y_1 y'_1, b_1 b'_1} c_{y_1 y'_1, y_2 y'_2}^{b_2 b'_2, b_1 b'_1} d_{y_1, y'_1}^{b_1, b'_1} \right) \\
&= \delta_{x_2, x'_2} \tilde{c}_{y_2, y'_2}^{b_2, b'_2}.
\end{aligned}$$

We may use a similar argument (relying on (13) and (15)) showing the existence of $\tilde{c}_{x_2, x'_2}^{a_2, a'_2} \in \mathbb{C}$ such that

$$\begin{aligned}
& \sum_{b_2} \sum_{\substack{x_1 x'_1, y_1 y'_1 \\ a_1 a'_1, b_1 b'_1}} \Gamma(x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2 | x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1) \mathcal{E}(a_1 a'_1, b_1 b'_1 | x_1 x'_1, y_1 y'_1) \\
&= \delta_{y_2, y'_2} \tilde{c}_{x_2, x'_2}^{a_2, a'_2}
\end{aligned}$$

for $x_2, x'_2 \in X_2, a_2, a'_2 \in A_2$. By (8), the former equality implies $L_{\omega_{XA}}(C) \in \mathcal{L}_{Y_2 B_2}$ for every $\omega_{XA} \in M_{X_2 A_2}$, while the latter equality implies $L_{\omega_{YB}}(C) \in \mathcal{L}_{X_2 A_2}$ for every $\omega_{YB} \in M_{Y_2 B_2}$. Thus, $C \in (\mathcal{L}_{X_2 A_2} \otimes \mathcal{L}_{Y_2 B_2}) \cap M_{X_2 Y_2 A_2 B_2}^+$; by the injectivity of the minimal operator system tensor product, $C \in (\mathcal{L}_{X_2 A_2} \otimes_{\min} \mathcal{L}_{Y_2 B_2})^+$. Argue now as [39, Theorem 6.2] to finish the proof.

(ii) Use Definition 3.7 (i) to obtain a Hilbert space H , stochastic operator matrices

$$\begin{aligned}
P_X &= (P_{x_2 x'_2, x_1 x'_1})_{x_2 x'_2, x_1 x'_1}, \\
P_Y &= (P_{y_2 y'_2, y_1 y'_1})_{y_2 y'_2, y_1 y'_1}, \\
Q_A &= (Q_{a_1 a'_1, a_2 a'_2})_{a_1 a'_1, a_2 a'_2}, \\
Q_B &= (Q_{b_1 b'_1, b_2 b'_2})_{b_1 b'_1, b_2 b'_2},
\end{aligned}$$

acting on K and a unit vector $\xi \in H$ such that $\Gamma = \Gamma_{P,Q,\xi}$. Let K be a Hilbert space, $E = (E_{x_1x'_1, a_1a'_1})_{x_1x'_1, a_1a'_1}$ and $F = (F^{y_1y'_1, b_1b'_1})_{y_1y'_1, b_1b'_1}$ stochastic operator matrices over (X_1, A_1) and (Y_1, B_1) with mutually commuting entries, and $\eta \in K$ a unit vector such that $\mathcal{E} = \mathcal{E}_{E,F,\eta}$. For $x_2, x'_2 \in X_2, a_2, a'_2 \in A_2, y_2, y'_2 \in Y_2$, and $b_2, b'_2 \in B_2$ set

$$\tilde{E}_{x_2x'_2, a_2a'_2} = \sum_{x_1, x'_1 \in X_1} \sum_{a_1, a'_1 \in A_1} P_{x_2x'_2, x_1x'_1} Q_{a_1a'_1, a_2a'_2} \otimes E_{x_1x'_1, a_1a'_1} \quad (22)$$

and

$$\tilde{F}_{y_2y'_2, b_2b'_2} = \sum_{y_1, y'_1 \in Y_1} \sum_{b_1, b'_1 \in B_1} P^{y_2y'_2, y_1y'_1} Q^{b_1b'_1, b_2b'_2} \otimes F^{y_1y'_1, b_1b'_1}. \quad (23)$$

We note that $\tilde{E} = (\tilde{E}_{x_2x'_2, a_2a'_2})_{x_2x'_2, a_2a'_2}$ and $\tilde{F} = (\tilde{F}_{y_2y'_2, b_2b'_2})_{y_2y'_2, b_2b'_2}$ are positive (as the entries of P_X, P_Y, Q_A, Q_B all commute with one another), and claim that $\tilde{E}$ (resp. $\tilde{F}$) is a stochastic operator matrix over (X_2, A_2) (resp. over (Y_2, B_2)). Indeed: as P_X, Q_A and E are all stochastic operator matrices, we have

$$\begin{aligned} \text{Tr}_{A_2}(\tilde{E}) &= \sum_{a_2} \sum_{x_2, x'_2} \sum_{x_1, x'_1} \sum_{a_1, a'_1} P_{x_2x'_2, x_1x'_1} Q_{a_1a'_1, a_2a'_2} \otimes E_{x_1x'_1, a_1a'_1} \\ &= \sum_{x_2, x'_2} \sum_{x_1, x'_1} \sum_{a_1, a'_1} P_{x_2x'_2, x_1x'_1} (\delta_{a_1, a'_1} I_H) \otimes E_{x_1x'_1, a_1a'_1} \\ &= \sum_{x_2, x'_2} \sum_{x_1, x'_1} \sum_{a_1} P_{x_2x'_2, x_1x'_1} \otimes E_{x_1x'_1, a_1a_1} \\ &= \sum_{x_2, x'_2} \sum_{x_1, x'_1} P_{x_2x'_2, x_1x'_1} \otimes (\delta_{x_1, x'_1} I_K) \\ &= \sum_{x_2, x'_2} \sum_{x_1} P_{x_2x'_2, x_1x_1} \otimes I_K \\ &= \sum_{x_2, x'_2} (\delta_{x_2, x'_2} I_H) \otimes I_K \\ &= I_{X_2} \otimes I_H \otimes I_K. \end{aligned}$$

Thus, $\tilde{E} \in M_{X_2A_2} \otimes \mathcal{B}(H \otimes K)$ is stochastic, as claimed. By symmetry, $\tilde{F} \in M_{Y_2B_2} \otimes \mathcal{B}(H \otimes K)$ is stochastic. We also note that

$$\tilde{E}_{x_2x'_2, a_2a'_2} \tilde{F}_{y_2y'_2, b_2b'_2} = \tilde{F}_{y_2y'_2, b_2b'_2} \tilde{E}_{x_2x'_2, a_2a'_2},$$

for all $x_2, x'_2 \in X_2, y_2, y'_2 \in Y_2, a_2, a'_2 \in A_2, b_2, b'_2 \in B_2$. Using (8), we see

$$\begin{aligned} &\Gamma[\mathcal{E}](a_2a'_2, b_2b'_2 | x_2x'_2, y_2y'_2) \\ &= \sum \Gamma(x_1x'_1, y_1y'_1, a_2a'_2, b_2b'_2 | x_2x'_2, y_2y'_2, a_1a'_1, b_1b'_1) \mathcal{E}(a_1a'_1, b_1b'_1 | x_1x'_1, y_1y'_1) \\ &= \sum \langle P_{x_2x'_2, x_1x'_1} P^{y_2y'_2, y_1y'_1} Q_{a_1a'_1, a_2a'_2} Q^{b_1b'_1, b_2b'_2} \xi, \xi \rangle \langle E_{x_1x'_1, a_1a'_1} F^{y_1y'_1, b_1b'_1} \eta, \eta \rangle \\ &= \langle \tilde{E}_{x_2x'_2, a_2a'_2} \tilde{F}_{y_2y'_2, b_2b'_2} (\xi \otimes \eta), \xi \otimes \eta \rangle, \end{aligned}$$

for all $x_2, x'_2 \in X_2, y_2, y'_2 \in Y_2, a_2, a'_2 \in A_2, b_2, b'_2 \in B_2$, where the latter sums range over all of X_1, Y_1, A_1 , and B_1 . Thus, $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{qc}}$.

(iii) This is a direct consequence of (iv).

(iv) Let $M = (M_{x_2 x'_2, a_1 a'_1}^{x_1 x'_1, a_2 a'_2})_{x_2 x'_2, a_1 a'_1, x_1 x'_1, a_2 a'_2}$ and $N = (N_{y_2 y'_2, b_1 b'_1}^{y_1 y'_1, b_2 b'_2})_{y_2 y'_2, b_1 b'_1, y_1 y'_1, b_2 b'_2}$ be quantum dilatable strongly stochastic operator matrices acting on H and K , respectively, and $\xi \in H \otimes K$ be a unit vector for which $\Gamma = \Gamma_{M \otimes N, \xi}$. Let $E = (E_{x_1 x'_1, a_1 a'_1})_{x_1 x'_1, a_1 a'_1}$ and $F = (F_{y_1 y'_1, b_1 b'_1})_{y_1 y'_1, b_1 b'_1}$ be finite-dimensionally acting stochastic operator matrices with η a unit vector such that $\mathcal{E} = \mathcal{E}_{E \otimes F, \eta}$. For $x_2, x'_2 \in X_2, a_2, a'_2 \in A_2, y_2, y'_2 \in Y_2$, and $b_2, b'_2 \in B_2$ set

$$\tilde{E}_{x_2 x'_2, a_2 a'_2} = \sum_{x_1, x'_1 \in X_1} \sum_{a_1, a'_1 \in A_1} M_{x_2 x'_2, a_1 a'_1}^{x_1 x'_1, a_2 a'_2} \otimes E_{x_1 x'_1, a_1 a'_1},$$

and

$$\tilde{F}_{y_2 y'_2, b_2 b'_2} = \sum_{y_1, y'_1 \in Y_1} \sum_{b_1, b'_1 \in B_1} N_{y_2 y'_2, b_1 b'_1}^{y_1 y'_1, b_2 b'_2} \otimes F_{y_1 y'_1, b_1 b'_1};$$

using an almost identical argument as in the proof of (ii), it is easy to verify that $\tilde{E} = (\tilde{E}_{x_2 x'_2, a_2 a'_2})_{x_2 x'_2, a_2 a'_2}$ and $\tilde{F} = (\tilde{F}_{y_2 y'_2, b_2 b'_2})_{y_2 y'_2, b_2 b'_2}$ are finite-dimensionally acting stochastic operator matrices such that

$$\Gamma[\mathcal{E}](a_2 a'_2, b_2 b'_2 | x_2 x'_2, y_2 y'_2) = \langle (\tilde{E}_{x_2 x'_2, a_2 a'_2} \otimes \tilde{F}_{y_2 y'_2, b_2 b'_2})(\xi \otimes \eta), \xi \otimes \eta \rangle.$$

Thus, $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{q}}$.

(v) If $\Phi_X : M_{X_2} \rightarrow M_{X_1}, \Phi_Y : M_{Y_2} \rightarrow M_{Y_1}, \Phi_A : M_{A_1} \rightarrow M_{A_2}$ and $\Phi_B : M_{B_1} \rightarrow M_{B_2}$ are quantum channels, it is easily verified (see [20]) that

$$(\Phi_X \otimes \Phi_Y \otimes \Phi_A \otimes \Phi_B)[\mathcal{E} \otimes \mathcal{F}] = (\Phi_X \otimes \Phi_A)[\mathcal{E}] \otimes (\Phi_Y \otimes \Phi_B)[\mathcal{F}]$$

for all quantum channels $\mathcal{E} : M_{X_1} \rightarrow M_{A_1}$ and $\mathcal{F} : M_{Y_1} \rightarrow M_{B_1}$. Using Remark 3.11, we may conclude that $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{loc}}$ as claimed. $\square$

Fix finite sets X and Y , and let $A = X, B = Y$.

Theorem 4.2. *Let Γ be an SQNS bicomrelation over (XY, XY, XY, XY) and $\mathcal{E}$ be a QNS bicomrelation over (X, Y, X, Y) . The following hold:*

- (i) $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{ns}}^{\text{bi}}$.
- (ii) if $\Gamma \in \mathcal{Q}_{\text{sqc}}^{\text{bi}}$ and $\mathcal{E} \in \mathcal{Q}_{\text{qc}}^{\text{bi}}$, then $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{qc}}^{\text{bi}}$.
- (iii) if $\Gamma \in \mathcal{Q}_{\text{sqa}}^{\text{bi}}$ and $\mathcal{E} \in \mathcal{Q}_{\text{qa}}^{\text{bi}}$, then $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{qa}}^{\text{bi}}$.
- (iv) if $\Gamma \in \mathcal{Q}_{\text{sq}}^{\text{bi}}$ and $\mathcal{E} \in \mathcal{Q}_{\text{q}}^{\text{bi}}$, then $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{q}}^{\text{bi}}$.
- (v) if $\Gamma \in \mathcal{Q}_{\text{sloc}}^{\text{bi}}$ and $\mathcal{E} \in \mathcal{Q}_{\text{loc}}^{\text{bi}}$, then $\Gamma[\mathcal{E}] \in \mathcal{Q}_{\text{loc}}^{\text{bi}}$.

Proof. (i) The claim follows as in Theorem 4.1, once we note that

$$\begin{aligned}
 & \Gamma[\mathcal{E}]^*(x_2x'_2, y_2y'_2 | a_2a'_2, b_2b'_2) \\
 &= \sum_{x_1x'_1, y_1y'_1} \sum_{a_1a'_1, b_1b'_1} \Gamma(x_1x'_1, y_1y'_1, a_2a'_2, b_2b'_2 | x_2x'_2, y_2y'_2, a_1a'_1, b_1b'_1) \mathcal{E}(a_1a'_1, b_1b'_1 | x_1x'_1, y_1y'_1) \\
 &= \sum_{a_1a'_1, b_1b'_1} \sum_{x_1x'_1, y_1y'_1} \Gamma^*(x_2x'_2, y_2y'_2, a_1a'_1, b_1b'_1 | x_1x'_1, y_1y'_1, a_2a'_2, b_2b'_2) \mathcal{E}^*(x_1x'_1, y_1y'_1 | a_1a'_1, b_1b'_1) \\
 &= \Gamma^*[\mathcal{E}^*](x_2x'_2, y_2y'_2 | a_2a'_2, b_2b'_2)
 \end{aligned}$$

for all $x_2, x'_2 \in X, y_2, y'_2 \in Y, a_2, a'_2 \in A, b_2, b'_2 \in B$. This means $\Gamma^*[\mathcal{E}^*] = \Gamma[\mathcal{E}]^*$.

(ii)-(v) follow as in the proof of Theorem 4.1 (ii)-(v), once we notice that the stochastic operator matrices defined there are bistochastic. $\square$

5. Perfect strategies for various quantum games

For an arbitrary Hilbert space H , we write $\overline{H}$ for the Banach space dual of H ; by the Riesz Representation Theorem, there exists a conjugate linear isometry $\partial : H \rightarrow \overline{H}$, such that $\partial(\xi)(\eta) = \langle \eta, \xi \rangle$, $\xi, \eta \in H$. In what follows, we will write $\overline{\mathbb{C}}^X = \overline{\mathbb{C}}^X$ for any finite set X . Set $\overline{\xi} = \partial(\xi)$, for $\xi \in H$. Given a linear operator $A : H \rightarrow K$, (where H, K are Hilbert spaces) let $\overline{A} : \overline{K} \rightarrow \overline{H}$ be its (Banach space) dual operator. Note that $\overline{A}(\overline{\xi}) = \overline{A^*\xi}$, for $\xi \in K$.

If H, K are finite dimensional, and we take vectors $\xi \in H, \eta \in K$ let $\eta\xi^* : H \rightarrow K$ be the rank one operator given by $(\eta\xi^*)(\xi') = \langle \xi', \xi \rangle \eta$. Let $\theta : \overline{H} \otimes K \rightarrow \mathcal{L}(H, K)$ be the linear isomorphism given by

$$\theta(\overline{\xi} \otimes \eta) = \eta\xi^*, \quad \xi \in H, \eta \in K.$$

If $\mathcal{V} \subseteq \overline{H} \otimes K$ is some subspace, we let $\overline{\mathcal{V}} := \theta(\mathcal{V}) \subseteq \mathcal{L}(H, K)$ be the corresponding subspace of linear operators.

Recall (see [21, Definition 4.1]) that if X, Y are finite sets, a quantum hypergraph over (X, Y) is any subspace $\mathcal{U} \subseteq \overline{\mathbb{C}}^X \otimes \mathbb{C}^Y$. For a classical hypergraph $E \subseteq X \times Y$, let

$$\mathcal{U}_E = \text{span}\{\overline{e}_x \otimes e_y : (x, y) \in E\}$$

be viewed as a quantum hypergraph over (X, Y) . Furthermore, following the notation established in [21] fix finite sets $X_i, Y_i, i = 1, 2$. Let $\mathcal{U}_1 \subseteq \overline{\mathbb{C}}^{X_1} \otimes \overline{\mathbb{C}}^{Y_1}$, $\mathcal{U}_2 \subseteq \overline{\mathbb{C}}^{X_2} \otimes \mathbb{C}^{Y_2}$ be quantum hypergraphs and set

$$\mathcal{U}_1 \Leftrightarrow \mathcal{U}_2 := (\mathcal{U}_1 \otimes \mathcal{U}_2) + (\mathcal{U}_1^\perp \otimes \mathcal{U}_2^\perp).$$

If

$$\sigma : \mathbb{C}^{X_1} \otimes \overline{\mathbb{C}}^{Y_1} \otimes \overline{\mathbb{C}}^{X_2} \otimes \mathbb{C}^{Y_2} \rightarrow \overline{\mathbb{C}}^{X_2} \otimes \overline{\mathbb{C}}^{Y_1} \otimes \mathbb{C}^{X_1} \otimes \mathbb{C}^{Y_2}$$

is the flip on the 1st and 3rd tensor legs, then we set

$$\mathcal{U}_1 \leftrightarrow \mathcal{U}_2 := \sigma(\mathcal{U}_1 \Leftrightarrow \mathcal{U}_2).$$

Given a quantum hypergraph $\mathcal{U} \subseteq \overline{\mathbb{C}}^X \otimes \mathbb{C}^Y$, we let

$$\mathcal{Q}(\mathcal{U}) := \left\{ \Phi : M_X \rightarrow M_Y : \text{a quantum channel with } \mathcal{K}_\Phi \subseteq \tilde{\mathcal{U}} \right\},$$

where $\mathcal{K}_\Phi$ is the Kraus space corresponding to channel Φ . For the next collection of results, we will need the following definition.

Definition 5.1 ([21]). Let $t \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$, with $\mathcal{U}_1 \subseteq \mathbb{C}^{X_1} \otimes \overline{\mathbb{C}}^{Y_1}$ and $\mathcal{U}_2 \subseteq \overline{\mathbb{C}}^{X_2} \otimes \mathbb{C}^{Y_2}$ quantum hypergraphs. We say that $\mathcal{U}_1$ is t -homomorphic to $\mathcal{U}_2$, written $\mathcal{U}_1 \rightarrow_t \mathcal{U}_2$ if there exists a quantum channel $\Phi : M_{X_2 Y_1} \rightarrow M_{X_1 Y_2}$ with $\Phi \in \mathcal{Q}_t$ such that $\Phi \in \mathcal{Q}(\mathcal{U}_1 \leftrightarrow \mathcal{U}_2)$.

5.1. Perfect strategies for quantum implication games. Let X, Y, A, B be finite sets. If $P \in M_{XY}, Q \in M_{AB}$ are projections, the quantum implication game (see [39]) $P \Rightarrow Q$ is the quantum non-local game $\varphi_{P \Rightarrow Q} : \mathcal{P}_{XY} \rightarrow \mathcal{P}_{AB}$ given by

$$\varphi_{P \Rightarrow Q}(\tilde{P}) = \begin{cases} Q, & \text{if } 0 \neq \tilde{P} \leq P, \\ 0, & \text{if } \tilde{P} = 0, \\ I, & \text{otherwise.} \end{cases}$$

In other words, a quantum implication game is a quantum non-local game which states that if a player is given an input state supported P , their output should be a state supported on Q . A QNS correlation $\Phi : M_{XY} \rightarrow M_{AB}$ is called a *perfect strategy* for $\varphi_{P \Rightarrow Q}$ if $\langle \Phi(P), Q^\perp \rangle = 0$. Equivalently, Φ is perfect if

$$\omega \in M_{XY}^+ \text{ and } \omega = P\omega P \Rightarrow \Phi(\omega) = Q\Phi(\omega)Q.$$

As both P and Q are finite-rank projections, we may find orthonormal basis $\{\xi_i\}_{i=1}^n \subseteq \mathbb{C}^{XY}$ (resp. $\{\gamma_\ell\}_{\ell=1}^m \subseteq \mathbb{C}^{AB}$) for $\text{rng}(P)$ (resp. $\text{rng}(Q)$). We may then associate to any quantum implication game the subspace

$$\mathcal{U}_{P,Q} := \text{span}\{\bar{\xi}_i \otimes \gamma_\ell : i \in [n], \ell \in [m]\},$$

considered as a quantum hypergraph over (XY, AB) . Note that if $S \in \tilde{\mathcal{U}}_{P,Q} \subseteq \mathcal{L}(\mathbb{C}^{XY}, \mathbb{C}^{AB})$, then $S = \sum_{j=1}^t \lambda_j \gamma_j \xi_j^*$ where $\xi_j \in \text{rng}(P), \gamma_j \in \text{rng}(Q)$ and $\lambda_j \in \mathbb{C}$ for $j = 1, \dots, t$.

If $\mathfrak{G} \subseteq \text{QC}(M_{XY}, M_{AB})$ is a convex subset of quantum channels from M_{XY} to M_{AB} , we let

$$\omega_{\mathfrak{G}}(P, Q) = \sup_{\Phi \in \mathfrak{G}} \text{Tr}(\Phi(P)Q) \quad (24)$$

be the $\mathfrak{G}$ -value of the quantum implication game $P \Rightarrow Q$. Specifically, if $\mathfrak{G} = \mathcal{Q}_t$ where $t \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$ we set $\omega_t(P, Q) = \omega_{\mathcal{Q}_t}(P, Q)$; one may easily check that $\omega_t(P, Q) = 1$ if and only if there exists a perfect t -strategy Φ for $P \Rightarrow Q$.

Lemma 5.2. *A QNS correlation $\Phi : M_{XY} \rightarrow M_{AB}$ is a perfect strategy for $\varphi_{P \rightarrow Q}$ if and only if $\mathcal{K}_\Phi \subseteq \tilde{\mathcal{U}}_{P,Q}$.*

Proof. First, assume that Φ is a perfect strategy for the implication game $\varphi_{P \rightarrow Q}$. Let $\Phi(S) = \sum_{j=1}^t M_j S M_j^*$ be a Kraus decomposition for Φ ; as Φ is perfect,

$$\begin{aligned} 0 &= \text{Tr}(\Phi(P)Q^\perp) \\ &= \sum_{j=1}^t \text{Tr}(M_j P M_j^* Q^\perp) \\ &= \sum_{j=1}^t \text{Tr}(Q^\perp M_j P^2 M_j^* Q^\perp) \\ &= \sum_{j=1}^t \text{Tr}((Q^\perp M_j P)(Q^\perp M_j P)^*), \end{aligned}$$

which implies $Q^\perp M_j P = 0$ for each $j = 1, \dots, t$. Taking adjoints, we also have that $P M_j^* Q^\perp = 0$ for each $j = 1, \dots, t$. If we then fix $1 \leq j \leq t$ and pick $\beta \in \text{rng}(P), \alpha \in \text{rng}(Q^\perp)$, we see

$$\langle M_j \beta, \alpha \rangle = \langle M_j P \beta, Q^\perp \alpha \rangle = \langle Q^\perp M_j P \beta, \alpha \rangle = 0.$$

As $\alpha \in \text{rng}(Q^\perp)$ was arbitrary, this implies $M_j \beta \in \text{rng}(Q)$ for $\beta \in \text{rng}(P)$. A similar argument using the adjoint relation $P M_j^* Q^\perp = 0$ implies that $M_j^* \alpha \in \text{rng}(P^\perp)$ for each $\alpha \in \text{rng}(Q^\perp)$. Thus, $M_j \in \tilde{\mathcal{U}}_{P,Q}$. Unfixing our choice of $1 \leq j \leq t$, we see $\mathcal{K}_\Phi \subseteq \tilde{\mathcal{U}}_{P,Q}$.

Now assume $\mathcal{K}_\Phi \subseteq \tilde{\mathcal{U}}_{P,Q}$. By the comment before the lemma statement, one may easily see that for any $M \in \mathcal{K}_\Phi$ we have $Q^\perp M P = 0$. If $M_1, \dots, M_t$ are Kraus operators for Φ , reversing the steps in the previous paragraph we have that

$$0 = \text{Tr}(\Phi(P)Q^\perp) = \langle \Phi(P), Q^\perp \rangle,$$

which shows that Φ is a perfect strategy for the game $\varphi_{P \rightarrow Q}$. $\square$

Theorem 5.3. *Let $X_i, Y_i, A_i, B_i, i = 1, 2$ be finite sets, $P_i \in M_{X_i Y_i}, Q_i \in M_{A_i B_i}, i = 1, 2$ projections, and $t \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$. If $\bar{\mathcal{U}}_{P_1, Q_1} \rightarrow_{\text{st}} \mathcal{U}_{P_2, Q_2}$ via Γ , and if $\mathcal{E} : M_{X_1 Y_1} \rightarrow M_{A_1 B_1}$ is a perfect t -strategy for $\varphi_{P_1 \rightarrow Q_1}$, then $\Gamma[\mathcal{E}]$ is a perfect t -strategy for $\varphi_{P_2 \rightarrow Q_2}$.*

Proof. Suppose that $\Gamma(T) = \sum_{p=1}^t N_p T N_p^*$ and $\mathcal{E}(S) = \sum_{r=1}^s M_r S M_r^*$. By Lemma

5.2, as $\mathcal{E}$ is a perfect strategy for $\varphi_{P_1 \rightarrow Q_1}$ then $M_r \in \tilde{\mathcal{U}}_{P_1, Q_1}$ for each $r = 1, \dots, s$. By construction (see [21]), the Kraus operators of $\Gamma[\mathcal{E}]$ are given by operators $N_p[M_r] : \mathbb{C}^{X_2 Y_2} \rightarrow \mathbb{C}^{A_2 B_2}$, dependent on N_p and M_r for each $p = 1, \dots, t, r = 1, \dots, s$. Furthermore, as $\bar{\mathcal{U}}_{P_1, Q_1} \rightarrow \mathcal{U}_{P_2, Q_2}$ via Γ by [21, Theorem 5.5] we know that $N_p[M_r] \in \tilde{\mathcal{U}}_{P_2, Q_2}$ for each $p = 1, \dots, t, r = 1, \dots, s$. This means $\mathcal{K}_{\Gamma[\mathcal{E}]} \subseteq$

$\tilde{\mathcal{U}}_{P_2, Q_2}$; applying Lemma 5.2 once again, we see that $\Gamma[\mathcal{E}]$ is a perfect strategy for $\varphi_{P_2 \rightarrow Q_2}$. Furthermore, by Theorem 4.1 we have that $\Gamma[\mathcal{E}] \in \mathcal{Q}_t$, whenever $\Gamma \in \mathcal{Q}_{st}$ with $\mathcal{E} \in \mathcal{Q}_t$. $\square$

5.2. Classical-to-quantum non-local games. In this subsection, we will consider a way to transfer strategies from classical non-local games to quantum ones, and how a perfect strategy remains perfect under this transfer even when both games are not quantum. For a fixed $t \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$ to make sense of how we can take a classical strategy $\mathcal{E} \in \mathcal{C}_t$ and “apply” it to a legitimately quantum game, the most natural way is to first use [39, Remark 8.1] to consider $\mathcal{E} \in \mathcal{Q}_t$, and then use $\Gamma \in \mathcal{Q}_{st}$ as defined in Definition 3.7 to construct a quantum strategy $\Gamma[\mathcal{E}]$ via Theorem 4.1.

Lemma 5.4. *Let X_1, Y_1, A_1, B_1 be finite sets, and $E \subseteq X_1 Y_1 \times A_1 B_1$ be a non-local game. Then $\mathcal{E} : \mathcal{D}_{X_1 Y_1} \rightarrow \mathcal{D}_{A_1 B_1}$ is a perfect strategy for the game E if and only if $\tilde{\mathcal{K}}_{\mathcal{E}} \subseteq \tilde{\mathcal{U}}_E$.*

Proof. First, assume $\mathcal{E}$ is a perfect strategy for classical non-local game E . Viewing $\mathcal{E}$ as a quantum channel, by [21, Remark 4.3]

$$\tilde{\mathcal{K}}_{\mathcal{E}} = \text{span}\{e_{a_1} e_{x_1}^* \otimes e_{b_1} e_{y_1}^* : (x_1, y_1, a_1, b_1) \in \text{supp}(\mathcal{E})\}.$$

As $\text{supp}(\mathcal{E}) \subseteq E$, clearly $\tilde{\mathcal{K}}_{\mathcal{E}} \subseteq \tilde{\mathcal{U}}_E$.

Now assume $\tilde{\mathcal{K}}_{\mathcal{E}} \subseteq \tilde{\mathcal{U}}_E$. This means that for any Kraus operator M of $\mathcal{E}$, $M \in \tilde{\mathcal{U}}_E$. Thus, $M = \sum_{j=1}^n \lambda_j e_{a_j} e_{x_j}^* \otimes e_{b_j} e_{y_j}^*$, where $(x_j, y_j, a_j, b_j) \in E$ for $j = 1, \dots, n$. If we assume $\mathcal{E}(S) = \sum_{\ell=1}^m M_{\ell} S M_{\ell}^*$, as

$$\begin{aligned} \mathcal{E}(a_1, b_1 | x_1, y_1) &= \text{Tr}(\mathcal{E}(\epsilon_{x_1 x_1} \otimes \epsilon_{y_1 y_1})(\epsilon_{a_1 a_1} \otimes \epsilon_{b_1 b_1})) \\ &= \sum_{\ell=1}^m \text{Tr}(M_{\ell}(\epsilon_{x_1 x_1} \otimes \epsilon_{y_1 y_1})M_{\ell}^*(\epsilon_{a_1 a_1} \otimes \epsilon_{b_1 b_1})), \end{aligned}$$

where each M_{ℓ} is of the form previously described, it is easy to see that $\mathcal{E}(a_1, b_1 | x_1, y_1) \neq 0$ only if $(x_1, y_1, a_1, b_1) \in E$. Thus, $\text{supp}(\mathcal{E}) \subseteq E$, which means $\mathcal{E}$ is a perfect strategy for E . $\square$

Theorem 5.5. *Let $X_i, Y_i, A_i, B_i, i = 1, 2$ be finite sets, $E \subseteq X_1 Y_1 \times A_1 B_1$ a classical non-local game, $P \in M_{X_2 Y_2}, Q \in M_{A_2 B_2}$ be projections, and let $t \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$. If $\tilde{\mathcal{U}}_E \rightarrow_{st} \mathcal{U}_{P, Q}$ via Γ , and if $\mathcal{E} : \mathcal{D}_{X_1 Y_1} \rightarrow \mathcal{D}_{A_1 B_1}$ is a perfect t -strategy for the game E , then $\Gamma[\mathcal{E}]$ is a perfect t -strategy for quantum non-local game $\varphi_{P \rightarrow Q}$.*

Proof. Assuming the notation and setup as in the proof of Theorem 5.3, by Lemma 5.4 we know that $M_r \in \hat{\mathcal{U}}_E^*$ for each $r = 1, \dots, s$. If the Kraus operators for $\Gamma[\mathcal{E}]$ are given by $N_p[M_r]$ for $p = 1, \dots, t$ and $r = 1, \dots, s$ then by [21, Theorem 5.5] we know that $N_p[M_r] \in \tilde{\mathcal{U}}_{P, Q}$ for each $p = 1, \dots, t$ and $r = 1, \dots, s$. By Lemma 5.2, this means $\Gamma[\mathcal{E}]$ is a perfect strategy for $\varphi_{P \rightarrow Q}$. Furthermore, by

Theorem 4.1 and Proposition 3.10 we have that $\Gamma[\mathcal{E}] \in \mathcal{Q}_t$ whenever $\Gamma \in \mathcal{Q}_{st}$ with $\mathcal{E} \in \mathcal{C}_t$. $\square$

5.3. Quantum graph games. Quantum graphs, and games played on quantum graphs, have generated considerable interest in the last decade. While there are several connected concepts in the literature of what a quantum graph “should be” (see [5, 13, 15]), we will use the concept discussed in [6, 7, 37, 39].

For a finite set X , let $\mathbf{m} : \mathbb{C}^X \otimes \mathbb{C}^X \rightarrow \mathbb{C}$ be the map given by

$$\mathbf{m}(\xi) = \left\langle \xi, \sum_{x \in X} e_x \otimes e_x \right\rangle.$$

Similarly, let $\mathbf{f} : \mathbb{C}^X \otimes \mathbb{C}^X \rightarrow \mathbb{C}^X \otimes \mathbb{C}^X$ be the flip operation, where $\mathbf{f}(\xi \otimes \eta) = \eta \otimes \xi$, for $\xi, \eta \in \mathbb{C}^X$.

Definition 5.6. A quantum graph with vertex set X is a linear subspace $\mathcal{U} \subseteq \mathbb{C}^X \otimes \mathbb{C}^X$ which is skew- in that $\mathbf{m}(\mathcal{U}) = \{0\}$ - and symmetric- in that $\mathbf{f}(\mathcal{U}) = \mathcal{U}$.

For the remainder of this section, for any subspace $\mathcal{U} \subseteq \mathbb{C}^X \otimes \mathbb{C}^X$ we denote by $P_{\mathcal{U}}$ the orthogonal projection onto $\mathcal{U}$. If G is a classical graph on vertex set X , there is a corresponding quantum graph $\mathcal{U}_G$ given by the subspace

$$\mathcal{U}_G := \text{span}\{e_x \otimes e_y : x \sim y\},$$

where we write $P_G = P_{\mathcal{U}_G}$. If $\mathcal{U} \subseteq \mathbb{C}^X \otimes \mathbb{C}^X$ and $\mathcal{V} \subseteq \mathbb{C}^A \otimes \mathbb{C}^A$ are quantum graphs, the quantum graph homomorphism game $\mathcal{U} \rightarrow \mathcal{V}$ is the quantum implication game $\varphi_{P_{\mathcal{U}} \rightarrow P_{\mathcal{V}}}$. As mentioned, QNS correlation $\Gamma : M_{XX} \rightarrow M_{AA}$ is perfect for $\mathcal{U} \rightarrow \mathcal{V}$ if

$$\omega \in M_{XX}^+ \text{ and } P_{\mathcal{U}} \omega P_{\mathcal{U}} \Rightarrow \Phi(\omega) = P_{\mathcal{V}} \Phi(\omega) P_{\mathcal{V}}.$$

In the event that $X = A$, then Φ is a perfect strategy for the quantum graph isomorphism game $\mathcal{U} \cong \mathcal{V}$ if Φ is a bicorrelation, with Φ a perfect strategy for $\mathcal{U} \rightarrow \mathcal{V}$ and Φ^* a perfect strategy for $\mathcal{V} \rightarrow \mathcal{U}$.

Theorem 5.7. Let $\mathcal{U}_i \subseteq \mathbb{C}^{X_i X_i}$, $\mathcal{V}_i \subseteq \mathbb{C}^{A_i A_i}$, $i = 1, 2$ be quantum graphs, with $P_i = P_{\mathcal{U}_i}$, $Q_i = P_{\mathcal{V}_i}$, $i = 1, 2$ their corresponding projections.

- (i) If $\mathcal{E} : M_{X_1 X_1} \rightarrow M_{A_1 A_1}$ is a perfect strategy for the quantum graph homomorphism game $\mathcal{U}_1 \rightarrow_t \mathcal{V}_1$ and $\overline{\mathcal{U}}_{P_1, Q_1} \rightarrow_{st} \mathcal{U}_{P_2, Q_2}$ via Γ , then $\Gamma[\mathcal{E}]$ is a perfect strategy for the quantum graph homomorphism game $\mathcal{U}_2 \rightarrow_t \mathcal{V}_2$;
- (ii) If $X_i = A_i$, $i = 1, 2$ and $\mathcal{E} : M_{X_1 X_1} \rightarrow M_{X_1 X_1}$ is a perfect strategy for the quantum graph isomorphism game $\mathcal{U}_1 \cong_t \mathcal{V}_1$ and $\overline{\mathcal{U}}_{P_1, Q_1} \cong_{st} \mathcal{U}_{P_2, Q_2}$ via Γ , then $\Gamma[\mathcal{E}]$ is a perfect strategy for the quantum graph isomorphism game $\mathcal{U}_2 \cong_t \mathcal{V}_2$.

Proof. (i) This follows as a special consequence of Theorem 5.3.

(ii) Assume the notation and setup as in the proof of Theorem 5.3; by the aforementioned theorem, we already know that $\Gamma[\mathcal{E}]$ is a perfect strategy for the graph homomorphism game $\mathcal{U}_2 \rightarrow \mathcal{V}_2$. To show that $\Gamma[\mathcal{E}]^*$ is a perfect

strategy for $\mathcal{V}_2 \rightarrow \mathcal{U}_2$, first note that $\Gamma[\mathcal{E}]^* = \Gamma^*[\mathcal{E}^*]$ (by construction of simulated channels). Furthermore, by definition of $\mathcal{U}_{P_2, Q_2}$ we may easily verify that $\mathcal{U}_{Q_2, P_2} = \overline{\mathcal{U}_{P_2, Q_2}}$. With the use of [21, Lemma 2.1], and as $\theta^{-1}(N_p[M_r]) \in \mathcal{U}_{P_2, Q_2}$, we have that $\theta^{-1}((N_p[M_r])^*) = \theta^{-1}(N_p^*[M_r^*]) \in \overline{\mathcal{U}_{P_2, Q_2}} = \mathcal{U}_{Q_2, P_2}$, $p = 1, \dots, t$ and $r = 1, \dots, s$. Thus, by Lemma 5.2 once more we see $\mathcal{K}_{\Gamma[\mathcal{E}]^*} \subseteq \mathcal{U}_{Q_2, P_2}$. This means $\Gamma[\mathcal{E}]^*$ is a perfect strategy for $\mathcal{V}_2 \rightarrow \mathcal{U}_2$. That $\Gamma[\mathcal{E}], \Gamma[\mathcal{E}]^*$ are both in $\mathcal{Q}_t$ follows from Theorem 4.2. Together, these show $\mathcal{U}_2 \cong_t \mathcal{V}_2$ via $\Gamma[\mathcal{E}]$, as claimed. $\square$

6. Characterization of SQNS correlations and applications to concurrent games

In the final section, we wish to link strongly quantum no-signalling correlations to the (multi-variate) tensor product of operator systems. In order to do so, we first briefly recall some facts about operator systems and related constructions. If $\mathcal{S}$ and $\mathcal{T}$ are operator systems, we call $\mathcal{S}$ and $\mathcal{T}$ *completely order isomorphic* and write $\mathcal{S} \cong_{\text{c.o.i.}} \mathcal{T}$ if there exists a unital completely positive bijection $\phi : \mathcal{S} \rightarrow \mathcal{T}$ with completely positive inverse. We write $\mathcal{S} \subseteq_{\text{c.o.i.}} \mathcal{T}$ if $\mathcal{S} \subseteq \mathcal{T}$ and the inclusion map $\iota : \mathcal{S} \hookrightarrow \mathcal{T}$ is a complete order isomorphism onto its range. The three main types of operator system tensor products that will be used in the sequel are given as follows:

- (i) the *minimal* operator system tensor product $\mathcal{S} \otimes_{\min} \mathcal{T}$ arises from viewing $\mathcal{S} \otimes \mathcal{T}$ as a subspace of $\mathcal{B}(H \otimes K)$, when we concretely realize $\mathcal{S} \subseteq \mathcal{B}(H)$ and $\mathcal{T} \subseteq \mathcal{B}(K)$ for Hilbert spaces H, K ;
- (ii) the *commuting* tensor product $\mathcal{S} \otimes_c \mathcal{T}$ has the smallest family of matricial cones which make the maps $\phi \cdot \psi$, where $\phi : \mathcal{S} \rightarrow \mathcal{B}(H)$ and $\psi : \mathcal{T} \rightarrow \mathcal{B}(H)$ are completely positive with commuting ranges, completely positive; note that $(\phi \cdot \psi)(x \otimes y) = \phi(x)\psi(y)$, $x \in \mathcal{S}, y \in \mathcal{T}$;
- (iii) the *maximal* operator system tensor product $\mathcal{S} \otimes_{\max} \mathcal{T}$ has matricial cones generated by the elementary tensors of the form $S \otimes T$, where $S \in M_n(\mathcal{S})^+$ and $T \in M_m(\mathcal{T})^+$, $n, m \in \mathbb{N}$.

More details about each tensor product may be found in [27]; the construction of the multivariate tensor product of each type, with explicit descriptions of their matricial cones, may be found in [20, Section 7].

We recall the notion of a coproduct of operator systems: if $\mathcal{S}$ and $\mathcal{T}$ are two operator systems, their coproduct $\mathcal{S} \oplus_1 \mathcal{T}$ is the unique (up to isomorphism) operator system equipped with complete order embeddings $\iota_{\mathcal{S}} : \mathcal{S} \rightarrow \mathcal{S} \oplus_1 \mathcal{T}$ and $\iota_{\mathcal{T}} : \mathcal{T} \rightarrow \mathcal{S} \oplus_1 \mathcal{T}$ which satisfies the following universal property: For each ucp map $\phi : \mathcal{S} \rightarrow \mathcal{R}$ and $\psi : \mathcal{T} \rightarrow \mathcal{R}$, where $\mathcal{R}$ is an operator system, there exists a unique ucp map $\varphi : \mathcal{S} \oplus_1 \mathcal{T} \rightarrow \mathcal{R}$ such that $\varphi(\iota_{\mathcal{S}}(s)) = \phi(s)$ and $\varphi(\iota_{\mathcal{T}}(t)) = \psi(t)$ for every $s \in \mathcal{S}, t \in \mathcal{T}$. For more properties of the coproduct of operator systems, we refer the reader to [25, Section 8].

We also will need to recall the universal operator system for stochastic operator matrices, introduced in [39]. A *ternary ring of operators (TRO)* is a

subspace $\mathcal{V} \subseteq \mathcal{B}(H, K)$ for some Hilbert spaces H and K , such that $ST^*R \in \mathcal{V}$ whenever $S, T, R \in \mathcal{V}$ (see [3, Section 4.3] or [41]). Let X and A be finite sets, and let $\mathcal{V}_{X,A}$ be the universal TRO generated by the entries $v_{a,x}$ of a block operator isometry $V = (v_{a,x})_{a \in A, x \in X}$. That is, $\mathcal{V}_{X,A}$ is the universal TRO with generators $v_{a,x}$, $a \in A, x \in X$, ternary operator $[\cdot, \cdot, \cdot] : \mathcal{V}_{X,A} \times \mathcal{V}_{X,A} \times \mathcal{V}_{X,A} \rightarrow \mathcal{V}_{X,A}$ given by $[u, v, w] = uv^*w$ for $u, v, w \in \mathcal{V}_{X,A}$, and relations

$$\sum_{a \in A} [v_{a'',x''}, v_{a,x}, v_{a,x'}] = \delta_{x,x'} v_{a'',x''}, \quad x, x', x'' \in X, a'' \in A.$$

Let $\mathfrak{C}_{X,A}$ be the unital $*$ -algebra, generated by the set

$$\{v_{a,x}^* v_{a',x'} : x, x' \in X, a, a' \in A\},$$

and $\mathcal{C}_{X,A}$ be the *right C^* -algebra* of $\mathcal{V}_{X,A}$; up to a $*$ -isomorphism, we may view

$$\mathcal{C}_{X,A} = \overline{\text{span}\{\theta(S)^*\theta(T) : S, T \in \mathcal{V}_{X,A}\}}$$

for any faithful ternary representation $\theta : \mathcal{V}_{X,A} \rightarrow \mathcal{B}(H, K)$ (where H, K are Hilbert spaces). Set

$$e_{x,x',a,a'} = v_{a,x}^* v_{a',x'}, \quad x, x' \in X, a, a' \in A$$

where the latter is considered either as an element in $\mathfrak{C}_{X,A}$ or $\mathcal{C}_{X,A}$, depending on the context. The *Brown-Cuntz operator system* (see [39]) inside $\mathcal{C}_{X,A}$ is given by

$$\mathcal{T}_{X,A} := \text{span}\{e_{x,x',a,a'} : x, x' \in X, a, a' \in A\}.$$

We also will consider the space

$$\mathcal{S}_{X,A} := \text{span}\{e_{x,x,a,a} : x \in X, a \in A\},$$

viewed as an operator subsystem inside $\mathcal{T}_{X,A}$. To help distinguish between operator systems $\mathcal{T}_{X,A}$ and $\mathcal{T}_{Y,B}$, we will denote the canonical generators of $\mathcal{T}_{X,A}$ by $e_{x,x',a,a'}, x, x' \in X, a, a' \in A$ and $\mathcal{T}_{Y,B}$ by $f_{y,y',b,b'}, y, y' \in Y, b, b' \in B$.

Similarly, there are canonical operator algebras and operator systems which corresponding to bistochastic operator matrices (first introduced in [7, Section 3]); these will be needed to show an analogous result to Proposition 6.1. Let $\mathcal{V}_X$ be the universal TRO generated by the entries $v_{a,x}$ of a block operator bi-isometry $V = (v_{a,x})_{a \in A, x \in X}$ (see [7]). Let $\mathcal{C}_X$ be the right C^* -algebra of $\mathcal{V}_X$, set $e_{x,x',a,a'} = v_{a,x}^* v_{a,x}$ (where the latter is considered as an element of $\mathcal{C}_X$) and

$$\mathcal{T}_X := \text{span}\{e_{x,x',a,a'} : x, x' \in X, a, a' \in A\}$$

be viewed as an operator system in $\mathcal{C}_X$. Furthermore, we let

$$\mathcal{S}_X := \text{span}\{e_{x,x,a,a} : x \in X, a \in A\}$$

be viewed as an operator subsystem of $\mathcal{T}_X$.

Proposition 6.1. *If $P = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'}$ is a dilatable strongly stochastic operator matrix acting on the Hilbert space H , then there exists a unital completely positive map $\gamma : \mathcal{T}_{X,A} \otimes_c \mathcal{T}_{Y,B} \rightarrow \mathcal{B}(H)$ such that $\gamma(e_{x,x',a,a'} \otimes f_{y,y',b,b'}) = P_{xx',yy'}^{aa',bb'}$. Conversely, if $\gamma : \mathcal{T}_{X,A} \otimes_c \mathcal{T}_{Y,B} \rightarrow \mathcal{B}(H)$ is a unital completely positive map, then $(\gamma(e_{x,x',a,a'} \otimes f_{y,y',b,b'}))_{xx',yy',aa',bb'}$ is a dilatable strongly stochastic operator matrix.*

Proof. Let K be a Hilbert space, $V : H \rightarrow K$ be an isometry, and

$$(E_{xx',aa'})_{x,x' \in X, a,a' \in A}, \quad (F_{yy',bb'})_{y,y' \in Y, b,b' \in B}$$

be mutually commuting stochastic operator matrices on K satisfying (9). Define linear maps $\phi : \mathcal{T}_{X,A} \rightarrow \mathcal{B}(K)$ (resp. $\psi : \mathcal{T}_{Y,B} \rightarrow \mathcal{B}(K)$) by $\phi(e_{x,x',a,a'}) = E_{xx',aa'}$ (resp. $\psi(f_{y,y',b,b'}) = F_{yy',bb'}$). By [39, Theorem 5.2], both ϕ and ψ are unital and completely positive. By the definition of the commuting tensor product of operator spaces, the map $\phi \cdot \psi : \mathcal{T}_{X,A} \otimes_c \mathcal{T}_{Y,B} \rightarrow \mathcal{B}(K)$ given by $(\phi \cdot \psi)(u \otimes v) = \phi(u)\psi(v)$ is (unital and) completely positive as well. Set

$$\gamma(w) = V^*(\phi \cdot \psi)(w)V, \quad w \in \mathcal{T}_{X,A} \otimes_c \mathcal{T}_{Y,B};$$

it is easy to verify that γ is unital and completely positive. Furthermore, we have $\gamma(e_{x,x',a,a'} \otimes f_{y,y',b,b'}) = P_{xx',yy'}^{aa',bb'}$, $x, x' \in X, y, y' \in Y, a, a' \in A, b, b' \in B$.

Conversely, suppose that $\gamma : \mathcal{T}_{X,A} \otimes_c \mathcal{T}_{Y,B} \rightarrow \mathcal{B}(H)$ is a unital completely positive map. By [39, Corollary 5.3] and [27, Theorem 6.4], $\mathcal{T}_{X,A} \otimes_c \mathcal{T}_{Y,B} \subseteq_{c.o.i} \mathcal{C}_{X,A} \otimes_{\max} \mathcal{C}_{Y,B}$. Apply Arveson's Extension Theorem to obtain a completely positive map $\tilde{\gamma} : \mathcal{C}_{X,A} \otimes_{\max} \mathcal{C}_{Y,B} \rightarrow \mathcal{B}(H)$ extending γ . If we then apply Stinespring's Theorem, we may write

$$\tilde{\gamma}(w) = V^*\pi(w)V, \quad w \in \mathcal{C}_{X,A} \otimes_{\max} \mathcal{C}_{Y,B},$$

where $\pi : \mathcal{C}_{X,A} \otimes_{\max} \mathcal{C}_{Y,B} \rightarrow \mathcal{B}(K)$ is a $*$ -representation on some Hilbert space K , and $V : H \rightarrow K$ is an isometry. Set $E_{xx',aa'} = \pi(e_{x,x',a,a'} \otimes 1)$, $F_{yy',bb'} = \pi(1 \otimes f_{y,y',b,b'})$ for $x, x' \in X, a, a' \in A, y, y' \in Y$ and $b, b' \in B$. Doing so gives us a dilatable representation of the matrix $(\gamma(e_{x,x',a,a'} \otimes f_{y,y',b,b'}))_{xx',yy',aa',bb'}$. $\square$

Remark 6.2. By Remark 3.2 and [20, Remark 5.3], there exist strongly stochastic operator matrices which are not dilatable.

Proposition 6.3. *If $P = (P_{xx',yy'}^{aa',bb'})_{xx',yy',aa',bb'}$ is a dilatable strongly bistochastic operator matrix acting on the Hilbert space H , then there exists a unital completely positive map $\gamma : \mathcal{T}_X \otimes_c \mathcal{T}_Y \rightarrow \mathcal{B}(H)$ such that $\gamma(e_{x,x',a,a'} \otimes f_{y,y',b,b'}) = P_{xx',yy'}^{aa',bb'}$. Conversely, if $\gamma : \mathcal{T}_X \otimes_c \mathcal{T}_Y \rightarrow \mathcal{B}(H)$ is a unital completely positive map, then $(\gamma(e_{x,x',a,a'} \otimes f_{y,y',b,b'}))_{xx',yy',aa',bb'}$ is a dilatable strongly bistochastic operator matrix.*

Proof. Using [7, Theorem 3.4] and [27, Theorem 6.4], argue exactly as in Proposition 6.1 but with bistochastic operator matrices in place of stochastic operator matrices and $\mathcal{T}_X$ (resp. $\mathcal{T}_Y$) in place of $\mathcal{T}_{X,A}$ (resp. $\mathcal{T}_{Y,B}$). $\square$

6.1. Representations of SQNS correlations via operator systems. Let $\mathcal{S}$ be an operator system. Recall that the *universal C^* -cover* of $\mathcal{S}$ (see [28]) is the pair $(C_u^*(\mathcal{S}), \iota)$ where $C_u^*(\mathcal{S})$ is a unital C^* -algebra, $\iota : \mathcal{S} \rightarrow C_u^*(\mathcal{S})$ is a unital complete order embedding such that $\iota(\mathcal{S})$ generates $C_u^*(\mathcal{S})$ as a C^* -algebra, and whenever H is a Hilbert space with $\phi : \mathcal{S} \rightarrow \mathcal{B}(H)$ a unital completely positive map, there exists a $*$ -representation $\pi_\phi : C_u^*(\mathcal{S}) \rightarrow \mathcal{B}(H)$ such that $\pi_\phi \circ \iota = \phi$. We will briefly introduce a multivariate extension of both the maximal and the commuting tensor product (as discussed in the beginning of this section) in the category of operator systems. If $\mathcal{S}_1, \dots, \mathcal{S}_k$ are operator systems, as the maximal tensor product of operator systems is associative (via [27, Theorem 5.5]) we may give an unambiguous meaning to the k -fold maximal tensor product

$$\max - \otimes_{j=1}^k \mathcal{S}_j := \mathcal{S}_1 \otimes_{\max} \cdots \otimes_{\max} \mathcal{S}_k.$$

The k -fold commuting tensor product was initially defined in [20, Section 7]: if H is a Hilbert space and $\phi_j : \mathcal{S}_j \rightarrow \mathcal{B}(H)$ are completely positive maps for $j = 1, \dots, k$, we call the family $(\phi_j)_{j=1}^k$ *commuting* if ϕ_i and ϕ_j have mutually commuting ranges whenever $i \neq j$, for $1 \leq i, j \leq k$. Given a commuting family $(\phi_j)_{j=1}^k$, we define the linear map $\prod_{j=1}^k \phi_j : \otimes_{j=1}^k \mathcal{S}_j \rightarrow \mathcal{B}(H)$ via

$$\left(\prod_{j=1}^k \phi_j \right) (\otimes_{j=1}^k u_j) := \prod_{j=1}^k \phi_j(u_j), \quad u_j \in \mathcal{S}_j, j \in [k].$$

The positive cones for $\mathcal{C} - \otimes_{j=1}^k \mathcal{S}_j$ are then determined by all elements $u \in M_n(\otimes_{j=1}^k \mathcal{S}_j)$ which are positive in $M_n(\mathcal{B}(H))$ under all mutually commuting families and for all choices of Hilbert space H .

The following result shows how we can understand the structure of the multivariate commuting tensor product of operator systems when we view them inside the maximal tensor product of their universal C^* -covers.

Theorem 6.4. *Let $\mathcal{S}_1, \dots, \mathcal{S}_k, k \in \mathbb{N}$ be operator systems. The operator system arising from the inclusion of $\otimes_{j=1}^k \mathcal{S}_j$ into $\max - \otimes_{j=1}^k C_u^*(\mathcal{S}_j)$ coincides with $\mathcal{C} - \otimes_{j=1}^k \mathcal{S}_j$.*

Proof. For the sake of brevity, set $\mathcal{S} = \mathcal{C} - \otimes_{j=1}^k \mathcal{S}_j$. First, suppose $u \in M_n(\mathcal{S})^+$ for some $n \in \mathbb{N}$. We wish to show that $u \in M_n(\max - \otimes_{j=1}^k C_u^*(\mathcal{S}_j))^+$. By [27, Lemma 4.1], it is sufficient to prove that $\phi^{(n)}(u) \geq 0$ for each unital completely positive $\phi : \max - \otimes_{j=1}^k C_u^*(\mathcal{S}_j) \rightarrow \mathcal{B}(H)$. By Stinespring's Theorem, we may also without loss of generality assume ϕ is a $*$ -homomorphism. By [20, Proposition 7.4], associativity, and the universal property of the maximal tensor product of C^* -algebras, each such ϕ is equivalent to $\prod_{j=1}^k \pi_j$, where $\pi_j : C_u^*(\mathcal{S}_j) \rightarrow \mathcal{B}(H)$ is a $*$ -homomorphism for $j = 1, \dots, k$ and all have mutually commuting ranges. As the restriction of π_j to $\mathcal{S}_j$ remains completely positive for $j = 1, \dots, k$, the result follows.

Conversely, let τ be the operator system structure on $\bigotimes_{j=1}^k \mathcal{S}_j$ arising from the inclusion $\bigotimes_{j=1}^k \mathcal{S}_j \subseteq \max - \bigotimes_{j=1}^k C_u^*(\mathcal{S}_j)$. Suppose that $u \in M_n(\tau - \bigotimes_{j=1}^k \mathcal{S}_j)^+$, with $\phi_j : \mathcal{S}_j \rightarrow \mathcal{B}(H)$ completely positive maps with mutually commuting ranges for $j = 1, \dots, k$. By the definition of $C_u^*(\mathcal{S}_j)$, there exists unique $*$ -homomorphisms $\pi_j : C_u^*(\mathcal{S}_j) \rightarrow \mathcal{B}(H)$ extending ϕ_j , for $j = 1, \dots, k$. As $\mathcal{S}_j$ generates $C_u^*(\mathcal{S}_j)$ as a C^* -algebra, we may then conclude the ranges of π_j are mutually commuting as well for $j = 1, \dots, k$. Thus, $(\bigotimes_{j=1}^k \phi_j)^{(n)}(u) \geq 0$, which implies $u \in M_n(\mathcal{S})^+$ - completing the proof. $\square$

Recall (see [33]) that for any Archimedean ordered unit (AOU) space V , there exists a unique operator system $\text{OMIN}(V)$ (respectively, $\text{OMAX}(V)$) with underlying space V , called the *minimal* (respectively, *maximal*) operator system of V which has the universal property that every positive map $\phi : \mathcal{T} \rightarrow V$ (respectively, $\phi : V \rightarrow \mathcal{T}$) where $\mathcal{T}$ is an operator system, is completely positive from $\mathcal{T} \rightarrow \text{OMIN}(V)$ (respectively, $\text{OMAX}(V) \rightarrow \mathcal{T}$).

Lemma 6.5. *Let $V_1, \dots, V_k$, $k \in \mathbb{N}$ be finite dimensional AOU spaces, with units e_j , $j = 1, \dots, k$ respectively. An element $u \in \max - \bigotimes_{j=1}^k \text{OMAX}(V_j)$ is positive if and only if $u = \sum_{i=1}^m v_i^{(1)} \otimes \dots \otimes v_i^{(k)}$, for some $v_i^{(j)} \in V_j^+$, $i = 1, \dots, m$ and $j = 1, \dots, k$.*

Proof. This proof relies on similar ideas as the proof of [39, Lemma 6.6]; we include the details for the convenience of the reader. We only consider the case when $k = 3$; all others will follow similarly. Let D be the set containing all sums of elementary tensors $v_1 \otimes v_2 \otimes v_3$ with $v_i \in V_i^+$, $i = 1, 2, 3$. We claim that if, for every $\epsilon > 0$, there exists $u_\epsilon \in D$ such that $\|u_\epsilon\| \rightarrow 0$ as $\epsilon \rightarrow 0$ and $u + u_\epsilon \in D$ for each $\epsilon > 0$, then $u \in D$. We may, without loss of generality, further assume that $\|u_\epsilon\| \leq 1$ for all $\epsilon > 0$. Set $K = 2\dim(V_1)\dim(V_2)\dim(V_3) + 1$ and, using Carathéodory's Theorem, start by writing

$$u + u_\epsilon = \sum_{j=1}^K v_{1,j}^{(\epsilon)} \otimes v_{2,j}^{(\epsilon)} \otimes v_{3,j}^{(\epsilon)},$$

where $v_{i,j}^{(\epsilon)} \in V_i^+$ and $\|v_{1,j}^{(\epsilon)}\| = \|v_{2,j}^{(\epsilon)}\| = \|v_{3,j}^{(\epsilon)}\|$ for $i = 1, 2, 3$, $j = 1, \dots, K$ and all $\epsilon > 0$. As $v_{1,j}^{(\epsilon)} \otimes v_{2,j}^{(\epsilon)} \otimes v_{3,j}^{(\epsilon)} \leq u + u_\epsilon$ and $\|u + u_\epsilon\| \leq \|u\| + 1$ for all $\epsilon > 0$, we get that $\|v_{i,j}^{(\epsilon)}\| \leq \sqrt[3]{\|u\| + 1}$ for $i = 1, 2, 3$ and $j = 1, \dots, K$. Using the fact that all of our AOU spaces are finite-dimensional, by compactness we can also assume $v_{i,j}^{(\epsilon)} \rightarrow v_{i,j}$ as $\epsilon \rightarrow 0$ for $i = 1, 2, 3$ and all $j = 1, \dots, K$. We may then conclude that $u = \sum_{j=1}^K v_{1,j} \otimes v_{2,j} \otimes v_{3,j} \in D$.

Let

$$S_0 = \sum_{p=1}^{\ell} a_p \otimes v_p^{(1)}, \quad T_0 = \sum_{q=1}^s b_q \otimes v_q^{(2)}, \quad U_0 = \sum_{r=1}^t c_r \otimes v_r^{(3)}, \quad (25)$$

for some $a_p \in M_n^+, v_p^{(1)} \in V_1^+, p = 1, \dots, \ell, b_q \in M_m^+, v_q^{(2)} \in V_2^+, q = 1, \dots, s$ and $c_r \in M_d^+, v_r^{(3)} \in V_3^+, r = 1, \dots, t$. If $\alpha \in M_{1,nmd}$, then

$$\alpha(S_0 \otimes T_0 \otimes U_0)\alpha^* = \sum_{p=1}^{\ell} \sum_{q=1}^s \sum_{r=1}^t (\alpha(a_p \otimes b_q \otimes c_r)\alpha^*) v_p^{(1)} \otimes v_q^{(2)} \otimes v_r^{(3)} \in D.$$

Now suppose first that $S \in M_n(\text{OMAX}(V_1))^+$ and $\alpha \in M_{1,nmd}$. By the description of the multivariate maximal tensor product as provided in [20, Proposition 7.2], if $\epsilon > 0$ then $S + \epsilon 1_n$ has the form of S_0 as in (25). Therefore,

$$\alpha(S \otimes T_0 \otimes U_0)\alpha^* + \epsilon \alpha(1_n \otimes T_0 \otimes U_0)\alpha^* = \alpha((S + \epsilon 1_n) \otimes T_0 \otimes U_0)\alpha^* \in D.$$

Since $\alpha(1_n \otimes T_0 \otimes U_0)\alpha^* \in D$ (using [33]), using our initial arguments we conclude that

$$\alpha(S \otimes T_0 \otimes U_0)\alpha^* \in D.$$

If we now pick $T \in M_m(\text{OMAX}(V_2))^+$ and $U \in M_d(\text{OMAX}(V_3))^+$, using similar arguments as before we may conclude that $\alpha(S \otimes T \otimes U)\alpha^* \in D$.

Let $u \in \max - \bigotimes_{j=1}^3 \text{OMAX}(V_j)$ be positive; by the definition of the multivariate maximal tensor product [20], for every $\epsilon > 0$ there exists $n, m, d \in \mathbb{N}, S \in M_n(\text{OMAX}(V_1))^+, T \in M_m(\text{OMAX}(V_2))^+, U \in M_d(\text{OMAX}(V_3))^+$ and $\alpha \in M_{1,nmd}$ such that $u + \epsilon 1 = \alpha(S \otimes T \otimes U)\alpha^*$. By the previous and first paragraphs, this implies $u \in D$. $\square$

For a linear functional

$$s : \mathcal{T}_{X_2, X_1} \otimes \mathcal{T}_{Y_2, Y_1} \otimes \mathcal{T}_{A_1, A_2} \otimes \mathcal{T}_{B_1, B_2} \rightarrow \mathbb{C},$$

let $\Gamma_s : M_{X_2 Y_2 A_1 B_1} \rightarrow M_{X_1 Y_1 A_2 B_2}$ be the linear map whose Choi matrix coincides with

$$(s(e_{x_2, x'_2, x_1, x'_1} \otimes e_{y_2, y'_2, y_1, y'_1} \otimes e_{a_1, a'_1, a_2, a'_2} \otimes e_{b_1, b'_1, b_2, b'_2}))_{x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1}^{x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2} \quad (26)$$

Theorem 6.6. *The map $s \mapsto \Gamma_s$ is an affine isomorphism from*

(i) *the state space of*

$$\mathcal{T}_{X_2, X_1} \otimes_{\max} \mathcal{T}_{Y_2, Y_1} \otimes_{\max} \mathcal{T}_{A_1, A_2} \otimes_{\max} \mathcal{T}_{B_1, B_2}$$

onto $\mathcal{Q}_{\text{sns}}$;

(ii) *the state space of*

$$\mathcal{T}_{X_2, X_1} \otimes_{\mathbb{C}} \mathcal{T}_{Y_2, Y_1} \otimes_{\mathbb{C}} \mathcal{T}_{A_1, A_2} \otimes_{\mathbb{C}} \mathcal{T}_{B_1, B_2}$$

onto $\mathcal{Q}_{\text{sqc}}$;

(iii) *the state space of*

$$\mathcal{T}_{X_2, X_1} \otimes_{\min} \mathcal{T}_{Y_2, Y_1} \otimes_{\min} \mathcal{T}_{A_1, A_2} \otimes_{\min} \mathcal{T}_{B_1, B_2}$$

onto $\mathcal{Q}_{\text{sqa}}$.

(iv) *the state space of*

$$\text{OMIN}(\mathcal{T}_{X_2, X_1}) \otimes_{\min} \text{OMIN}(\mathcal{T}_{Y_2, Y_1}) \otimes_{\min} \text{OMIN}(\mathcal{T}_{A_1, A_2}) \otimes_{\min} \text{OMIN}(\mathcal{T}_{B_1, B_2})$$

onto $\mathcal{Q}_{\text{sloc}}$.

Proof. (i) Let $\Gamma \in \mathcal{Q}_{\text{sns}}$. If

$$C = (C_{x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1}^{x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2})$$

is the Choi matrix of Γ (where the indices above range over the corresponding sets $X_i, Y_i, A_i, B_i, i = 1, 2$), then $C \in M_{X_2 Y_2 A_1 B_1, X_1 Y_1 A_2 B_2}^+$. We also note that condition (12) implies that for $y_i, y'_i \in Y_i, a_i, a'_i \in A_i$ and $b_i, b'_i \in B_i, i = 1, 2$ there exists a constant $C_{y_2 y'_2, a_1 a'_1, b_1 b'_1}^{y_1 y'_1, a_2 a'_2, b_2 b'_2} \in \mathbb{C}$ such that

$$\sum_{x_1} C_{x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1}^{x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2} = \delta_{x_2, x'_2} C_{y_2 y'_2, a_1 a'_1, b_1 b'_1}^{y_1 y'_1, a_2 a'_2, b_2 b'_2},$$

for all $x_2, x'_2 \in X_2$. This equality implies

$$L_\rho(C) \in \mathcal{L}_{X_2, X_1} \text{ for all } \rho \in M_{Y_2 Y_1 A_1 A_2 B_1 B_2}.$$

Similarly, using conditions (13)-(15) we can make a similar argument showing

$$L_\rho(C) \in \mathcal{L}_{Y_2, Y_1} \text{ for all } \rho \in M_{X_2 X_1 A_1 A_2 B_1 B_2},$$

$$L_\rho(C) \in \mathcal{L}_{A_1, A_2} \text{ for all } \rho \in M_{X_2 X_1 Y_2 Y_1 B_1 B_2},$$

and

$$L_\rho(C) \in \mathcal{L}_{B_1, B_2} \text{ for all } \rho \in M_{X_2 X_1 Y_2 Y_1 A_1 A_2}.$$

Thus,

$$C \in (\mathcal{L}_{X_2, X_1} \otimes \mathcal{L}_{Y_2, Y_1} \otimes \mathcal{L}_{A_1, A_2} \otimes \mathcal{L}_{B_1, B_2}) \cap M_{X_2 Y_2 A_1 B_1, X_1 Y_1 A_2 B_2}^+;$$

by the injectivity of the minimal operator system tensor product,

$$C \in (\mathcal{L}_{X_2, X_1} \otimes_{\min} \mathcal{L}_{Y_2, Y_1} \otimes_{\min} \mathcal{L}_{A_1, A_2} \otimes_{\min} \mathcal{L}_{B_1, B_2})^+.$$

By [17, Proposition 1.9], and [39, Proposition 5.5],

$$(\mathcal{T}_{X_2, X_1} \otimes_{\max} \mathcal{T}_{Y_2, Y_1} \otimes_{\max} \mathcal{T}_{A_1, A_2} \otimes_{\max} \mathcal{T}_{B_1, B_2})^d \cong_{\text{c.o.i.}} \mathcal{L}_{X_2, X_1} \otimes_{\min} \mathcal{L}_{Y_2, Y_1} \otimes_{\min} \mathcal{L}_{A_1, A_2} \otimes_{\min} \mathcal{L}_{B_1, B_2}.$$

Arguing now as in [39, Theorem 6.2], this establishes the claim.

(ii) First suppose that

$$s : \mathcal{T}_{X_2, X_1} \otimes_{\text{c}} \mathcal{T}_{Y_2, Y_1} \otimes_{\text{c}} \mathcal{T}_{A_1, A_2} \otimes_{\text{c}} \mathcal{T}_{B_1, B_2} \rightarrow \mathbb{C}$$

is a state. By Theorem 6.4, we may consider the state s as the restriction of a state

$$\tilde{s} : \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{Y_2, Y_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \otimes_{\max} \mathcal{C}_{B_1, B_2} \rightarrow \mathbb{C}.$$

Applying the GNS construction to $\tilde{s}$ and using [39, Theorem 5.2], we obtain a Hilbert space H , a unit vector $\xi \in H$ and mutually commuting stochastic operator matrices

$$\begin{aligned} (E_{x_2, x'_2, x_1, x'_1})_{x_2, x'_2}^{x_1, x'_1} &\in M_{X_2 X_1} \otimes \mathcal{B}(H), & (E_{y_2, y'_2, y_1, y'_1})_{y_2, y'_2}^{y_1, y'_1} &\in M_{Y_2 Y_1} \otimes \mathcal{B}(H), \\ (F_{a_1, a'_1, a_2, a'_2})_{a_1, a'_1}^{a_2, a'_2} &\in M_{A_1 A_2} \otimes \mathcal{B}(H), & (F_{b_1, b'_1, b_2, b'_2})_{b_1, b'_1}^{b_2, b'_2} &\in M_{B_1 B_2} \otimes \mathcal{B}(H) \end{aligned}$$

satisfying

$$\begin{aligned} s(e_{x_2, x'_2, x_1, x'_1} \otimes e_{y_2, y'_2, y_1, y'_1} \otimes e_{a_1, a'_1, a_2, a'_2} \otimes e_{b_1, b'_1, b_2, b'_2}) = \\ \langle E_{x_2, x'_2, x_1, x'_1} E_{y_2, y'_2, y_1, y'_1} F_{a_1, a'_1, a_2, a'_2} F_{b_1, b'_1, b_2, b'_2}, \xi \xi^* \rangle \end{aligned}$$

for all $x_i, x'_i, y_i, y'_i, a_i, a'_i, b_i, b'_i, i = 1, 2$. Setting $E_{x_2, x'_2, y_2, y'_2}^{x_1, x'_1, y_1, y'_1} = E_{x_2, x'_2, x_1, x'_1} E_{y_2, y'_2, y_1, y'_1}$ and $F_{a_1, a'_1, b_1, b'_1}^{a_2, a'_2, b_2, b'_2} = F_{a_1, a'_1, a_2, a'_2} F_{b_1, b'_1, b_2, b'_2}$, we have that the stochastic operator matrices

$$E = (E_{x_2, x'_2, y_2, y'_2}^{x_1, x'_1, y_1, y'_1})_{x_2, x'_2, y_2, y'_2}^{x_1, x'_1, y_1, y'_1}, \quad F = (F_{a_1, a'_1, b_1, b'_1}^{a_2, a'_2, b_2, b'_2})_{a_1, a'_1, b_1, b'_1}^{a_2, a'_2, b_2, b'_2}$$

are dilatable and have mutually commuting entries. Thus, $\Gamma_s = \Gamma_{E, F, \xi}$ with $\Gamma_s \in \mathcal{Q}_{\text{sqc}}$.

Conversely, suppose that $\Gamma \in \mathcal{Q}_{\text{sqc}}$; by definition, there exists a Hilbert space K , a unit vector $\eta \in K$ and mutually commuting strongly stochastic operator matrices

$$\begin{aligned} E_X &= (E_{x_2, x'_2, x_1, x'_1})_{x_2, x'_2, x_1, x'_1}, & E_Y &= (E_{y_2, y'_2, y_1, y'_1})_{y_2, y'_2, y_1, y'_1}, \\ F_A &= (F_{a_1, a'_1, a_2, a'_2})_{a_1, a'_1, a_2, a'_2}, & F_B &= (F_{b_1, b'_1, b_2, b'_2})_{b_1, b'_1, b_2, b'_2} \end{aligned}$$

acting on K so that $\Gamma = \Gamma_{E, F, \eta}$, for $E = E_X \cdot E_Y$ and $F = F_A \cdot F_B$. Let π_X, π_Y, π_A and π_B be the (unital) $*$ -representations of $\mathcal{C}_{X_2, X_1}, \mathcal{C}_{Y_2, Y_1}, \mathcal{C}_{A_1, A_2}$ and $\mathcal{C}_{B_1, B_2}$ on $\mathcal{B}(K)$ arising from E_X, E_Y, F_A , and F_B respectively (see [39, Theorem 5.2]). Then $\pi := \pi_X \otimes \pi_Y \otimes \pi_A \otimes \pi_B$ is a unital $*$ -representation of $\mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{Y_2, Y_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \otimes_{\max} \mathcal{C}_{B_1, B_2}$ on $\mathcal{B}(K)$. Using Theorem 6.4 once more, if we let s be the restriction to $\mathcal{T}_{X_2, X_1} \otimes_{\mathbb{C}} \mathcal{T}_{Y_2, Y_1} \otimes_{\mathbb{C}} \mathcal{T}_{A_1, A_2} \otimes_{\mathbb{C}} \mathcal{T}_{B_1, B_2}$ of the state

$$\begin{aligned} \tilde{s} : \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{Y_2, Y_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \otimes_{\max} \mathcal{C}_{B_1, B_2} &\rightarrow \mathbb{C}, \\ w &\mapsto \langle \pi(w)\eta, \eta \rangle, \end{aligned}$$

it is clear that $\Gamma = \Gamma_s$.

(iii) First, let Γ be a quantum SQNS correlation. By definition, there exists stochastic operator matrices

$$\begin{aligned} M_X &= (M_{x_2, x'_2, x_1, x'_1})_{x_2, x'_2, x_1, x'_1}, & M_A &= (M_{a_1, a'_1, a_2, a'_2})_{a_1, a'_1, a_2, a'_2}, \\ N_Y &= (N_{y_2, y'_2, y_1, y'_1})_{y_2, y'_2, y_1, y'_1}, & N_B &= (N_{b_1, b'_1, b_2, b'_2})_{b_1, b'_1, b_2, b'_2} \end{aligned}$$

acting on finite dimensional Hilbert spaces H_X, H_A, H_Y and H_B , respectively, along with unit vector $\eta \in H_X \otimes H_A \otimes H_Y \otimes H_B$ such that

$$\begin{aligned} & \langle \Gamma(\epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2} \otimes \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1}), \epsilon_{x_1 x'_1} \otimes \epsilon_{y_1 y'_1} \otimes \epsilon_{a_2 a'_2} \otimes \epsilon_{b_2 b'_2} \rangle = \\ & \left\langle (M_{x_2 x'_2, x_1 x'_1} \otimes M_{a_1 a'_1, a_2 a'_2} \otimes N_{y_2 y'_2, y_1 y'_1} \otimes N_{b_1 b'_1, b_2 b'_2}), \eta \eta^* \right\rangle \end{aligned}$$

for $x_i, x'_i \in X_i, y_i, y'_i \in Y_i, a_i, a'_i \in A_i, b_i, b'_i \in B_i, i = 1, 2$. Let $\pi_X : \mathcal{C}_{X_2, X_1} \rightarrow \mathcal{B}(H_X), \pi_Y : \mathcal{C}_{Y_2, Y_1} \rightarrow \mathcal{B}(H_Y), \pi_A : \mathcal{C}_{A_1, A_2} \rightarrow \mathcal{B}(H_A)$ and $\pi_B : \mathcal{C}_{B_1, B_2} \rightarrow \mathcal{B}(H_B)$ be the unital $*$ -representations arising from M_X, M_A, N_Y , and N_B respectively. Let $\pi := \pi_X \otimes \pi_Y \otimes \pi_A \otimes \pi_B, \tilde{\eta} \in H_X \otimes H_Y \otimes H_A \otimes H_B$ be the unit vector obtained from applying the canonical shuffle to $\eta, \tilde{s}$ be the state given by

$$\begin{aligned} \tilde{s} : \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{Y_2, Y_1} \otimes_{\min} \mathcal{C}_{A_1, A_2} \otimes_{\min} \mathcal{C}_{B_1, B_2} &\rightarrow \mathbb{C}, \\ w &\mapsto \langle \pi(w) \tilde{\eta}, \tilde{\eta} \rangle, \end{aligned}$$

and s be the restriction of $\tilde{s}$ to $\mathcal{T}_{X_2, X_1} \otimes_{\min} \mathcal{T}_{Y_2, Y_1} \otimes_{\min} \mathcal{T}_{A_1, A_2} \otimes_{\min} \mathcal{T}_{B_1, B_2}$; it is clear that $\Gamma = \Gamma_s$ in this case.

If $\Gamma \in \mathcal{Q}_{\text{sqa}}$, let $(\Gamma_n)_{n \in \mathbb{N}}$ be a sequence of quantum SQNS correlations with $\Gamma_n \rightarrow \Gamma$ as $n \rightarrow \infty$. By the previous paragraph, for each $n \in \mathbb{N}$ we may pick a state $s_n : \mathcal{T}_{X_2, X_1} \otimes_{\min} \mathcal{T}_{Y_2, Y_1} \otimes_{\min} \mathcal{T}_{A_1, A_2} \otimes_{\min} \mathcal{T}_{B_1, B_2} \rightarrow \mathbb{C}$ such that $\Gamma_n = \Gamma_{s_n}$. If s is a cluster point (in the weak * topology) of the sequence of states $(s_n)_{n \in \mathbb{N}}$, we have $\Gamma = \Gamma_s$.

Now let $s : \mathcal{T}_{X_2, X_1} \otimes_{\min} \mathcal{T}_{Y_2, Y_1} \otimes_{\min} \mathcal{T}_{A_1, A_2} \otimes_{\min} \mathcal{T}_{B_1, B_2} \rightarrow \mathbb{C}$ be a state, and using the fact that the minimal operator system tensor product is injective (see [20]) let $\tilde{s} : \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{Y_2, Y_1} \otimes_{\min} \mathcal{C}_{A_1, A_2} \otimes_{\min} \mathcal{C}_{B_1, B_2} \rightarrow \mathbb{C}$ be an extension of s . By [24, Corollary 4.3.10], $\tilde{s}$ can be approximated in the weak * topology by elements of the convex hull of vector states on $\pi_X(\mathcal{C}_{X_2, X_1}) \otimes_{\min} \pi_Y(\mathcal{C}_{Y_2, Y_1}) \otimes_{\min} \pi_A(\mathcal{C}_{A_1, A_2}) \otimes_{\min} \pi_B(\mathcal{C}_{B_1, B_2})$ (where π_X, π_Y, π_A , and π_B are unital $*$ -representations of $\mathcal{C}_{X_2, X_1}, \mathcal{C}_{Y_2, Y_1}, \mathcal{C}_{A_1, A_2}$ and $\mathcal{C}_{B_1, B_2}$, respectively). Using an argument similar to the proof of [7, Theorem 5.6] or [39, Theorem 6.5], we can show that Γ is a limit of quantum SQNS correlations.

(iv) This proof is along the lines of the proof for [39, Theorem 6.7]; we include the details for the convenience of the reader. We first let s be a state on

$$\text{OMIN}(\mathcal{T}_{X_2, X_1}) \otimes_{\min} \text{OMIN}(\mathcal{T}_{Y_2, Y_1}) \otimes_{\min} \text{OMIN}(\mathcal{T}_{A_1, A_2}) \otimes_{\min} \text{OMIN}(\mathcal{T}_{B_1, B_2}).$$

By [26, Theorem 9.9] and [17, Proposition 1.9], we may consider s as an element of

$$\left(\text{OMAX}(\mathcal{T}_{X_2, X_1}) \otimes_{\max} \text{OMAX}(\mathcal{T}_{Y_2, Y_1}) \otimes_{\max} \text{OMAX}(\mathcal{T}_{A_1, A_2}) \otimes_{\max} \text{OMAX}(\mathcal{T}_{B_1, B_2}) \right)^+.$$

By Lemma 6.5, there exist states $\phi_X^{(j)} \in (\mathcal{T}_{X_2, X_1})^+, \phi_Y^{(j)} \in (\mathcal{T}_{Y_2, Y_1})^+, \phi_A^{(j)} \in (\mathcal{T}_{A_1, A_2})^+$ and $\phi_B^{(j)} \in (\mathcal{T}_{B_1, B_2})^+$, and non-negative scalars $\lambda_j, j = 1, \dots, k$ such

that $s = \sum_{j=1}^k \lambda_j \phi_X^{(j)} \otimes \phi_Y^{(j)} \otimes \phi_A^{(j)} \otimes \phi_B^{(j)}$. Set

$$\begin{aligned} E_X^{(j)} &= (\phi_X^{(j)}(e_{x_2, x'_2, x_1, x'_1}))_{x_2 x'_2, x_1 x'_1}, & E_Y^{(j)} &= (\phi_Y^{(j)}(e_{y_2, y'_2, y_1, y'_1}))_{y_2 y'_2, y_1 y'_1}, \\ E_A^{(j)} &= (\phi_A^{(j)}(e_{a_1, a'_1, a_2, a'_2}))_{a_1 a'_1, a_2 a'_2}, & E_B^{(j)} &= (\phi_B^{(j)}(e_{b_1, b'_1, b_2, b'_2}))_{b_1 b'_1, b_2 b'_2}, \end{aligned}$$

and let $\Phi_X^{(j)} : M_{X_2} \rightarrow M_{X_1}$, $\Phi_Y^{(j)} : M_{Y_2} \rightarrow M_{Y_1}$, $\Phi_A^{(j)} : M_{A_1} \rightarrow M_{A_2}$, and $\Phi_B^{(j)} : M_{B_1} \rightarrow M_{B_2}$ be the quantum channels with Choi matrices $E_X^{(j)}$, $E_Y^{(j)}$, $E_A^{(j)}$, and $E_B^{(j)}$ (respectively) for $j = 1, \dots, k$. Clearly,

$$\Gamma_s = \sum_{j=1}^k \lambda_j \Phi_X^{(j)} \otimes \Phi_Y^{(j)} \otimes \Phi_A^{(j)} \otimes \Phi_B^{(j)}. \quad (27)$$

By Remark 3.11, this shows $\Gamma_s \in \mathcal{Q}_{\text{sloc}}$.

Now suppose Γ is of the form (27), with s a functional on $\mathcal{T}_{X_2, X_1} \otimes \mathcal{T}_{Y_2, Y_1} \otimes \mathcal{T}_{A_1, A_2} \otimes \mathcal{T}_{B_1, B_2}$ such that $\Gamma = \Gamma_s$. Let $E_X^{(j)} \in (M_{X_2} \otimes M_{X_1})^+$ (resp. $E_Y^{(j)} \in (M_{Y_2} \otimes M_{Y_1})^+$, $E_A^{(j)} \in (M_{A_1} \otimes M_{A_2})^+$, and $E_B^{(j)} \in (M_{B_1} \otimes M_{B_2})^+$) be the Choi matrix of $\Phi_X^{(j)}$ (resp. $\Phi_Y^{(j)}$, $\Phi_A^{(j)}$ and $\Phi_B^{(j)}$); then $E_X^{(j)}$, $E_Y^{(j)}$, $E_A^{(j)}$, and $E_B^{(j)}$ are stochastic operator matrices acting on $\mathbb{C}$. By [39, Theorem 5.2], there exist positive functionals $\phi_X^{(j)} : \mathcal{T}_{X_2, X_1} \rightarrow \mathbb{C}$, $\phi_Y^{(j)} : \mathcal{T}_{Y_2, Y_1} \rightarrow \mathbb{C}$, $\phi_A^{(j)} : \mathcal{T}_{A_1, A_2} \rightarrow \mathbb{C}$, and $\phi_B^{(j)} : \mathcal{T}_{B_1, B_2} \rightarrow \mathbb{C}$ such that

$$\begin{aligned} E_X^{(j)} &= (\phi_X^{(j)}(e_{x_2, x'_2, x_1, x'_1}))_{x_2 x'_2, x_1 x'_1}, & E_Y^{(j)} &= (\phi_Y^{(j)}(e_{y_2, y'_2, y_1, y'_1}))_{y_2 y'_2, y_1 y'_1}, \\ E_A^{(j)} &= (\phi_A^{(j)}(e_{a_1, a'_1, a_2, a'_2}))_{a_1 a'_1, a_2 a'_2}, & E_B^{(j)} &= (\phi_B^{(j)}(e_{b_1, b'_1, b_2, b'_2}))_{b_1 b'_1, b_2 b'_2}, \end{aligned}$$

for $j = 1, \dots, k$. It is thus straightforward to see that s is the functional corresponding to

$$\sum_{j=1}^k \lambda_j \phi_X^{(j)} \otimes \phi_Y^{(j)} \otimes \phi_A^{(j)} \otimes \phi_B^{(j)},$$

and thus, by Lemma 6.5 is a state on

$$\text{OMIN}(\mathcal{T}_{X_2, X_1}) \otimes_{\min} \text{OMIN}(\mathcal{T}_{Y_2, Y_1}) \otimes_{\min} \text{OMIN}(\mathcal{T}_{A_1, A_2}) \otimes_{\min} \text{OMIN}(\mathcal{T}_{B_1, B_2}),$$

as claimed. $\square$

Remark 6.7. We note that, using an almost identical argument as in (iv), we have an affine isomorphism between the state space of $\text{OMIN}(\mathcal{S}_{X_2, X_1}) \otimes_{\min} \text{OMIN}(\mathcal{S}_{Y_2, Y_1}) \otimes_{\min} \text{OMIN}(\mathcal{S}_{A_1, A_2}) \otimes_{\min} \text{OMIN}(\mathcal{S}_{B_1, B_2})$ and $\mathcal{C}_{\text{sloc}}$; the classical analogue for cases (i)-(iii) are addressed in [20, Theorem 7.11].

Corollary 6.8. *The set $\mathcal{Q}_{\text{sqc}}$ is closed and convex.*

Proof. By Theorem 6.6, it is straightforward to show that the affine mapping $s \mapsto \Gamma_s$ is also homeomorphism when the state space of $\mathcal{T}_{X_2 X_1} \otimes_c \mathcal{T}_{Y_2 Y_1} \otimes_c \mathcal{T}_{A_1 A_2} \otimes_c \mathcal{T}_{B_1 B_2}$ is equipped with the weak*-topology. As the state space will be weak*-compact, the range of this homeomorphism must be (convex and) closed. $\square$

Remark 6.9. By [20, Theorem 7.11], we may use an identical argument as in Corollary 6.8 to show that $\mathcal{C}_{\text{sqc}}$ is closed and convex.

Remark 6.10. Suppose that for finite sets $X_i, Y_i, A_i, B_i, i = 1, 2$ we make the additional restriction that $Y_i = B_i = [1], i = 1, 2$. One may easily verify that for $t \in \{\text{loc}, q, \text{qa}, \text{qc}, \text{ns}\}$ we have the reduction(s)

$$\begin{aligned}\mathcal{Q}_{\text{st}}(X_2[1], A_1[1], X_1[1], A_2[1]) &= \mathcal{Q}_t(X_2, A_1, X_1, A_2), \\ \mathcal{C}_{\text{st}}(X_2[1], A_1[1], X_1[1], A_2[1]) &= \mathcal{C}_t(X_2, A_1, X_1, A_2).\end{aligned}$$

Theorem 6.11. For all finite sets $X_i, Y_i, A_i, B_i, i = 1, 2$ of sufficiently large cardinality, the following hold true:

- (i) $\mathcal{Q}_{\text{sqa}}(X_2 Y_2, A_1 B_1, X_1 Y_1, A_2 B_2) \neq \mathcal{Q}_{\text{sqc}}(X_2 Y_2, A_1 B_1, X_1 Y_1, A_2 B_2)$.
- (ii) $\mathcal{T}_{X_2, X_1} \otimes_c \mathcal{T}_{Y_2, Y_1} \otimes_c \mathcal{T}_{A_1, A_2} \otimes_c \mathcal{T}_{B_1, B_2} \neq \mathcal{T}_{X_2, X_1} \otimes_{\min} \mathcal{T}_{Y_2, Y_1} \otimes_{\min} \mathcal{T}_{A_1, A_2} \otimes_{\min} \mathcal{T}_{B_1, B_2}$.
- (iii) $\mathcal{C}_{\text{sqa}}(X_2 Y_2, A_1 B_1, X_1 Y_1, A_2 B_2) \neq \mathcal{C}_{\text{sqc}}(X_2 Y_2, A_1 B_1, X_1 Y_1, A_2 B_2)$.
- (iv) $\mathcal{S}_{X_2, X_1} \otimes_c \mathcal{S}_{Y_2, Y_1} \otimes_c \mathcal{S}_{A_1, A_2} \otimes_c \mathcal{S}_{B_1, B_2} \neq \mathcal{S}_{X_2, X_1} \otimes_{\min} \mathcal{S}_{Y_2, Y_1} \otimes_{\min} \mathcal{S}_{A_1, A_2} \otimes_{\min} \mathcal{S}_{B_1, B_2}$.

Proof. Statement (i) follows from Remark 6.10, and an application of [39, Theorem 8.3]. Statement (ii) follows from (i), and Theorem 6.6. Statements (iii) and (iv) follow similarly, when paired with [39, Remark 8.1]. $\square$

As in the setup before Theorem 6.6, for a linear functional

$$s : \mathcal{T}_X \otimes \mathcal{T}_Y \otimes \mathcal{T}_A \otimes \mathcal{T}_B \rightarrow \mathbb{C},$$

let $\Gamma_s : M_{XYAB} \rightarrow M_{XYAB}$ be the linear map whose Choi matrix coincides with (26).

Theorem 6.12. The map $s \mapsto \Gamma_s$ is an affine isomorphism from

- (i) the state space of $\mathcal{T}_X \otimes_{\max} \mathcal{T}_Y \otimes_{\max} \mathcal{T}_A \otimes_{\max} \mathcal{T}_B$ onto $\mathcal{Q}_{\text{sns}}^{\text{bi}}$;
- (ii) the state space of $\mathcal{T}_X \otimes_c \mathcal{T}_Y \otimes_c \mathcal{T}_A \otimes_c \mathcal{T}_B$ onto $\mathcal{Q}_{\text{sqc}}^{\text{bi}}$;
- (iii) the state space of $\mathcal{T}_X \otimes_{\min} \mathcal{T}_Y \otimes_{\min} \mathcal{T}_A \otimes_{\min} \mathcal{T}_B$ onto $\mathcal{Q}_{\text{sqa}}^{\text{bi}}$;
- (iv) the state space of $\text{OMIN}(\mathcal{T}_X) \otimes_{\min} \text{OMIN}(\mathcal{T}_Y) \otimes_{\min} \text{OMIN}(\mathcal{T}_A) \otimes_{\min} \text{OMIN}(\mathcal{T}_B)$ onto $\mathcal{Q}_{\text{sloc}}^{\text{bi}}$.

Proof. (i) Let $\Gamma \in \mathcal{Q}_{\text{sns}}^{\text{bi}}$. If

$$C = (C_{x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1}^{x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2})$$

is the Choi matrix of Γ and

$$\tilde{C} = (\tilde{C}_{x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2}^{x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1})$$

is the Choi matrix of Γ^* (where the indices above range over the corresponding sets X, Y, A , and B) then $C, \tilde{C} \in M_{XYABXYAB}^+$. As Γ and Γ^* are both SNS, for $y_i, y'_i \in Y, a_i, a'_i \in A, b_i, b'_i \in B, i = 1, 2$ there exist complex constants $C_{y_1 y'_1, a_1 a'_1, b_1 b'_1}^{y_2 y'_2, a_2 a'_2, b_2 b'_2}$ and $\tilde{C}_{y_1 y'_1, a_2 a'_2, b_2 b'_2}^{y_2 y'_2, a_1 a'_1, b_1 b'_1}$ such that

$$\sum_{x_1} C_{x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1}^{x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2} = \delta_{x_2, x'_2} C_{y_2 y'_2, a_1 a'_1, b_1 b'_1}^{y_1 y'_1, a_2 a'_2, b_2 b'_2},$$

$$\sum_{x_2} \tilde{C}_{x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2}^{x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1} = \delta_{x_1, x'_1} \tilde{C}_{y_1 y'_1, a_2 a'_2, b_2 b'_2}^{y_2 y'_2, a_1 a'_1, b_1 b'_1},$$

for all $x_i, x'_i \in X, i = 1, 2$. Note that

$$\tilde{C}_{x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2}^{x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1} = C_{x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1}^{x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2},$$

for $x_i, x'_i \in X, y_i, y'_i \in Y, a_i, a'_i \in A, b_i, b'_i \in B, i = 1, 2$. These together imply

$$L_\rho(C) \in \mathcal{L}_X \text{ for all } \rho \in M_{YAB}.$$

We then may use the other SNS conditions for Γ and Γ^* to show that

$$L_\rho(C) \in \mathcal{L}_Y \text{ for all } \rho \in M_{XAB},$$

$$L_\rho(C) \in \mathcal{L}_A \text{ for all } \rho \in M_{XYB},$$

and

$$L_\rho(C) \in \mathcal{L}_B \text{ for all } \rho \in M_{XYA}.$$

Together, these imply $C \in (\mathcal{L}_X \otimes \mathcal{L}_Y \otimes \mathcal{L}_A \otimes \mathcal{L}_B)^+$. Use [7, Proposition 3.6] and argue as in Theorem 6.6 (i) to finish.

(ii)-(iv) The proofs for (ii)-(iv) follow exactly the same as in the proofs of Theorem 6.6 (ii)-(iv), just by replacing the use of [39, Theorem 5.2] with [7, Theorem 3.4], the use of Remark 3.11 with Remark 3.13, and by replacing $\mathcal{T}_{X_2, X_1}$ and $\mathcal{C}_{X_2, X_1}$ (resp. $\mathcal{T}_{Y_2, Y_1}$ and $\mathcal{C}_{Y_2, Y_1}$, $\mathcal{T}_{A_1, A_2}$ and $\mathcal{C}_{A_1, A_2}$, $\mathcal{T}_{B_1, B_2}$ and $\mathcal{C}_{B_1, B_2}$) with $\mathcal{T}_X$ and $\mathcal{C}_X$ (resp. $\mathcal{T}_Y$ and $\mathcal{C}_Y$, $\mathcal{T}_A$ and $\mathcal{C}_A$, $\mathcal{T}_B$ and $\mathcal{C}_B$). $\square$

We end this subsection with the following (obvious) result.

Corollary 6.13. *The set $\mathcal{Q}_{\text{sqc}}^{\text{bi}}$ is closed and convex.*

6.2. Concurrent strategies. The goal of this subsection is to apply the results connecting subclasses of strongly QNS correlations with state spaces on tensor products of canonical operator systems previously developed, to the study of a particular class of quantum input-output game which are called *concurrent games*. A quantum non-local game $\varphi : \mathcal{P}_{XX} \rightarrow \mathcal{P}_{AA}$ is *concurrent* if $\varphi(\tilde{J}_X) = \tilde{J}_A$. A QNS correlation $\Phi \in \mathcal{Q}_{\text{ns}}$ is called concurrent if $\Phi(\tilde{J}_X) = \tilde{J}_A$; equivalently, if Φ is a perfect strategy for the (trivial) implication game $\varphi_{\tilde{J}_X \rightarrow \tilde{J}_A}$.

The study of concurrent games and their perfect strategies has grown in recent years (see [6, 7, 39]), and is of particular interest for its connections to the study of quantum automorphism groups and compact quantum groups, along

with its ramifications for the study of synchronous games; indeed— concurrent games were introduced as a “quantization” of synchronous games. The reasoning behind this claim comes from [6, Remark 2.1]: for classical non-local game $\lambda : X \times X \times A \times A \rightarrow \{0, 1\}$, then λ is synchronous if and only if $\varphi_\lambda(J_X^{\text{cl}}) \leq J_A^{\text{cl}}$. Concurrency allows us to see how a correlation affects the “non-classical” portion of the (unnormalized) maximally entangled state— ideally, sending it to a state supported on $\tilde{J}_A$.

Let $\tau : \mathcal{C}_{X,A} \rightarrow \mathbb{C}$ be a tracial state; the linear map $\Gamma_\tau : M_{XX} \rightarrow M_{AA}$ given by

$$\Gamma_\tau(\epsilon_{xx'} \otimes \epsilon_{yy'}) = \sum_{a,a',b,b' \in A} \tau(e_{x,x',a,a'} e_{y',y,b',b}) \epsilon_{aa'} \otimes \epsilon_{bb'},$$

is a QNS correlation. This particular type of a QNS correlation is called *tracial*, and was first introduced in [39]. Subclasses of *quantum tracial* and *locally tracial* QNS correlations are defined by requiring that τ factors through a finite-dimensional and abelian $*$ -representation, respectively. We focus on such QNS correlations for their connection to concurrent games: it was first established in [6, Theorem 4.1] that any perfect strategy Γ for a concurrent game φ must be tracial, in the sense defined above. In the rest of this section, we will develop a notion of when SQNS correlations are tracial, and describe their structure.

Let $\Gamma : M_{X_2 X_2 \times A_1 A_1} \rightarrow M_{X_1 X_1 \times A_2 A_2}$ be an SQNS correlation, and let $\sigma \in M_{A_1 A_1}$ be an arbitrary state. Define the mapping $\Gamma_{X_2 X_2 \rightarrow X_1 X_1}^\sigma : M_{X_2 X_2} \rightarrow M_{X_1 X_1}$ via

$$\Gamma_{X_2 X_2 \rightarrow X_1 X_1}^\sigma(\rho) := \text{Tr}_{A_2 A_2} \Gamma(\rho \otimes \sigma), \quad \rho \in M_{X_2 X_2}.$$

One easily shows that $\Gamma_{X_2 X_2 \rightarrow X_1 X_1}^\sigma$ is a well-defined QNS correlation; furthermore, $\Gamma_{X_2 X_2 \rightarrow X_1 X_1}^\sigma = \Gamma_{X_2 X_2 \rightarrow X_1 X_1}^{\sigma'}$ for any states $\sigma, \sigma' \in M_{A_1 A_1}$. Thus, we let $\Gamma_{X_2 X_2 \rightarrow X_1 X_1}$ denote the right marginal channel of Γ , for an arbitrary fixed state $\sigma \in M_{A_1 A_1}$. An analogous argument shows that $\Gamma^{A_1 A_1 \rightarrow A_2 A_2}$ is a well-defined left marginal QNS correlation for Γ , for an arbitrary fixed state $\rho \in M_{X_2 X_2}$.

Remark 6.14. Note that, in the event $\Gamma \in \mathcal{C}_{\text{st}}$ for $t \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}, \text{ns}\}$, then the marginal channels previously discussed (when considering Γ inside $\mathcal{Q}_{\text{st}}$) coincide with the marginal channels discussed in [20, Section 5].

Definition 6.15. An SQNS correlation Γ over $(X_2 X_2, A_1 A_1, X_1 X_1, A_2 A_2)$ is called jointly tracial if $\Gamma_{X_2 X_2 \rightarrow X_1 X_1}$ and $\Gamma^{A_1 A_1 \rightarrow A_2 A_2}$ are tracial QNS correlations.

Remark 6.16. There exists a surjective $*$ -homomorphism

$$\pi : \mathcal{C}_{X_2, A_2} \rightarrow \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \otimes_{\max} \mathcal{C}_{X_1, A_1}.$$

Indeed: for $x_2, x'_2 \in X_2, a_2, a'_2 \in A_2$ let

$$\tilde{e}_{x_2 x'_2, a_2 a'_2} := \sum_{x_1, x'_1} \sum_{a_1, a'_1} e_{x_2 x'_2, x_1 x'_1} \otimes e_{a_1 a'_1, a_2 a'_2} \otimes f_{x_1 x'_1, a_1 a'_1}.$$

One easily checks that $\tilde{e}_{x_2x_2',a_2a_2'} \geq 0$ and $\sum_{a_2} \tilde{e}_{x_2x_2',a_2a_2'} = \delta_{x_2x_2'} 1$ for all $x_2, x_2' \in X_2, a_2 \in A_2$. Therefore, by universality there exists a surjective $*$ -homomorphism $\pi : \mathcal{C}_{X_2,A_2} \rightarrow \mathcal{C}_{X_2,X_1} \otimes_{\max} \mathcal{C}_{A_1,A_2} \otimes_{\max} \mathcal{C}_{X_1,A_1}$ which sends $\pi(e_{x_2x_2',a_2a_2'}) = \tilde{e}_{x_2x_2',a_2a_2'}$, $x_2, x_2' \in X_2, a_2, a_2' \in A_2$.

For a linear functional Γ on $\mathcal{C}_{X_2,X_1} \otimes \mathcal{C}_{A_1,A_2}$, set

$$\Gamma_{\Gamma}(x_1x_1', y_1y_1', a_2a_2', b_2b_2' | x_2x_2', y_2y_2', a_1a_1', b_1b_1') = \\ \Gamma(e_{x_2x_2',x_1x_1'} e_{y_2y_2',y_1y_1'} \otimes e_{a_1a_1',a_2a_2'} e_{b_1b_1',b_2b_2'}),$$

where $x_i, x_i', y_i, y_i' \in X_i, a_i, a_i', b_i, b_i' \in A_i, i = 1, 2$.

Theorem 6.17. *The following hold:*

- (i) *If Γ is a jointly tracial and quantum commuting SQNS correlation, then there exists a trace Γ_{Γ} on $\mathcal{C}_{X_2,X_1} \otimes_{\max} \mathcal{C}_{A_1,A_2}$ such that $\Gamma = \Gamma_{\Gamma}$.*
- (ii) *If Γ is a jointly tracial and approximately quantum SQNS correlation, then there exists an amenable trace Γ_{Γ} on $\mathcal{C}_{X_2,X_1} \otimes_{\min} \mathcal{C}_{A_1,A_2}$ such that $\Gamma = \Gamma_{\Gamma}$.*
- (iii) *If Γ is a jointly tracial and quantum SQNS correlation, then there exists a trace Γ_{Γ} factoring through a finite-dimensional $*$ -representation of $\mathcal{C}_{X_2,X_1} \otimes_{\min} \mathcal{C}_{A_1,A_2}$ such that $\Gamma = \Gamma_{\Gamma}$.*
- (iv) *If Γ is a jointly tracial and local SQNS correlation, then there exists a trace Γ_{Γ} factoring through an abelian $*$ -representation of $\mathcal{C}_{X_2,X_1} \otimes_{\max} \mathcal{C}_{A_1,A_2}$ such that $\Gamma = \Gamma_{\Gamma}$.*

Proof. (i) First, assume $\Gamma \in \mathcal{Q}_{\text{sqc}}$ is jointly tracial. For notational simplification, write

$$\mathfrak{B} = \mathcal{C}_{X_2,X_1} \otimes_{\max} \mathcal{C}_{X_2,X_1} \otimes_{\max} \mathcal{C}_{A_1,A_2} \otimes_{\max} \mathcal{C}_{A_1,A_2}$$

and

$$\mathfrak{U} = \mathcal{C}_{X_2,X_1} \otimes_{\max} \mathcal{C}_{A_1,A_2}.$$

We note that, up to a flip of tensor legs, $\mathfrak{B} = \mathfrak{U} \otimes_{\max} \mathfrak{U}$; in the sequel, we will use this identification without explicitly mentioning it.

By (the proof of) Theorem 6.6, there exists a state $\tilde{s} : \mathfrak{B} \rightarrow \mathbb{C}$ such that $\Gamma = \Gamma_{\tilde{s}}$. For $v, w \in \mathfrak{U} \otimes_{\max} \mathfrak{U}$, write $v \sim w$ if $\tilde{s}(v - w) = 0$. Let $V = (v_{x_1,x_2})_{x_1,x_2}$ be the isometry such that $e_{x_2x_2',x_1x_1'} = v_{x_1,x_2}^* v_{x_1',x_2'}$. Then

$$VV^* = \left(\sum_{x_2} v_{x_1,x_2} v_{x_1',x_2}^* \right)_{x_1,x_1'}$$

is a projection, with $\sum_{x_2} v_{x_1,x_2} v_{x_1',x_2}^* \leq 1$ for all $x_1 \in X_1$. Using this, we see that

$$\begin{aligned} \sum_{x_2 \in X_2} e_{x_2'x_2,x_1'x_1'} e_{x_2x_2',x_1x_1'} &= \sum_{x_2 \in X_2} v_{x_1',x_2}^* v_{x_1,x_2} v_{x_1,x_2}^* v_{x_1',x_2'} \\ &\leq v_{x_1',x_2}^* v_{x_1,x_2} = e_{x_2'x_2,x_1'x_1'}. \end{aligned}$$

From this, we compute

$$\sum_{x_2, x'_2 \in X_2} \sum_{x_1, x'_1 \in X_1} e_{x'_2 x_2, x'_1 x_1} e_{x_2 x'_2, x_1 x'_1} \leq |X_2| |X_1| 1. \quad (28)$$

Let τ_X be the tracial state on $\mathcal{C}_{X_2, X_1}$ corresponding to the marginal channel $\Gamma_{X_2 X_2 \rightarrow X_1 X_1}$. Without loss of generality, fix $a_1, b_1 \in A_1$. We see

$$\begin{aligned} & \tilde{s}(e_{x_2 x'_2, x_1 x'_1} \otimes 1 \otimes 1 \otimes 1) \\ &= \sum_{y_1 \in X_1} \sum_{a_2, b_2 \in A_2} \tilde{s}(e_{x_2 x'_2, x_1 x'_1} \otimes e_{x_2 x_2, y_1 y_1} \otimes e_{a_1 a_1, a_2 a_2} \otimes e_{b_1 b_1, b_2 b_2}) \\ &= \sum_{a_2, b_2 \in A_2} \sum_{y_1 \in X_1} \tau_X(e_{x_2 x'_2, x_1 x'_1} e_{x_2 x_2, y_1 y_1}) \\ &= \tau_X(e_{x_2 x'_2, x_1 x'_1}). \end{aligned}$$

Similarly,

$$\begin{aligned} & \tilde{s}(1 \otimes e_{x'_2 x_2, x'_1 x_1} \otimes 1 \otimes 1) \\ &= \sum_{y_1 \in X_1} \sum_{a_2, b_2 \in A_2} \tilde{s}(e_{x'_2 x_2, y_1 y_1} \otimes e_{x'_2 x_2, x'_1 x_1} \otimes e_{a_1 a_1, a_2 a_2} \otimes e_{b_1 b_1, b_2 b_2}) \\ &= \sum_{a_2, b_2 \in A_2} \sum_{y_1 \in X_1} \tau_X(e_{x'_2 x_2, y_1 y_1} e_{x_2 x'_2, x_1 x'_1}) \\ &= \tau_X(e_{x_2 x'_2, x_1 x'_1}). \end{aligned}$$

Therefore,

$$e_{x_2 x'_2, x_1 x'_1} \otimes 1 \otimes 1 \otimes 1 \sim 1 \otimes e_{x'_2 x_2, x'_1 x_1} \otimes 1 \otimes 1, \quad x_i, x'_i \in X_i, i = 1, 2.$$

Write

$$h_{x_2 x'_2, x_1 x'_1} = e_{x_2 x'_2, x_1 x'_1} \otimes 1 \otimes 1 \otimes 1 - 1 \otimes e_{x'_2 x_2, x'_1 x_1} \otimes 1 \otimes 1,$$

for $x_i, x'_i \in X_i, i = 1, 2$; we note that

$$\begin{aligned} & h_{x_2 x'_2, x_1 x'_1}^* h_{x_2 x'_2, x_1 x'_1} \\ &= (e_{x'_2 x_2, x'_1 x_1} e_{x_2 x'_2, x_1 x'_1} \otimes 1 \otimes 1 \otimes 1 + 1 \otimes e_{x_2 x'_2, x_1 x'_1} e_{x'_2 x_2, x'_1 x_1} \otimes 1 \otimes 1) \\ &- (e_{x'_2 x_2, x'_1 x_1} \otimes e_{x_2 x'_2, x_1 x'_1} \otimes 1 \otimes 1 + e_{x_2 x'_2, x_1 x'_1} \otimes e_{x'_2 x_2, x'_1 x_1} \otimes 1 \otimes 1). \end{aligned}$$

By (28), we have

$$\begin{aligned} & \sum_{x_2, x'_2, x_1, x'_1} \tilde{s}(h_{x_2 x'_2, x_1 x'_1}^* h_{x_2 x'_2, x_1 x'_1}) \\ &= \sum_{x_2, x'_2, x_1, x'_1} \tilde{s}(e_{x'_2 x_2, x'_1 x_1} e_{x_2 x'_2, x_1 x'_1} \otimes 1 \otimes 1 \otimes 1 + 1 \otimes e_{x_2 x'_2, x_1 x'_1} e_{x'_2 x_2, x'_1 x_1} \otimes 1 \otimes 1) \\ &- \tilde{s}(e_{x'_2 x_2, x'_1 x_1} \otimes e_{x_2 x'_2, x_1 x'_1} \otimes 1 \otimes 1 + e_{x_2 x'_2, x_1 x'_1} \otimes e_{x'_2 x_2, x'_1 x_1} \otimes 1 \otimes 1) \\ &\leq 2|X_2| |X_1| 1 - 2|X_2| |X_1| 1 = 0. \end{aligned}$$

From this, we have

$$\tilde{s}(h_{x_2x'_2, x_1x'_1}^* h_{x_2x'_2, x_1x'_1}) = 0, \quad x_i, x'_i \in X_i, i = 1, 2. \quad (29)$$

By applying Cauchy-Schwarz inequality in conjunction with (29), we have $wh_{x_2x'_2, x_1x'_1} \sim 0$ and $h_{x_2x'_2, x_1x'_1}w \sim 0$ for all $w \in \mathfrak{B}$. In particular, for all $x_i, x'_i \in X_i, i = 1, 2$ we have

$$ze_{x_2x'_2, x_1x'_1} \otimes 1 \otimes v \sim z \otimes e_{x'_2x_2, x'_1x_1} \otimes v \sim e_{x_2x'_2, x_1x'_1} z \otimes 1 \otimes v, \quad (30)$$

for $z \in \mathcal{C}_{X_2, X_1}, v \in \mathfrak{U}$. If instead we use τ_A for the tracial state corresponding to marginal channel $\Gamma^{A_1A_1 \rightarrow A_2A_2}$, we can make an almost identical argument to show that

$$u \otimes ze_{a_1a'_1, a_2a'_2} \otimes 1 \sim u \otimes z \otimes e_{a'_1a_1, a'_2a_2} \sim u \otimes e_{a_1a'_1, a_2a'_2}, \quad (31)$$

for $a_i, a'_i \in A_i, z \in \mathcal{C}_{A_1, A_2}$ and $u \in \mathfrak{U}, i = 1, 2$. Equations (30) and (31) imply

$$ze_{x_2x'_2, x_1x'_1} \otimes 1 \otimes z'e_{a_1a'_1, a_2a'_2} \otimes 1 \sim z \otimes e_{x'_2x_2, x'_1x_1} \otimes z' \otimes e_{a'_1a_1, a'_2a_2} \quad (32)$$

$$\sim e_{x_2x'_2, x_1x'_1} z \otimes 1 \otimes e_{a_1a'_1, a_2a'_2} z' \otimes 1, \quad (33)$$

for all $z \in \mathcal{C}_{X_2, X_1}, z' \in \mathcal{C}_{A_1, A_2}$, and $x_i, x'_i \in X_i, a_i, a'_i \in A_i, i = 1, 2$. An induction argument on the lengths of the words w on $\{e_{x_2x'_2, x_1x'_1} : x_i, x'_i \in X_i, i = 1, 2\}$ and w' on $\{e_{a_1a'_1, a_2a'_2} : a_i, a'_i \in A_i, i = 1, 2\}$ whose base step is provided by (32) shows that

$$zw \otimes 1 \otimes z'w' \otimes 1 \sim wz \otimes 1 \otimes w'z' \otimes 1, \quad z, w \in \mathcal{C}_{X_2, X_1}, z', w' \in \mathcal{C}_{A_1, A_2}$$

(see, for instance, the proofs of [6, Theorem 3.2, Theorem 4.1]). We may then conclude that the functional T on $\mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2}$, given by

$$\mathsf{T}(u \otimes v) = \tilde{s}(u \otimes 1 \otimes v \otimes 1), \quad u \in \mathcal{C}_{X_2, X_1}, v \in \mathcal{C}_{A_1, A_2},$$

is a tracial state. Furthermore, (32) implies $\Gamma = \Gamma_{\mathsf{T}}$.

(ii) If Γ is an approximately quantum jointly tracial correlation, again by the proof of Theorem 6.6 there exists a state $s : \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{A_1, A_2} \otimes_{\min} \mathcal{C}_{A_1, A_2} \rightarrow \mathbb{C}$ such that $\Gamma = \Gamma_s$. As each approximately quantum SQNS correlation is quantum commuting, by (i) there exists some trace $\mathsf{T} : \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \rightarrow \mathbb{C}$ such that $\Gamma = \Gamma_{\mathsf{T}}$ as well. By [39, Lemma 9.2], for any finite sets X, Y there exists a $*$ -isomorphism $\partial : \mathcal{C}_{X, Y} \rightarrow \mathcal{C}_{X, Y}^{\text{op}}$ such that

$$\partial(e_{xx', yy'}) = e_{x'y', yx'}^{\text{op}}, \quad x, x' \in X, y, y' \in Y.$$

Let $\partial_X : \mathcal{C}_{X_2, X_1} \rightarrow \mathcal{C}_{X_2, X_1}^{\text{op}}$ and $\partial_A : \mathcal{C}_{A_1, A_2} \rightarrow \mathcal{C}_{A_1, A_2}^{\text{op}}$ be the $*$ -isomorphisms corresponding to each $\mathbb{C}^*$ -algebra. Let

$$\begin{aligned} \mathcal{F} : \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{A_1, A_2} \otimes_{\min} \mathcal{C}_{A_1, A_2} \\ \rightarrow \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{A_1, A_2} \otimes_{\min} \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{A_1, A_2} \end{aligned}$$

be the flip operation, and

$$q : \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \rightarrow \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{A_1, A_2}$$

be the quotient map. Define linear functional

$$\mu : (\mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2}) \otimes_{\min} (\mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2})^{\text{op}} \rightarrow \mathbb{C}$$

via

$$\mu = s \circ \mathcal{F} \circ (q \otimes (q \circ (\partial_X^{-1} \otimes \partial_A^{-1}))).$$

One may easily check that $\mu(u \otimes v^{\text{op}}) = T(uv)$. Indeed: on the canonical generators we have

$$\begin{aligned} & \mu(e_{x_2 x'_2, x_1 x'_1} \otimes e_{a_1 a'_1, a_2 a'_2} \otimes (e_{y'_2 y_2, y'_1 y_1} \otimes e_{b'_1 b_1, b'_2 b_2})^{\text{op}}) \\ &= s \circ \mathcal{F}(e_{x_2 x'_2, x_1 x'_1} \otimes e_{a_1 a'_1, a_2 a'_2} \otimes e_{y_2 y'_2, y_1 y'_1} \otimes e_{b_1 b'_1, b_2 b'_2}) \\ &= s(e_{x_2 x'_2, x_1 x'_1} \otimes e_{y_2 y'_2, y_1 y'_1} \otimes e_{a_1 a'_1, a_2 a'_2} \otimes e_{b_1 b'_1, b_2 b'_2}) \\ &= T(e_{x_2 x'_2, x_1 x'_1} e_{y'_2 y_2, y'_1 y_1} \otimes e_{a_1 a'_1, a_2 a'_2} e_{b'_1 b_1, b'_2 b_2}). \end{aligned}$$

By [9, Theorem 6.2.7], trace T is amenable.

(iii) Let Γ be a perfect concurrent strategy in $\mathcal{Q}_{\text{sq}}$. By the proof of Theorem 6.6 (iii), there exist finite-dimensional Hilbert spaces $H_X, H_{X'}, H_A, H_{A'}$, $*$ -representations $\pi_X : \mathcal{C}_{X_2, X_1} \rightarrow \mathcal{B}(H_X), \pi_{X'} : \mathcal{C}_{X_2, X_1} \rightarrow \mathcal{B}(H_{X'}), \pi_A : \mathcal{C}_{A_1, A_2} \rightarrow \mathcal{B}(H_A)$ and $\pi_{A'} : \mathcal{C}_{A_1, A_2} \rightarrow \mathcal{B}(H_{A'})$ along with a unit vector $\xi \in H_X \otimes H_{X'} \otimes H_A \otimes H_{A'}$ such that $\Gamma = \Gamma_{\tilde{s}}$, where

$$\tilde{s} : \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{A_1, A_2} \otimes_{\min} \mathcal{C}_{A_1, A_2} \rightarrow \mathbb{C}$$

is a state given by

$$\tilde{s}(u) = \langle (\pi_X \otimes \pi_{X'} \otimes \pi_A \otimes \pi_{A'})(u) \xi, \xi \rangle, \quad (34)$$

for $u \in \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{A_1, A_2} \otimes_{\min} \mathcal{C}_{A_1, A_2}$. By the proof of (i), the state $T : \mathcal{C}_{X_2, X_1} \otimes_{\min} \mathcal{C}_{A_1, A_2} \rightarrow \mathbb{C}$ constructed from $\tilde{s}$ is tracial, and in this case factors through a finite-dimensional $*$ -representation.

(iv) The proof of this statement is similar to the proof of Proposition 3.3; we include the details for the benefit of the reader. Assume $\Gamma \in \mathcal{Q}_{\text{sloc}}$ is a perfect concurrent strategy; by Remark 3.11, $\Gamma = \sum_{j=1}^k \lambda_j \Phi_X^{(j)} \otimes \Phi_{X'}^{(j)} \otimes \Phi_A^{(j)} \otimes \Phi_{A'}^{(j)}$ as a convex combination of quantum channels $\Phi_X^{(j)}, \Phi_{X'}^{(j)} : M_{X_2} \rightarrow M_{X_1}, \Phi_A^{(j)}, \Phi_{A'}^{(j)} : M_{A_1} \rightarrow M_{A_2}, j = 1, \dots, k$. Let

$$\begin{aligned} (\lambda_{x_2 x'_2, x_1 x'_1}^{(j)})_{x_1, x'_1} &= \Phi_X^{(j)}(\epsilon_{x_2 x'_2}), & (\lambda_{y_2 y'_2, y_1 y'_1}^{(j)})_{y_1, y'_1} &= \Phi_{X'}^{(j)}(\epsilon_{y_2 y'_2}), \\ (\mu_{a_1 a'_1, a_2 a'_2}^{(j)})_{a_2, a'_2} &= \Phi_A^{(j)}(\epsilon_{a_1 a'_1}), & (\mu_{b_1 b'_1, b_2 b'_2}^{(j)})_{b_2, b'_2} &= \Phi_{A'}^{(j)}(\epsilon_{b_1 b'_1}), \end{aligned}$$

for $x_2, x'_2, y_2, y'_2 \in X_2, a_1, a'_1, b_1, b'_1 \in A_1$, and $\pi_X^{(j)}, \pi_{X'}^{(j)} : \mathcal{C}_{X_2, X_1} \rightarrow \mathbb{C}, \pi_A^{(j)}, \pi_{A'}^{(j)} : \mathcal{C}_{A_1, A_2} \rightarrow \mathbb{C}$ be the $*$ -representations given by

$$\begin{aligned} \pi_X^{(j)}(e_{x_2 x'_2, x_1 x'_1}) &= \lambda_{x_2 x'_2, x_1 x'_1}^{(j)}, & \pi_{X'}^{(j)}(e_{y_2 y'_2, y_1 y'_1}) &= \lambda_{y_2 y'_2, y_1 y'_1}^{(j)}, \\ \pi_A^{(j)}(e_{a_1 a'_1, a_2 a'_2}) &= \mu_{a_1 a'_1, a_2 a'_2}^{(j)}, & \pi_{A'}^{(j)}(e_{b_1 b'_1, b_2 b'_2}) &= \mu_{b_1 b'_1, b_2 b'_2}^{(j)} \end{aligned}$$

for $j = 1, \dots, k$. Furthermore, let $\pi_X, \pi_{X'} : \mathcal{C}_{X_2, X_1} \rightarrow \mathcal{B}(\mathbb{C}^k)$, $\pi_A, \pi_{A'} : \mathcal{C}_{A_1, A_2} \rightarrow \mathcal{B}(\mathbb{C}^k)$ be the $*$ -representations given by

$$\begin{aligned}\pi_X(u) &= \sum_{j=1}^k \pi_X^{(j)}(u) \epsilon_{jj}, & \pi_{X'}(u) &= \sum_{j=1}^k \pi_{X'}^{(j)}(u) \epsilon_{jj}, \\ \pi_A(v) &= \sum_{j=1}^k \pi_A^{(j)}(v) \epsilon_{jj}, & \pi_{A'}(v) &= \sum_{j=1}^k \pi_{A'}^{(j)}(v) \epsilon_{jj},\end{aligned}$$

for $u \in \mathcal{C}_{X_2, X_1}, v \in \mathcal{C}_{A_1, A_2}$. Clearly, the images of $\pi_X, \pi_{X'}, \pi_A, \pi_{A'}$ are all abelian.

If we then set $\xi = \sum_{j=1}^k \sqrt{\lambda_j} e_j \otimes e_j \otimes e_j \otimes e_j \in \mathbb{C}^k \otimes \mathbb{C}^k \otimes \mathbb{C}^k \otimes \mathbb{C}^k$, we have

$$\begin{aligned}& \Gamma(\epsilon_{x_2 x'_2} \otimes \epsilon_{y_2 y'_2} \otimes \epsilon_{a_1 a'_1} \otimes \epsilon_{b_1 b'_1}) \\ &= \left(\langle (\pi_X(e_{x_2 x'_2, x_1 x'_1}) \otimes \pi_{X'}(e_{y_2 y'_2, y_1 y'_1}) \otimes \pi_A(e_{a_1 a'_1, a_2 a'_2}) \otimes \pi_{A'}(e_{b_1 b'_1, b_2 b'_2})) \xi, \xi \rangle \right)_{x_1 x'_1, y_1 y'_1}^{a_2 a'_2, b_2 b'_2},\end{aligned}$$

with corresponding state $\tilde{s}$ given by

$$\tilde{s}(u) = \langle (\pi_X \otimes \pi_{X'} \otimes \pi_A \otimes \pi_{A'})(u) \xi, \xi \rangle,$$

for $u \in \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \otimes_{\max} \mathcal{C}_{A_1, A_2}$. Argue now as in (iii) to conclude that tracial state Γ factors through an abelian $*$ -representation of $\mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2}$. $\square$

Theorem 6.18. *Let $t \in \{\text{loc}, q, \text{qa}, \text{qc}\}$. If $\Gamma \in \mathcal{Q}_{\text{st}}$ is jointly tracial, and $\mathcal{E} \in \mathcal{Q}_t$ is tracial, then $\Gamma[\mathcal{E}]$ is tracial.*

Proof. We handle the case when $t = \text{qc}$. By Theorem 6.17 and [6, Theorem 4.1], there exists tracial states $\Gamma : \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \rightarrow \mathbb{C}$ and $\tau : \mathcal{C}_{X_1, A_1} \rightarrow \mathbb{C}$ such that $\Gamma = \Gamma_\top$ and $\mathcal{E} = \mathcal{E}_\tau$. Let

$$\Gamma \odot \tau : \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \otimes_{\max} \mathcal{C}_{X_1, A_1} \rightarrow \mathbb{C}$$

be the linear map given by

$$(\Gamma \odot \tau)(u \otimes v \otimes w) = \Gamma(u \otimes v) \tau(w), \quad u \in \mathcal{C}_{X_2, X_1}, v \in \mathcal{C}_{A_1, A_2}, w \in \mathcal{C}_{X_1, A_1}.$$

That this is a state follows from [38, Proposition 4.23]; that it is also tracial follows from a standard argument using the (automatic) continuity of the mapping on the maximal tensor product, in conjunction with calculations on the dense subset of finite linear combinations of simple tensors. Define tracial state $\tilde{\tau}$ on $\mathcal{C}_{X_2, A_2}$ in the following way: by Remark 6.16, there exists a surjective $*$ -representation $\pi : \mathcal{C}_{X_2, A_2} \rightarrow \mathcal{C}_{X_2, X_1} \otimes_{\max} \mathcal{C}_{A_1, A_2} \otimes_{\max} \mathcal{C}_{X_1, A_1}$ sending $\pi(e_{x_2 x'_2, a_2 a'_2}) = \tilde{e}_{x_2 x'_2, a_2 a'_2}$ for $x_2, x'_2 \in X_2, a_2, a'_2 \in A_2$. Using π , we let

$$\tilde{\tau}(u) := ((\Gamma \odot \tau) \circ \pi)(u), \quad u \in \mathcal{C}_{X_2, A_2}. \quad (35)$$

We note

$$\begin{aligned}
& \Gamma[\mathcal{E}](a_2 a'_2, b_2 b'_2 | x_2 x'_2, y_2 y'_2) \\
&= \sum_{\substack{x_1, x'_1 \\ y_1, y'_1}} \sum_{\substack{a_1, a'_1 \\ b_1, b'_1}} \Gamma(x_1 x'_1, y_1 y'_1, a_2 a'_2, b_2 b'_2 | x_2 x'_2, y_2 y'_2, a_1 a'_1, b_1 b'_1) \mathcal{E}(a_1 a'_1, b_1 b'_1 | x_1 x'_1, y_1 y'_1) \\
&= \sum_{\substack{x_1, x'_1 \\ y_1, y'_1}} \sum_{\substack{a_1, a'_1 \\ b_1, b'_1}} (\tau \odot \tau)(e_{x_2 x'_2, x_1 x'_1} e_{y_2 y'_2, y_1 y'_1} \otimes e_{a_1 a'_1, a_2 a'_2} e_{b_1 b'_1, b_2 b'_2} \otimes f_{x_1 x'_1, a_1 a'_1} f_{y_1 y'_1, b_1 b'_1}) \\
&= (\tau \odot \tau)(\pi(e_{x_2 x'_2, a_2 a'_2} e_{y_2 y'_2, b_2 b'_2})) \\
&= \tilde{\tau}(e_{x_2 x'_2, a_2 a'_2} e_{y_2 y'_2, b_2 b'_2}),
\end{aligned}$$

for $x_2, x'_2, y_2, y'_2 \in X_2, a_2, a'_2, b_2, b'_2 \in A_2$. This shows $\Gamma[\mathcal{E}] = \Gamma_{\tilde{\tau}}$, and so by [6, Theorem 4.10] once more we conclude that $\Gamma[\mathcal{E}]$ is tracial. For $t \in \{\text{loc}, \text{q}, \text{qa}\}$, the argument is similar once we use Theorem 6.17 (ii)-(iv). $\square$

For finite sets X, A , let

$$E_{XA} = (e_{x, x', a, a'})_{x, x', a, a'}, \quad E_{XA}^{\text{op}} = (e_{x', x, a', a})_{x, x', a, a'},$$

be considered as elements in $M_{XA} \otimes \mathfrak{G}_{X,A}$ and $M_{XA} \otimes \mathfrak{G}_{X,A}^{\text{op}}$, respectively. If $P \in \mathcal{P}_{XX}, Q \in \mathcal{P}_{AA}$, define a linear map

$$\gamma_{P,Q} : M_{XX} \otimes M_{AA} \otimes \mathfrak{G}_{X,A} \otimes \mathfrak{G}_{X,A}^{\text{op}} \rightarrow \mathfrak{G}_{X,A} \quad (36)$$

by setting

$$\gamma_{P,Q}(\omega \otimes u \otimes v^{\text{op}}) = \text{Tr}(\omega(P \otimes Q))uv, \quad \omega \in M_{XX} \otimes M_{AA}, u, v \in \mathfrak{G}_{X,A}.$$

If $\varphi : \mathcal{P}_{XX} \rightarrow \mathcal{P}_{AA}$ is any quantum non-local game, let

$$\mathfrak{J}(\varphi) = \left\langle \gamma_{P, \varphi(P)}(E_{XA} \otimes E_{XA}^{\text{op}}) : P \in \mathcal{P}_{XX} \right\rangle$$

be the generated $*$ -ideal in $\mathfrak{G}_{X,A}$, and $J(\varphi)$ be the corresponding ideal considered in $\mathcal{C}_{X,A}$. Finally, write $\mathfrak{C}(\varphi) = \mathfrak{G}_{X,A} / \mathfrak{J}(\varphi)$ (resp. $\mathcal{C}(\varphi) = \mathcal{C}_{X,A} / J(\varphi)$) for the quotient $*$ -algebra (resp. quotient C^* -algebra). Perfect strategies for a concurrent quantum game φ were shown to correspond to tracial states acting on $*$ -representations of $\mathcal{C}_{X,A}$ which annihilate $\mathfrak{J}(\varphi)$ or $J(\varphi)$ in [6, 7]. In light of previous results, we may conclude by giving an algebraic characterization of our transfer of perfect strategies between quantum games.

Theorem 6.19. *Let $X_i, A_i, i = 1, 2$ be finite sets, $P_i \in M_{X_i X_i}, Q_i \in M_{A_i A_i}, i = 1, 2$ be projections, and $t \in \{\text{loc}, \text{q}, \text{qa}, \text{qc}\}$. If $\overline{\mathcal{U}}_{P_1, Q_1} \rightarrow_{\text{st}} \mathcal{U}_{P_2, Q_2}$ via jointly tracial Γ_T , then for any tracial state τ on $\mathcal{C}(\varphi_{P_1 \rightarrow Q_1})$, the tracial state $\tilde{\tau}$ given in (35) restricts to a tracial state on $\mathcal{C}(\varphi_{P_2 \rightarrow Q_2})$.*

Proof. As corresponding QNS correlation $\mathcal{E}_\tau$ is both tracial and a perfect strategy for $\varphi_{P_1 \rightarrow Q_1}$ (via [6, Corollary 4.4]), by Theorem 6.18 we know $\Gamma_\tau[\mathcal{E}_\tau]$ is tracial, and corresponds to tracial state $\tilde{\tau}$ given in (35). Furthermore, by Theorem 5.3 $\Gamma_\tau[\mathcal{E}_\tau]$ is a perfect strategy for $\varphi_{P_2 \rightarrow Q_2}$; thus, $\tilde{\tau}$ annihilates $J(\varphi_{P_2 \rightarrow Q_2})$ as claimed. $\square$

Remark 6.20. We wish to comment on the specific case when both games are the quantum graph isomorphism game. Let $X_i = A_i, i = 1, 2$, let $\mathcal{U}_i \subseteq \mathbb{C}^{X_i} \otimes \mathbb{C}^{X_i}, i = 1, 2$ be quantum graphs, and set $P_i := P_{\mathcal{U}_i}, i = 1, 2$.

Recall, that, for finite set X , the C^* -algebra of the free unitary quantum group $C(\mathcal{U}_X^+)$ is the universal unital C^* -algebra generated by the entries $u_{x,a}$ of an $X \times X$ matrix $U = (u_{x,a})_{x,a \in X}$ under the condition that U and U^t are unitary (known otherwise as a bi-unitary matrix). Subalgebra $C(\mathbb{P}\mathcal{U}_X^+)$ is generated by length two words of the form $u_{x,a}^* u_{x',a'}, x, x', a, a' \in X$ - and was shown in [1, Corollary 4.1] to be the proper quantization of the automorphism group of M_X . In [7, Theorem 6.7], concurrent bicorrelations of types $t \in \{\text{loc}, \text{q}, \text{qc}\}$ (and therefore, strategies for the quantum graph isomorphism game) were shown to be in correspondence with different tracial states on $C(\mathbb{P}\mathcal{U}_X^+)$.

Abusing notation, for $S, T \in M_{XX}$ let

$$\gamma_{S,T} : M_{XX} \otimes C(\mathbb{P}\mathcal{U}_X^+) \otimes M_{XX} \otimes C(\mathbb{P}\mathcal{U}_X^+)^{\text{op}} \rightarrow C(\mathbb{P}\mathcal{U}_X^+)$$

be defined as in (36). We let $\tilde{U} = (u_{x,x',a,a'})_{x,x',a,a'} \in M_{XX}(C(\mathbb{P}\mathcal{U}_X^+))$, and for $P \in \text{Proj}(M_{XX})$ set

$$J_{P,P} := \left\langle \gamma_{P,P^\perp}(\tilde{U} \otimes \tilde{U}^{\text{op}}), \gamma_{P^\perp,P}(\tilde{U} \otimes \tilde{U}^{\text{op}}) \right\rangle.$$

Finally, set $\mathcal{A}_{P,P} := C(\mathbb{P}\mathcal{U}_X^+)/J_{P,P}$. In [7, Remark 7.12], it was shown that C^* -algebra $\mathcal{A}_{P,P}$ can be endowed with a natural co-associative comultiplication $\Delta_P : \mathcal{A}_{P,P} \rightarrow \mathcal{A}_{P,P} \otimes \mathcal{A}_{P,P}$; this means $\mathcal{A}_{P,P}$ can be viewed as a compact quantum group. By [7, Theorem 7.10] and Theorem 5.7, we see that the existence of a quantum hypergraph isomorphism Γ between $\overline{\mathcal{U}}_{P_1,P_1}$ and $\mathcal{U}_{P_2,P_2}$ ensures a way to construct $*$ -representations of compact quantum group $\mathcal{A}_{P_2,P_2}$ from $*$ -reps of $\mathcal{A}_{P_1,P_1}$; that is, a way to transfer quantum automorphisms between quantum graphs.

References

- [1] BANICA, T., Symmetries of a generic coaction, *Math. Ann.* **314** (1999), no. 4, 763-780. [MR1709109](#) (2001g:46146), [Zbl 0928.46038](#), doi: [10.1007/s002080050315](#). 1653
- [2] BELL, J. S., On the Einstein Podolsky Rosen paradox, *Phys. Phys. Fiz.* **1** (1964), no. 3, 195-200. [MRMR3790629](#), [Zbl 0152.23605](#), doi: [10.1103/PhysicsPhysiqueFizika.1.195](#). 1608
- [3] BLECHER, D.; LE-MERDY, C., Operator algebras and their modules— an operator space approach. , *The Clarendon Press, Oxford University Press, Oxford*, 2004. x+387pp. ISBN: 0-19-852659-8 [MR2111973](#) (2006a:46070), [Zbl 1061.47002](#), doi: [10.1093/acprof:oso/9780198526599.001.0001](#). 1635
- [4] BOCHNIAK, A.; KASPRZAK, P.; SOŁTAN, P. M., Quantum correlations on quantum spaces, *Int. Math. Res. Not. IMRN* (2023), no.14, 12400-12440. [MR4615234](#), [Zbl 1520.58002](#), doi: [10.1093/imrn/rnac139](#). 1609

- [5] BRANNAN, M.; HAMIDI, M.; ISMERT, L.; NELSON, B.; WASILEWSKI, M., Quantum edge correspondences and quantum Cuntz-Krieger algebras, *J. London Math. Soc. (2)* **107** (2023), no. 3, 886-913. [MR4555986](#), [Zbl 1531.46031](#), doi: [10.1112/jlms.12702](#). [1633](#)
- [6] BRANNAN, M.; HARRIS, S.J.; TODOROV, I.G.; TUROWSKA, L., Synchronicity for quantum non-local games, *J. Funct. Anal.* **284** (2023), no. 2, Paper No. 109738, 54 pp. [MR4507619](#), [Zbl 1515.81117](#), doi: [10.1016/j.jfa.2022.109738](#). [1609](#), [1610](#), [1633](#), [1645](#), [1646](#), [1649](#), [1651](#), [1652](#), [1653](#)
- [7] BRANNAN, M.; HARRIS, S.J.; TODOROV, I.G.; TUROWSKA, L., Quantum no-signalling bicorrelations, *Adv. Math.* **449** (2024), Paper No. 109732, 81 pp. [MR4752739](#), [Zbl 1553.46073](#), doi: [10.1016/j.aim.2024.109732](#). [1609](#), [1610](#), [1612](#), [1613](#), [1617](#), [1624](#), [1633](#), [1635](#), [1636](#), [1642](#), [1645](#), [1652](#), [1653](#)
- [8] BRASSARD, G.; CLEVE, R.; TAPP, A., Cost of exactly simulating quantum entanglement with classical communication, *Phys. Rev. Lett.* **83** (1999). doi: [10.1103/PhysRevLett.83.1874](#). [1608](#)
- [9] BROWN, N. P.; OZAWA, N., *C*-algebras and finite-dimensional approximations*. Graduate Studies in Mathematics, 88. American Mathematical Society, Providence, RI, 2008. xvi+509 pp. ISBN: 978-0-8218-4381-9; 0-8218-4381-8 [MR2391387](#) (2009h:46101), [Zbl 1160.46001](#), doi: [10.1090/gsm/088](#). [1650](#)
- [10] CHOI, M. D., Completely positive linear maps on complex matrices, *Linear Algebra Appl.* **10** (1975), 285-290. [MR376726](#) (51 #12901), [Zbl 0327.15018](#), doi: [10.1016/0024-3795\(75\)90075-0](#).
- [11] COONEY, T.; JUNGE, M.; PALAZUELOS, C.; PÉREZ-GARCÍA, D., Rank-one quantum games, *Comput. Complexity* **24** (2015), no. 1, 133-196. [MR3320304](#), [Zbl 1328.81063](#), doi: [10.1007/s00037-014-0096-x](#). [1609](#)
- [12] CUBITT, T. S.; LEUNG, D.; MATTHEWS, W.; WINTER, A., Zero-error capacity and simulation assisted by non-local correlations, *IEEE Trans. Inform. Theory* **57** (2011), no. 8, 5509-5523. [MR2849372](#) (2012e:81041), [Zbl 1365.81024](#), doi: [10.1109/TIT.2011.2159047](#). [1614](#)
- [13] DAWS, M., Quantum graphs: different perspectives, homomorphisms, and quantum automorphisms, *Commun. Am. Math. Soc.* **4** (2024), 117-181. [MR4706978](#), [Zbl 1554.46049](#), doi: [10.1090/cams/30](#). [1617](#), [1633](#)
- [14] DOUGLAS, R. G., On majorization, factorization, and range inclusion of operators on Hilbert space, *Proc. Am. Math. Soc.* **17** (1966), 413-415. [MR203464](#) (34 #3315), [Zbl 0146.12503](#), doi: [10.2307/2035178](#).
- [15] DUAN, R.; WINTER, A., No-signalling assisted zero-error capacity of quantum channels and an information theoretic interpretation of the Lovász number, *IEEE Trans. Inform. Theory* **62** (2016), no. 2, 891-914. [MR3455905](#), [Zbl 1359.81062](#), doi: [10.1109/TIT.2015.2507979](#). [1609](#), [1612](#), [1614](#), [1625](#), [1633](#)
- [16] DYKEMA, K.; PAULSEN, V. I.; PRAKASH, J., Non-closure of the set of quantum correlations via graphs, *Comm. Math. Phys.* **365** (2019), no. 3, 1125-1142. [MR3916992](#), [Zbl 1408.81008](#), doi: [10.1007/s00220-019-03301-1](#). [1608](#)
- [17] FARENICK, D.; PAULSEN, V. I., Operator system quotients of matrix algebras and their tensor products, *Math. Scand.* **111** (2012), no. 2, 210-243. [MR3023524](#), [Zbl 1273.46038](#), doi: [10.7146/math.scand.a-15225](#). [1640](#), [1642](#)
- [18] FRITZ, T., Tsirelson's problem and Kirchberg's conjecture, *Rev. Math. Phys.* **24** (2012), no. 5, 1250012, 67 pp. [MR2928100](#), [Zbl 1250.81023](#), doi: [10.1142/S0129055X12500122](#). [1608](#)
- [19] HELTON, J. W.; MEYER, K. P.; PAULSEN, V. I.; SATRIANO, M., Algebras, synchronous games, and chromatic numbers of graphs, *New York J. Math.* **25** (2019), 328-361. [MR3933765](#), [Zbl 1410.05069](#). [1610](#)
- [20] HOEFER, G.; TODOROV, I. G., Quantum hypergraph homomorphisms and non-local games, *Dissertationes Math.* **588** (2023), 1-54. [MR4698861](#), [Zbl 1539.81030](#),

- doi: [10.4064/dm230309-14-11](https://doi.org/10.4064/dm230309-14-11). [1609](#), [1610](#), [1611](#), [1614](#), [1618](#), [1622](#), [1623](#), [1624](#), [1628](#), [1634](#), [1636](#), [1637](#), [1639](#), [1642](#), [1643](#), [1644](#), [1646](#)
- [21] HOEFER, G.; TODOROV, I. G., Homomorphisms of quantum hypergraphs, *J. Math. Anal. Appl.* **543** (2025), no. 2 (pt. 1), Paper No. 128907, 20. [MR4803784](#), [Zbl 1564.46057](#), doi: [10.1016/j.jmaa.2024.128907](https://doi.org/10.1016/j.jmaa.2024.128907). [1607](#), [1609](#), [1610](#), [1614](#), [1629](#), [1630](#), [1631](#), [1632](#), [1634](#)
 - [22] JI, Z.; NATARAJAN, A.; VIDICK, T.; WRIGHT, J.; YUEN, H., $MIP^*=RE$, *preprint* (2020), *arXiv:2001.04383*. [Zbl 1503.68075](#). [1608](#)
 - [23] JUNGE, M.; NAVASCUES, M.; PALAZUELOS, C.; PEREZ-GARCIA, D.; SCHOLZ, V. B.; WERNER, R. F., Connes' embedding problem and Tsirelson's problem, *J. Math. Phys.* **52** (2011), no. 1, 012102, 12 pp. [MR2790067](#) (2011m:46099), [Zbl 1314.81038](#), doi: [10.1063/1.3514538](https://doi.org/10.1063/1.3514538). [1608](#)
 - [24] KADISON, R. V.; RINGROSE, J. R., Fundamentals of the theory of operator algebras. Vol. I. Elementary theory. Graduate Studies in Mathematics, 16. *American Mathematical Society, Providence, RI*, 1997. pp. i-xxii and 399-1074. ISBN: 0-8218-0820-6 [MR1468230](#) (98f:46001a), [Zbl 0831.46060](#), doi: [10.1090/gsm/015](https://doi.org/10.1090/gsm/015). [1642](#)
 - [25] KAVRUK, A. S., Nuclearity related properties in operator systems, *J. Operator Theory* **71** (2014), no. 1, 95-156. [MR3173055](#), [Zbl 1349.46060](#), doi: [10.7900/jot.2011nov16.1977](https://doi.org/10.7900/jot.2011nov16.1977). [1634](#)
 - [26] KAVRUK, A. S., Tensor products of operator systems and applications. Thesis (Ph.D.)-University of Houston. *ProQuest LLC, Ann Arbor, MI*, 2011. 87 pp. ISBN: 978-1267-11028-2. [MR2992647](#). [1642](#)
 - [27] KAVRUK, A. S.; PAULSEN, V. I.; TODOROV, I. G.; TOMFORDE, M., Tensor products of operator systems, *J. Funct. Anal.* **261** (2011), no. 2, 267-299. [MR2793115](#) (2012h:46094), [Zbl 1235.46051](#), doi: [10.1016/j.jfa.2011.03.014](https://doi.org/10.1016/j.jfa.2011.03.014). [1634](#), [1636](#), [1637](#)
 - [28] KIRCHBERG, E.; WASSERMANN, S., C^* -algebras generated by operator systems, *J. Funct. Anal.* **155** (1998), no. 2, 324-351. [MR1624549](#) (99f:46085), [Zbl 0940.46038](#), doi: [10.1006/jfan.1997.3226](https://doi.org/10.1006/jfan.1997.3226). [1637](#)
 - [29] LUPINI, M.; MANČINSKA, L.; PAULSEN, V. I.; ROBERSON, D. E.; SCARPA, G.; SEVERINI, S.; TODOROV, I. G.; WINTER, A., Perfect strategies for non-local games, *Math. Phys. Anal. Geom.* **23** (2020), no. 1, Paper No. 7, 31 pp. [MR4069856](#), [Zbl 1436.91021](#), doi: [10.1007/s11040-020-9331-7](https://doi.org/10.1007/s11040-020-9331-7). [1610](#)
 - [30] MANČINSKA, L.; ROBERSON, D. E., Graph homomorphisms for quantum players, *9th Conference on the Theory of Quantum Computation, Communication, and Cryptography*, 212-216, LIPIcs. Leibniz Int. Proc. Inform. 27 *Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern*, 2014. [MR3354695](#), [Zbl 1360.91048](#).
 - [31] OZAWA, N., About the Connes' embedding conjecture: algebraic approaches, *Jpn. J. Math.* **8** (2013), no. 1, 147-183. [MR3067294](#), [Zbl 1278.46053](#), doi: [10.1007/s11537-013-1280-5](https://doi.org/10.1007/s11537-013-1280-5). [1608](#)
 - [32] PAULSEN, V. I., Completely bounded maps and operator algebras. Cambridge Studies in Advanced Mathematics, 78. *Cambridge University Press, Cambridge*, 2002. xii+300 pp. ISBN: 0-521-81669-6 [MR1976867](#) (2004c:46118), [Zbl 1029.47003](#), doi: [10.1017/CBO9780511546631](https://doi.org/10.1017/CBO9780511546631). [1612](#)
 - [33] PAULSEN, V. I.; TODOROV, I. G.; TOMFORDE, M., Operator system structures on ordered spaces, *Proc. Lond. Math. Soc.* (3) **102** (2011), no. 1, 25-49. [MR2747723](#) (2012h:47151), [Zbl 1213.46050](#), doi: [10.1112/plms/pdq011](https://doi.org/10.1112/plms/pdq011). [1638](#), [1639](#)
 - [34] REGEV, O.; VIDICK, T., Quantum XOR games, *ACM Trans. Comput. Theory* **7** (2015), no. 4, Art. 15, 43. [MR3414675](#), [Zbl 1348.81177](#), doi: [10.1145/2799560](https://doi.org/10.1145/2799560). [1609](#)
 - [35] SLOFSTRA, W., The set of quantum correlations is not closed, *Forum Math. Pi* **7** (2019), e1, 41. [MR3898717](#), [Zbl 1405.81021](#), doi: [10.1017/fmp.2018.3](https://doi.org/10.1017/fmp.2018.3). [1608](#)
 - [36] SLOFSTRA, W., Tsirelson's problem and an embedding theorem for groups arising from non-local games, *J. Amer. Math. Soc.* **33** (2020), no. 1, 1-56. [MR4066471](#), [Zbl 1480.20083](#), doi: [10.1090/jams/929](https://doi.org/10.1090/jams/929).

- [37] STAHLKE, D., Quantum zero-error source-channel coding and non-commutative graph theory, *IEEE Trans. Inform. Theory* **62** (2016), no. 1, 554-577. [MR3447998](#), [Zbl 1359.94454](#), doi: [10.1109/TIT.2015.2496377](#). [1633](#)
- [38] TAKESAKI, M., Theory of operator algebras. I., *Springer-Verlag, New York-Heidelberg*, 1979. vii+415pp. ISBN: 0-387-90391-7 [MR1873025](#) (2002m:46083), [Zbl 0990.46034](#), doi: [10.1007/978-1-4612-6188-9](#). [1651](#)
- [39] TODOROV, I. G.; TUROWSKA, L., Quantum no-signalling correlations and non-local games, *Comm. Math. Phys.* **405** (2024), no. 6, Paper No. 141, 65 pp. [MR4751723](#), [Zbl 07868800](#), doi: [10.1007/s00220-024-05001-x](#). [1609](#), [1610](#), [1612](#), [1613](#), [1615](#), [1617](#), [1623](#), [1626](#), [1630](#), [1632](#), [1633](#), [1634](#), [1635](#), [1636](#), [1638](#), [1640](#), [1641](#), [1642](#), [1643](#), [1644](#), [1645](#), [1646](#), [1649](#)
- [40] WATROUS, J., The theory of quantum information, *Cambridge University Press, Cambridge*, (2018), viii+590 pp. ISBN: 978-1-107-18056-7. [Zbl 1393.81004](#), doi: [10.1017/9781316848142](#).
- [41] ZETTL, H., A characterization of ternary rings of operators, *Adv. in Math.* **48** (1983), no. 2, 117-143. [MR700979](#) (84h:46093), [Zbl 0517.46049](#), doi: [10.1016/0001-8708\(83\)90083-X](#). [1635](#)

(Gage Hofer) DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755, USA

gage.hoefer@dartmouth.edu

This paper is available via <http://nyjm.albany.edu/j/2025/31-64.html>.